

LIFETIME DETERMINATION  
OF EXCITED STATES IN  $^{106}\text{Cd}^*$

S.F. ASHLEY<sup>a</sup>, A. LINNEMANN<sup>b</sup>, J. JOLIE<sup>b</sup>, P.H. REGAN<sup>a</sup>  
K. ANDGREN<sup>a,c</sup>, A. DEWALD<sup>b</sup>, E.A. MCCUTCHAN<sup>d</sup>, B. MELON<sup>b</sup>  
O. MÖLLER<sup>b</sup>, N.V. ZAMFIR<sup>d,e</sup>, L. AMON<sup>d,f</sup>, N. BOELAERT<sup>b,g</sup>  
R.B. CAKIRLI<sup>d,f</sup>, R.F. CASTEN<sup>d</sup>, R.M. CLARK<sup>h</sup>, C. FRANSEN<sup>b</sup>  
W. GELLETLY<sup>a</sup>, G. GÜRDAL<sup>d,i</sup>, M. HEIDEMANN<sup>b</sup>, K.L. KEYES<sup>j</sup>  
M.N-. ERDURAN<sup>f</sup>, D.A. MEYER<sup>d</sup>, A. PAPENBERG<sup>j</sup>, C. PLETTNER<sup>d</sup>  
G. RAINOVSKI<sup>k</sup>, R.V. RIBAS<sup>l</sup>, N.J. THOMAS<sup>a,c</sup>, J. VINSON<sup>d</sup>  
D.D. WARNER<sup>m</sup>, V. WERNER<sup>d</sup>, E. WILLIAMS<sup>d</sup>, K.O. ZELL<sup>b</sup>

<sup>a</sup>Department of Physics, University of Surrey, Guildford GU2 7XH, UK

<sup>b</sup>Institut für Kernphysik der Universität zu Köln, 50937 Köln, Germany

<sup>c</sup>Department of Physics, Royal Institute of Technology, Stockholm, Sweden

<sup>d</sup>WNSL, Yale University, New Haven, CT 06520, USA

<sup>e</sup>Institutul Național de Fizică și Inginerie Nucleară, București, Romania

<sup>f</sup>Department of Physics, Istanbul University, Istanbul, Turkey

<sup>g</sup>Universiteit Gent, Vakgroep Subatomaire en Stralingsfysica, Gent, Belgium

<sup>h</sup>Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

<sup>i</sup>Clark University, Worcester, MA 01610-1477, USA

<sup>j</sup>Institute of Physical Research, University of Paisley, Paisley PA1 2BE, UK

<sup>k</sup>Department of Physics and Astronomy, SUNY, Stony Brook, NY 11794, USA

<sup>l</sup>Instituto de Física, Universidade de São Paulo, C.P. 05315-970, Brazil

<sup>m</sup>CCLRC, Daresbury Laboratory, Daresbury, Warrington WA4 4AD, UK

*(Received November 11, 2006)*

Two separate experiments using the Differential Decay Curve Method have been performed to extract mean lifetimes of excited states in  $^{106}\text{Cd}$ . The medium-spin states of interest were populated by the  $^{98}\text{Mo}(^{12}\text{C}, 4n)^{106}\text{Cd}$  reaction performed at the Wright Nuclear Structure Lab., Yale University. From this experiment, two isomeric state mean lifetimes have been deduced. The low-lying states were populated by the  $^{96}\text{Mo}(^{13}\text{C}, 3n)^{106}\text{Cd}$  reaction performed at the Institut für Kernphysik, Universität zu Köln. The mean lifetime of the  $I^\pi = 2_1^+$  state was deduced, tentatively, as 16.4(9) ps. This value differs from the previously accepted literature value from Coulomb excitation of 10.43(9) ps.

PACS numbers: 21.10.Tg, 23.20.Lv, 25.70.Gh, 27.60.+j

---

\* Presented at the Zakopane Conference on Nuclear Physics, September 4–10, 2006, Zakopane, Poland.

## 1. Introduction

In terms of low-lying excitations, the cadmium nuclei are considered some of the best examples of quasi-vibrational nuclei (see reference [1] and references therein). However, from the systematics of the  $B(E2)$  values of the  $I^\pi = 2_1^+ \rightarrow 0_1^+$  and  $I^\pi = 4_1^+ \rightarrow 2_1^+$  transitions in  $^{104-110}\text{Cd}$  [2], the  $B(E2)$  values in  $^{106}\text{Cd}$  appear to be larger than the systematic trend of the light cadmium isotopes, whose  $B(E2)$  values decrease, approaching  $^{102}\text{Cd}$  [3]. Within the medium-spin regime, it is evident that there are collective structures with occupation of at least one  $\nu h_{11/2}$  orbital [4].

This paper summarises two experiments using the Recoil Distance Method (RDM) and Differential Decay Curve Method to determine  $B(E2)$  values for various transitions in  $^{106}\text{Cd}$ .

## 2. Experimental details

### 2.1. DDCM experiment of the medium-spin states in $^{106}\text{Cd}$

For population and analysis of the medium-spin states in  $^{106}\text{Cd}$ , an experiment was performed at the Wright Nuclear Structure Laboratory, using the New Yale Plunger Device [5] and SPEEDY  $\gamma$ -ray array [6] consisting of seven HPGe clover detectors, four at  $41.5^\circ$  and three at  $138.5^\circ$ , with both angles relative to the beam axis. The  $^{98}\text{Mo}(^{12}\text{C}, 4n)^{106}\text{Cd}$  reaction channel was utilised, with  $E(^{12}\text{C})_{\text{LAB}} = 60$  MeV. Further details of the experiment can be found in [7].

### 2.2. Lifetime determination of isomeric states in $^{106}\text{Cd}$

The deduction of the  $I^\pi = 9^-$  and  $8^-$  isomeric state lifetimes was performed using the  $330\mu\text{m}$  and  $2008\mu\text{m}$  target-stopper distances from the Yale experiment. The lifetime of the  $I^\pi = 9^-$  state at  $E_x = 3678$  keV in  $^{106}\text{Cd}$  was deduced by gating on the shifted component of the 646 keV,  $I^\pi = 11^- \rightarrow 9^-$  transition and projecting, fitting, deconvoluting and normalising the stopped and shifted components of the 269 keV,  $I^\pi = 9^- \rightarrow 7^-$  transition, as detailed in [8]. The deduced mean lifetime,  $\tau$ , of the  $I^\pi = 9^-$  state at  $E_x = 3678$  keV is  $0.89(20)$  ns.

A similar procedure was performed for the mean lifetime of the  $I^\pi = 8^-$  state at  $E_x = 3507$  keV in  $^{106}\text{Cd}$  by gating on the shifted component of the 598 keV,  $I^\pi = 10^- \rightarrow 8^-$  transition and projecting, fitting, deconvoluting and normalising the stopped and shifted peaks of the 188 keV,  $I^\pi = 8^- \rightarrow 6^-$  transition. The deduced mean lifetime of the  $I^\pi = 8^-$  state at  $E_x = 3507$  keV is  $1.7(5)$  ns.

### 2.3. DDCM experiment of the low-spin states in $^{106}\text{Cd}$

A second experiment was performed at the Institut für Kernphysik, Universität zu Köln, which utilised the Köln plunger and the  $^{96}\text{Mo}(^{13}\text{C}, 3n)^{106}\text{Cd}$  reaction at  $E(^{13}\text{C})_{\text{LAB}} = 43$  MeV. In this experiment, twenty distances

were measured, eight of which ( $6\ \mu\text{m}$ ,  $8\ \mu\text{m}$ ,  $13\ \mu\text{m}$ ,  $16\ \mu\text{m}$ ,  $18\ \mu\text{m}$ ,  $21\ \mu\text{m}$ ,  $25\ \mu\text{m}$  and  $37\ \mu\text{m}$ ) are used in the preliminary analysis presented here. The reaction  $\gamma$  rays were detected using seven individual segments of one germanium cluster detector (one segment was at an angle of  $0^\circ$  and the other six segments were at an angle of  $34.5^\circ$  relative to the beam axis) and five additional single crystal germanium detectors, each at an angle of  $141.5^\circ$  relative to the beam axis.

For both experiments, prompt coincidences were sorted into angle-dependent  $\gamma$ - $\gamma$  matrices and were analysed with the TV matrix viewer [9]. The lifetimes were deduced by using the Differential Decay Curve Method (DDCM) [10].

#### 2.4. Preliminary analysis of the $I^\pi = 2_1^+$ state lifetime

From the Köln experiment, three separate 1 keV wide energy coincidence gates were placed on the backward shifted component of the 861 keV,  $I^\pi = 4_1^+ \rightarrow 2_1^+$  transition. Projecting, fitting, deconvoluting and normalising the stopped and backward shifted components of the 633 keV,  $I^\pi = 2_1^+ \rightarrow 0_1^+$




Fig. 1. Left: Projection and deconvolution of the stopped and backward-shifted components of the 633 keV,  $I^\pi = 2_1^+ \rightarrow 0_1^+$  transition in  $^{106}\text{Cd}$ . The gate was set on the backward-shifted component of the 861 keV  $I^\pi = 4_1^+ \rightarrow 2_1^+$  transition for a series of distances between  $6\ \mu\text{m}$  and  $37\ \mu\text{m}$ . Right bottom: Normalised intensities of the stopped (decreasing as a function of distance) and shifted (increasing as a function of distance) components of the 633 keV  $I^\pi = 2_1^+ \rightarrow 0_1^+$  transition. Right top: Corresponding mean lifetimes at each individual distance measured. The weighted mean lifetime of this particular gate for the  $I^\pi = 2_1^+$  state at  $E_x = 633\ \text{keV}$  is 16.7(16) ps.

transition yields mean lifetimes of 15.5(14) ps, 16.7(16) ps (see Fig. 1) and 17.4(19) ps. The weighted mean of these values yielded a mean lifetime of the  $I^\pi = 2_1^+$  state of 16.4(9) ps.

### 3. Discussion and conclusion

For the isomeric states, the  $I^\pi = 9^-$  and  $I^\pi = 8^-$  mean lifetimes of 0.89(20) ns and 1.7(5) ns compare well to the previously reported values of 1.0(+2,-4) ns and 1.7(6) ns deduced by the ‘‘centroid shift method’’ [11]. For the  $I^\pi = 2_1^+$  state, the mean lifetime of 16.4(9) ps, presented here, differs from the literature value of 10.43(9) ps deduced from Coulomb excitation [2].

S.F.A. would like to acknowledge financial support from EPSRC DTG studentship. Work supported in part by the US DOE under grant nos DE-FG02-91ER-40609 and DE-FG02-88ER-40417. P.H.R. would like to acknowledge financial support from EPSRC and the Yale University Flint and Science Development Funds. J.J. and A.L. would like to acknowledge financial support from the Deutsche Forschungsgemeinschaft.

### REFERENCES

- [1] S.W. Yates, *J. Phys.* **G31**, S1393 (2005).
- [2]  $^{104}\text{Cd}$ : G.A. Müller *et al.*, *Phys. Rev.* **C64**, 014305 (2001);  
 $^{106}\text{Cd}$ : M. T. Esat *et al.*, *Nucl. Phys.* **A274**, 237 (1976);  
 $^{108}\text{Cd}$ : I. Thorslund *et al.*, *Nucl. Phys.* **A564**, 285 (1993);  
 $^{110}\text{Cd}$ : S. Harissopulos *et al.*, *Nucl. Phys.* **A683**, 157 (2001).
- [3] N. Boelaert, Masters Thesis, Universiteit Gent, Belgium, 2006.
- [4]  $^{106}\text{Cd}$ : P.H. Regan, *et al.*, *Nucl. Phys.* **A586**, 351 (1995);  
 $^{108}\text{Cd}$ : M. Piiparinen, *et al.*, *Nucl. Phys.* **A565**, 671 (1993);  
 $^{110}\text{Cd}$ : S. Juutinen *et al.*, *Z. Phys.* **A336**, 475 (1990);  
S. Juutinen, *et al.*, *Nucl. Phys.* **A573**, 306 (1994).
- [5] R. Krücken, *J. Res. Natl. Stand. Technol.* **105**, 53 (2000).
- [6] C.W. Beausang *et al.*, *Nucl. Instrum. Methods* **A452**, 431 (2000).
- [7] K. Andgren, *et al.*, *J. Phys. G* **31**, S1563 (2005).
- [8] S.F. Ashley, Ph.D. Thesis, University of Surrey, UK, to be submitted.
- [9] J. Theuerkauf, TV, unpublished, <http://www.ikp.uni-koeln.de/~fitz>
- [10] A. Dewald *et al.*, *Z. Phys.* **A364**, 163 (1989); G. Böhm, *et al.*, *Nucl. Instrum. Methods* **A329**, 248 (1993).
- [11] W. Andrejtscheff *et al.*, *Nucl. Phys.* **A437**, 167 (1985).