

LEPTON GENERATION-WEIGHTING FACTORS AND NEUTRINO MASS FORMULA

WOJCIECH KRÓLIKOWSKI

Institute of Theoretical Physics, Warsaw University
Hoża 69, 00-681 Warszawa, Poland

(Received March 7, 2005)

A candidate for the simple empirical neutrino mass formula is found, predicting the mass proportion $m_1 : m_2 : m_3 = 0 : 4 : 24$ and so, the mass ratio $\Delta m_{32}^2 / \Delta m_{21}^2 = 35$ not inconsistent with its experimental estimate. It involves only one free parameter and three generation-weighting factors suggested by the successful mass formula found previously for charged leptons (the simplest neutrino mass formula would predict $m_1 : m_2 : m_3 = 1 : 4 : 24$ and thus, $\Delta m_{32}^2 / \Delta m_{21}^2 \simeq 37$). A more involved variation of this equation follows from a special seesaw neutrino model with specifically “conspiring” Dirac and Majorana neutrino mass matrices. In this variation $m_1 : m_2 : m_3 \simeq \varepsilon^{(\nu)} : 4 : 24$, where $O(\varepsilon^{(\nu)}) = 10^{-2}$.

PACS numbers: 12.15.Ff, 14.60.Pq, 12.15.Hh

Some time ago we found an efficient empirical mass formula for charged leptons $e_i = e^-, \mu^-, \tau^-$ [1]. This formula reads

$$m_{e_i} = \mu^{(e)} \rho_i \left(N_i^2 + \frac{\varepsilon^{(e)} - 1}{N_i^2} \right), \quad (1)$$

where

$$N_i = 1, 3, 5, \quad (2)$$

and

$$\rho_i = \frac{1}{29}, \frac{4}{29}, \frac{24}{29}, \quad (3)$$

($\sum_i \rho_i = 1$). Here, $\mu^{(e)} > 0$ and $\varepsilon^{(e)} > 0$ are constants. In fact, with the experimental values $m_e = 0.510999$ MeV and $m_\mu = 105.658$ MeV as an

input, the formula (1), rewritten explicitly as

$$m_e = \frac{\mu^{(e)}}{29} \varepsilon^{(e)}, \quad m_\mu = \frac{\mu^{(e)}}{29} \frac{4}{9} (80 + \varepsilon^{(e)}), \quad m_\tau = \frac{\mu^{(e)}}{29} \frac{24}{25} (624 + \varepsilon^{(e)}), \quad (4)$$

leads to *the prediction*

$$m_\tau = \frac{6}{125} (351m_\mu - 136m_e) = 1776.80 \text{ MeV} \quad (5)$$

and also determines both constants

$$\mu^{(e)} = \frac{29(9m_\mu - 4m_e)}{320} = 85.9924 \text{ MeV}, \quad \varepsilon^{(e)} = \frac{320m_e}{9m_\mu - 4m_e} = 0.172329. \quad (6)$$

The prediction (5) is really close to the experimental value $m_\tau^{\text{exp}} = 1776.99^{+0.29}_{-0.26} \text{ MeV}$ [2].

Though the formula (1) has essentially the empirical character, there exists a speculative background for it based on a Kähler-like extension of Dirac equation that the interested reader may find in Ref. [1]. In particular, the numbers N_i and ρ_i ($i = 1, 2, 3$) given in Eqs. (2) and (3) are interpreted there. Let us only mention that $N_i - 1 = 0, 2, 4$ is the number of *additional* bispinor indices appearing in the extended Dirac equation and obeying Fermi statistics that enforces their antisymmetrization and so, restricts to zero the related additional spin. This Fermi statistics is also the reason, why there are precisely *three* Standard Model fermion generations *i.e.*, $N_i - 1 = 0, 2, 4$, since any additional bispinor index can assume *four* values, what implies that $N_i - 1 \leq 4$ (after the antisymmetrization of additional bispinor indices). Thus, an analogue of Pauli principle works (intrinsically), restricting the number of additional bispinor indices to ≤ 4 and so, resulting into three and only three generations of leptons and quarks (all with spin 1/2). The generation-weighting factors ρ_i multiplied by 29, $29\rho_i = 1, 4, 24$ ($\sum_i \rho_i = 1$), tell us, how many times the lepton or quark wave functions of three generations are realized (up to the factor ± 1) by the extended Dirac equation.

Now, it is tempting to seek in the same framework an efficient empirical mass formula for mass neutrinos $\nu_i = \nu_1, \nu_2, \nu_3$ (being the mass states of the flavor neutrinos $\nu_\alpha = \nu_e, \nu_\mu, \nu_\tau$).

As is well known, the mass neutrinos display a less hierarchical spectrum than the charged leptons. In fact, neutrino oscillation experiments give actually the following estimates [3] for $\Delta m_{ji}^2 \equiv m_{\nu_j}^2 - m_{\nu_i}^2$: the ranges

$$7.2 < \frac{\Delta m_{21}^2}{(10^{-5} \text{ eV}^2)} < 9.1, \quad 1.9 < \frac{\Delta m_{32}^2}{(10^{-3} \text{ eV}^2)} < 3.0 \quad (7)$$

and the best fits

$$\Delta m_{21}^2 \sim 8.1 \times 10^{-5} \text{ eV}^2, \quad \Delta m_{32}^2 \sim 2.4 \times 10^{-3} \text{ eV}^2. \quad (8)$$

For the values (8) $\Delta m_{32}^2/\Delta m_{21}^2 \sim 30$. Notice that $1.9/0.091 \sim 21$ and $3.0/0.072 \sim 42$ and so, the experimental limits are $21 < \Delta m_{32}^2/\Delta m_{21}^2 < 42$.

Thus, let us tentatively try for the neutrino mass formula the simplest conjecture

$$m_{\nu_i} = \mu^{(\nu)} \rho_i, \quad (9)$$

where the generation-weighting factors ρ_i as given in Eq. (3) still appear, while the numbers N_i numerating the generations and defined in Eq. (2) are absent. Here, $\mu^{(\nu)} > 0$ is a constant.

The tentative mass formula (9), rewritten as

$$m_{\nu_1} = \frac{1}{29} \mu^{(\nu)}, \quad m_{\nu_2} = \frac{4}{29} \mu^{(\nu)}, \quad m_{\nu_3} = \frac{24}{29} \mu^{(\nu)}, \quad (10)$$

implies that

$$m_{\nu_1} : m_{\nu_2} : m_{\nu_3} = 1 : 4 : 24 \quad (11)$$

and

$$\mu^{(\nu)} = m_{\nu_1} + m_{\nu_2} + m_{\nu_3} = 29 m_{\nu_1} = \frac{29}{4} m_{\nu_2} = \frac{29}{24} m_{\nu_3}. \quad (12)$$

From Eq. (11)

$$\frac{\Delta m_{32}^2}{\Delta m_{21}^2} = \frac{112}{3} = 37.3333. \quad (13)$$

Thus, using the experimental range (7) of Δm_{21}^2 and its experimental best fit (8) as an input, we get the following *prediction*: the range

$$2.7 < \frac{\Delta m_{32}^2}{10^{-3} \text{ eV}^2} < 3.4 \quad (14)$$

and the best fit:

$$\Delta m_{32}^2 \sim 3.0 \times 10^{-3} \text{ eV}^2. \quad (15)$$

The predicted range (14) of Δm_{32}^2 is not inconsistent with its experimental range (7), but its predicted best fit (15) appears too large in comparison with the experimental best fit (8) (though the predicted ratio (13) remains within its experimental limits $21 < \Delta m_{32}^2/\Delta m_{21}^2 < 42$). Note that making use of the best fit (15) for Δm_{32}^2 , we would *predict* from Eq. (11)

$$m_{\nu_1} \sim 2.3 \times 10^{-3} \text{ eV}, \quad m_{\nu_2} \sim 9.3 \times 10^{-3} \text{ eV}, \quad m_{\nu_3} \sim 5.6 \times 10^{-2} \text{ eV} \quad (16)$$

and determine from Eq. (12)

$$\mu^{(\nu)} \sim 6.7 \times 10^{-2} \text{ eV} . \quad (17)$$

Here, the only input is the experimental estimate (8) of Δm_{21}^2 .

We may argue that the tentative mass formula (9) requires a correction for the smallest neutrino mass m_{ν_1} , if the neutrino masses are related (*grosso modo*) to the additional bispinor indices in the general Dirac equation applied to the neutrino triplet. Then, for the ν_1 neutrino — that does not involve additional indices — we ought to expect $m_{\nu_1} = 0$ (at least approximately). This conjecture may lead to the correction factor $1 - \delta_{i1}$ in the mass equation (9). In consequence, the corrected neutrino mass formula may read

$$m_{\nu_i} = \mu^{(\nu)} \rho_i (1 - \delta_{i1}) . \quad (18)$$

This mass formula, rewritten as

$$m_{\nu_1} = 0 , \quad m_{\nu_2} = \frac{4}{29} \mu^{(\nu)} , \quad m_{\nu_3} = \frac{24}{29} \mu^{(\nu)} , \quad (19)$$

gives

$$m_{\nu_1} : m_{\nu_2} : m_{\nu_3} = 0 : 4 : 24 \quad (20)$$

and

$$\mu^{(\nu)} = \frac{29}{28} (m_{\nu_2} + m_{\nu_3}) = \frac{29}{4} m_{\nu_2} = \frac{29}{24} m_{\nu_3} . \quad (21)$$

From Eq. (20)

$$\frac{\Delta m_{32}^2}{\Delta m_{21}^2} = 35 . \quad (22)$$

Hence, making use of the experimental range (7) of Δm_{21}^2 and its experimental best fit (8) as an input, we obtain the following *prediction*: the range

$$2.5 < \frac{\Delta m_{32}^2}{10^{-3} \text{ eV}^2} < 3.2 \quad (23)$$

and the best fit

$$\Delta m_{32}^2 \sim 2.8 \times 10^{-3} \text{ eV}^2 . \quad (24)$$

The predicted range (23) of Δm_{32}^2 is a little closer to its experimental range (7) than the previous range (14) (both being not inconsistent with (7)). Also the predicted best fit (24) is a bit closer to its actual experimental best fit (8) than the previous best fit (15) (both being too large, though the predicted ratios (13) and (22) remain within their actual experimental limits

$21 < \Delta m_{32}^2 / \Delta m_{21}^2 < 42$). Note that using the best fit (24) for Δm_{32}^2 , we would predict from Eq. (20)

$$m_{\nu_1} \sim 0, \quad m_{\nu_2} \sim 9.0 \times 10^{-3} \text{ eV}, \quad m_{\nu_3} \sim 5.4 \times 10^{-2} \text{ eV} \quad (25)$$

and determine from Eq. (21)

$$\mu^{(\nu)} \sim 6.5 \times 10^{-2} \text{ eV}. \quad (26)$$

The experimental best fit (8) for Δm_{21}^2 is the only input here.

Naturally, the actual experimental best fit $\Delta m_{21}^2 \sim 8.1 \times 10^{-5} \text{ eV}^2$ (giving $\Delta m_{32}^2 \sim 3.0 \times 10^{-3} \text{ eV}^2$ through Eq. (22)) may change in the course of further experiments. For instance, if (drastically) it turned out as small as $\Delta m_{21}^2 \sim (6.9 - 7.2) \times 10^{-5} \text{ eV}^2$, we would predict from Eq. (22) that $\Delta m_{32}^2 \sim (2.4 - 2.5) \times 10^{-3} \text{ eV}^2$. Then, from Eq. (20)

$$m_{\nu_1} \sim 0, \quad m_{\nu_2} \sim (8.3 - 8.5) \times 10^{-3} \text{ eV}, \quad m_{\nu_3} \sim (5.0 - 5.1) \times 10^{-2} \text{ eV} \quad (27)$$

and from Eq. (21)

$$\mu^{(\nu)} \sim (6.0 - 6.1) \times 10^{-2} \text{ eV}. \quad (28)$$

Similarly, the actual experimental best fit $\Delta m_{32}^2 \sim 2.4 \times 10^{-3} \text{ eV}^2$ (giving $\Delta m_{21}^2 \sim 6.9 \times 10^{-5} \text{ eV}^2$ by means of Eq. (22)) may change. For example, if (drastically) it appeared as large as $\Delta m_{32}^2 \sim (2.8 - 2.9) \times 10^{-3} \text{ eV}^2$, we would predict from Eq. (22) that $\Delta m_{21}^2 \sim (8.0 - 8.3) \times 10^{-5} \text{ eV}^2$. Then, from Eq. (20)

$$m_{\nu_1} \sim 0, \quad m_{\nu_2} \sim (9.0 - 9.1) \times 10^{-3} \text{ eV}, \quad m_{\nu_3} \sim (5.4 - 5.5) \times 10^{-2} \text{ eV} \quad (29)$$

and from Eq. (21)

$$\mu^{(\nu)} \sim (6.5 - 6.6) \times 10^{-2} \text{ eV}. \quad (30)$$

The neutrino mass formula (18) is not of the seesaw form. At any rate, no seesaw elements were used in its formulation. However, we constructed recently [4] a special seesaw neutrino model — with the Dirac and Majorana neutrino mass matrices living in a specific “conspiracy” [5] — that leads to the neutrino mass formula

$$m_{\nu_i} = \mu^{(\nu)} \rho_i \left(1 + \frac{\varepsilon^{(\nu)} - 1}{N_i^4} \right), \quad (31)$$

where $\varepsilon^{(\nu)} > 0$ is a new constant. In Ref. [4] this constant gets the small value

$$\varepsilon^{(\nu)} \sim 7.35 \times 10^{-3}. \quad (32)$$

We can see that the mass spectra (18) and (31) are practically identical for m_{ν_2} and m_{ν_3} , but differ for m_{ν_1} which becomes now nonzero since the mass formula (31) implies

$$m_{\nu_1} : m_{\nu_2} : m_{\nu_3} \simeq \varepsilon^{(\nu)} : 4 \cdot \frac{80}{81} : 24 \cdot \frac{624}{625} \quad (33)$$

and

$$\begin{aligned} \mu^{(\nu)} &\simeq \frac{29}{\varepsilon^{(\nu)} + 28} (m_{\nu_1} + m_{\nu_2} + m_{\nu_3}) \\ &\simeq \frac{29}{28} (m_{\nu_1} + m_{\nu_2} + m_{\nu_3}) = \frac{29}{\varepsilon^{(\nu)}} m_{\nu_1} = \frac{29}{4} \frac{81}{80} m_{\nu_2} = \frac{29}{24} \frac{625}{624} m_{\nu_3}. \end{aligned} \quad (34)$$

From Eq. (33)

$$\frac{\Delta m_{32}^2}{\Delta m_{21}^2} \simeq 36, \quad (35)$$

what is larger by 1 than the value (22). Thus, using the experimental range (7) of Δm_{21}^2 and its experimental best fit (8) as an input, we obtain as a *prediction* the range very similar to (23):

$$2.6 < \frac{\Delta m_{32}^2}{10^{-3} \text{ eV}^2} < 3.3 \quad (36)$$

and the best fit very similar to (24):

$$\Delta m_{32}^2 \sim 2.9 \times 10^{-3} \text{ eV}^2. \quad (37)$$

Notice that making use of the best fit (37) for Δm_{32}^2 , we would predict from Eqs. (33) and (32)

$$m_{\nu_1} \sim 1.7 \times 10^{-5} \text{ eV}, \quad m_{\nu_2} \sim 9.0 \times 10^{-3} \text{ eV}, \quad m_{\nu_3} \sim 5.5 \times 10^{-2} \text{ eV} \quad (38)$$

and from Eq. (34)

$$\mu^{(\nu)} \sim 6.6 \times 10^{-2} \text{ eV}. \quad (39)$$

In conclusion, it is exciting that the generation-weighting factors ρ_i , so efficient in the case of charged-lepton masses, can be also useful for neutrino masses, namely, for predicting their ratio $\Delta m_{32}^2/\Delta m_{21}^2$ up to the deviation 35–30 or 36–30 from its actual experimental estimation 30 (its predicted value 35 or 36 still remains within the actual experimental limits $21 < \Delta m_{32}^2/\Delta m_{21}^2 < 42$). This suggests the hypothesis that the proposed simple mass formula (18) or its seesaw variation (31) describes, at least approximately, the true character of neutrino mass spectrum.

Supplement

One can achieve the full agreement with the actual experimental estimate $\Delta m_{32}^2/\Delta m_{21}^2 \sim 30$ by introducing an appropriate second free parameter, but then the prediction for this ratio is lost. For example, the simplest mass formula

$$m_{\nu_i} = \mu^{(\nu)} \rho_i (1 - \beta \delta_{i3}) \quad (40)$$

evolving from Eq. (9), where $\beta > 0$ is the second free parameter, gives

$$m_{\nu_1} : m_{\nu_2} : m_{\nu_3} = 1 : 4 : 24(1 - \beta) \quad (41)$$

and

$$\mu^{(\nu)} = \frac{29}{5 + 24(1 - \beta)} (m_{\nu_1} + m_{\nu_2} + m_{\nu_3}) = 29m_{\nu_1} = \frac{29}{4}m_{\nu_2} = \frac{29}{24(1 - \beta)}m_{\nu_3}. \quad (42)$$

From Eq. (41)

$$\frac{\Delta m_{32}^2}{\Delta m_{21}^2} = \frac{16 [36(1 - \beta)^2 - 1]}{15}. \quad (43)$$

This leads to the value ~ 30 if

$$\beta \sim 0.10. \quad (44)$$

So, β is a small parameter ($\Delta m_{32}^2/\Delta m_{21}^2 = 112/3 \simeq 37$ for $\beta = 0$).

Using the experimental range (7) of Δm_{21}^2 and its experimental best fit (8), one gets

$$2.2 < \frac{\Delta m_{32}^2}{10^{-3} \text{ eV}^2} < 2.7 \quad (45)$$

and

$$\Delta m_{32}^2 \sim 2.4 \times 10^{-3} \text{ eV}^2. \quad (46)$$

Then, from Eq. (41)

$$m_{\nu_1} \sim 2.3 \times 10^{-3} \text{ eV}, \quad m_{\nu_2} \sim 9.2 \times 10^{-3} \text{ eV}, \quad m_{\nu_3} \sim 5.0 \times 10^{-2} \text{ eV} \quad (47)$$

and from Eq. (42)

$$\mu^{(\nu)} \sim 6.7 \times 10^{-2} \text{ eV}. \quad (48)$$

Here, the experimental estimates of Δm_{21}^2 and $\Delta m_{32}^2/\Delta m_{21}^2$ are both the input. But, one of three neutrino masses is still a prediction.

REFERENCES

- [1] W. Królikowski, *Acta Phys. Pol. B* **33**, 2559 (2002) [hep-ph/0203107]; and references therein.
- [2] Particle Data Group, *Review of Particle Physics*, *Phys. Lett.* **B592** (2004).
- [3] For a review *cf.* J.W.F. Valle, hep-ph/0410103; D. Kielczewska, *Nucl. Phys. (Proc. Suppl.)* **B136**, 77 (2004).
- [4] W. Królikowski, *Acta Phys. Pol. B* **35**, 2095 (2004) [hep-ph/0401101].
- [5] S.M. Barr, I. Dorsner, *Nucl. Phys.* **B585**, 79 (2000).