

IDENTIFICATION OF ISOMERIC STATES ‘SOUTH’ OF ^{208}Pb VIA PROJECTILE FRAGMENTATION*

S.J. STEER^a, ZS. PODOLYÁK^a, S. PIETRI^a, P.H. REGAN^a
 D. RUDOLPH^b, E. WERNER-MALENTO^{c,d}, A.B. GARNSWORTHY^{a,e}
 R. HOISCHEN^b, M. GÓRSKA^c, J. GERL^c, H.J. WOLLERSHEIM^c
 F. BECKER^c, P. BEDNARCZYK^{c,f}, L. CÁ CERES^{c,g}, P. DOORNENBAL^c
 H. GEISSEL^c, J. GRĘBOSZ^{c,f}, A. KELIC^c, N. KURZ^c
 F. MONTES^c, W. PROKOPOWICZ^{c,f}, T. SAITO^c, H. SCHAFFNER^c
 S. TACHENOV^c, A. HEINZ^e, M. PFÜTZNER^d, T. KURTUKIAN-NIETO^h
 G. BENZONIⁱ, A. JUNGCLAUS^g, D.L. BALABANSKI^{j,k}, C. BRANDAU^{a,c}
 A.M. BRUCE^l, W.N. CATFORD^a, I.J. CULLEN^a, ZS. DOMBRADI^m
 E. ESTEVEZ^h, W. GELLETLY^a, G. ILIE^{o,p}, J. JOLIEⁿ, G.A. JONES^a
 M. KMIĘCIK^f, F.G. KONDEV^p, S. LALKOVSKI^{g,l}, Z. LIU^a, A. MAJ^f
 S. MYALSKI^f, T. SHIZUMA^{a,s}, S. SCHWERTEL^t, P.M. WALKER^a
 O. WIELANDⁱ

^aDepartment of Physics, University of Surrey, Guildford, GU2 7XH, UK

^bDepartment of Physics, Lund University, S-22100 Lund, Sweden

^cGSi, Planckstrasse 1, D-64291, Darmstadt, Germany

^dInstitute of Experimental Physics, Warsaw University, Poland

^eWNSL, Yale University, New Haven, CT, USA

^fThe Institute of Nuclear Physics PAN, Kraków, Poland

^gDept. de Física Teórica, Universidad Autónoma de Madrid, Madrid, Spain

^hUniversidad de Santiago de Compostela, Santiago de Compostela, Spain

ⁱINFN, and Università degli Studi di Milano, Italy

^jINRNE, Bulgarian Academy of Sciences, Sofia, Bulgaria

^kDipartimento di Fisica, Università di Camerino, Italy

^lSchool of Engineering, University of Brighton, Brighton, BN2 4GJ, UK

^mInstitute of Nuclear Research, Debrecen, Hungary

ⁿIKP, Universität zu Köln, Germany

^oNational Institute of Physics and Nuclear Engineering, Bucharest, Romania

^pNED, Argonne National Laboratory, Argonne, USA

^rFaculty of Physics, University of Sofia, Sofia, Bulgaria

^sJapan Atomic Energy Agency, Kyoto, Japan

^tPhysik Department E12, Technische Universität München, Garching, Germany

(Received November 3, 2006)

* Presented at the Zakopane Conference on Nuclear Physics, September 4–10, 2006, Zakopane, Poland.

Relativistic fragmentation of ^{208}Pb has been used to produce excited states in neutron-rich nuclei with $N \approx 126$. Spectroscopic information for a range of nuclei has been obtained through observing delayed γ -ray emissions from isomeric states. Preliminary results for $^{203,204}\text{Pt}$ nuclei are presented. For the first time, excited states have been observed in ^{203}Pt and ^{204}Pt . The yrast structure of ^{204}Pt up to spin-parity, $I^\pi = (10^+)$ has been tentatively inferred from the internal decay of two isomeric states.

PACS numbers: 21.10.-k, 23.20.-g, 25.70.Mn, 27.80.+w

1. Introduction

The Rare Isotopes Investigations at GSI (RISING) project has begun the Stopped Beam phase of experiments [1]; reported here are the first results from one of these measurements. Rare, exotic, neutron-rich nuclei in the vicinity of $N = 126$ were produced in the relativistic projectile fragmentation of a ^{208}Pb beam.

A number of $N \approx 126$ nuclei were produced during the fragmentation process. These present an ideal testing ground for the shell model, because of the limited number of valence nucleons involved in forming the low lying, yrast and near yrast excited states in these nuclei. The analysis of such states permits the extraction of information on the single-particle energies and residual shell-model interactions in this region.

2. Experimental details and results

The 1 GeV/nucleon ^{208}Pb beam, provided by the SIS-18 accelerator at GSI, was fragmented on a beryllium target (thickness 2.5 g/cm²). The FRagment Separator (FRS) [2], operated in standard achromatic mode, was used to separate and identify the desired nuclei (see Fig 1). Once identified, nuclei were brought to a halt in a 7 mm plastic stopper. The delayed γ rays were detected over a 100 μs time range after implantation using the high efficiency RISING array [1], which, in its Stopped Beam configuration, surrounded the implantation target.

Delayed γ -ray spectra associated with $^{202,203,204}\text{Pt}$ are shown in Fig. 2. The observation of the previously identified isomeric state in ^{202}Pt [3] confirms the particle identification. Gamma rays belonging to ^{203}Pt and ^{204}Pt have been seen for the first time in this experiment. The ground state of ^{203}Pt was previously identified in projectile fragmentation at GSI [5], but the isomer in ^{203}Pt represents the first spectroscopic information on this $N = 125$ nucleus. Similarly, this is the first observation of excited states in ^{204}Pt ; transitions from two isomeric states have been observed, with three γ rays (96, 1061, 1158 keV) originating from a shorter lived metastable state and two γ rays (872, 1123 keV) originating from a lower-lying, longer-lived isomer.




Fig. 1. Fragment identification. (A) Energy loss at the middle focal plane *versus* energy loss at the final focal plane. Charge state changes are distinguished here, allowing selection between nuclei that undergo no change, the loss of or gaining of an electron in the S2 achromatic degrader. (B) A/Q *versus* S4 position tangential to the beam direction. Only $\Delta Q = 0$ nuclei are plotted. Locations of the previously observed ^{202}Pt [3] and the newly identified ^{204}Pt nuclei are highlighted.




Fig. 2. Delayed γ -ray spectra associated with $^{202,203,204}\text{Pt}$ and ^{206}Hg . Note that the non-physical background suppression of low energy γ rays (up to ~ 300 keV) is a result of data analysis techniques. Similar γ -ray spectra are observed for ^{206}Hg and ^{204}Pt , which exhibit 2-proton hole and 4-proton hole structures, respectively.

In producing ^{204}Pt four protons are removed from doubly closed shell ^{208}Pb . Like ^{206}Hg , ^{204}Pt has two isomeric states (see Fig. 2). The higher lying isomer in both cases is expected to be of a $(\pi_{h_{\frac{11}{2}}})_{10^+}^{-2}$ character [4]. This structure is associated with seniority two proton-hole states in the $Z = 82$ shell closure. A more detailed interpretation will be the subject of a future publication.

In conclusion, new spectroscopic information has been obtained on a range of heavy, neutron-rich nuclei, populated in relativistic energy projectile fragmentation. For the first time, excited states in $N = 125$ ^{203}Pt and the $N = 126$ ^{204}Pt have been presented, including the identification of three isomeric states.

This work is sponsored by the EPSRC (UK), the Swedish Research Council, the Polish Ministry of Science and Higher Education (grants 1-P03B-030-30 and 620/E-77/SPB/GSI/P-03/DWM105/2004-2007), the Bulgarian Science Fund VUF06/05, the US Department of Energy (grants DE-FG02-91ER-40609 and W-31-109-ENG-38), the German Federal Ministry of Education and Research under grant 06KY205I and EURONS (European Commission contract number 506065). ABG would like to acknowledge financial support from Nexia Solutions Ltd, FGK acknowledges the support of the US DOE, Office of Nuclear Physics, under Contract No. DE-AC02-06CH11357 and AJ acknowledges the spanish Ministerio de Educacion y Ciencia under project number FPA2005-00696.

REFERENCES

- [1] S. Pietri, *et al.*, *Acta Phys. Pol. B* **38**, 1255 (2007) these proceedings.
- [2] H. Geissel, *Nucl. Instrum. Methods Phys. Res.* **B70**, 286 (1992).
- [3] M. Caamaño, *et al.*, *Eur. Phys. J.* **A23**, 201 (2005); Zs. Podolyák, *et al.*, *Nucl. Phys.* **A722**, 273c (2003).
- [4] B. Fornal, *et al.*, *Phys. Rev. Lett.* **87**, 215201 (2001); A.R. Poletti, *et al.*, *Nucl. Phys.* **A580**, 64 (1994).
- [5] T. Kurtukian-Nieto, *et al.*, *AIP Conf. Proc.* **802**, 73 (2005).