

A SIMPLE METHOD OF CALCULATING EFFECTIVE OPERATORS

NAOYUKI HABA, KUNIO KANETA

Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

SHIGEKI MATSUMOTO

IPMU, TODIAS, University of Tokyo, Kashiwa, 277-8583, Japan

TAKEHIRO NABESHIMA

Department of Physics, University of Toyama, Toyama 930-8555, Japan

(Received July 18, 2011)

It is important to obtain effective operators by integrating out high energy degrees of freedom in physics. We suggest a general method of calculating accurate irrelevant operators in a scattering process without use of equation of motions. By using this method, for example, we will represent a complete set of dimension six operators in QCD, which are induced from physics beyond the standard model, supersymmetry and universal extra dimension. We will also show an example of effective anomalous 4-Fermi interactions induced from a little Higgs model.

DOI:10.5506/APhysPolB.43.405

PACS numbers: 11.15.Bt

1. Introduction

It is important to obtain effective operators by integrating out high energy degrees of freedom in physics. In a quantum field theory, we can obtain effective Lagrangian by integrating out high energy momentum and heavy particles. And their effects are introduced into irrelevant operators in a low energy effective theory. Therefore, it is important to obtain effective irrelevant operators accurately for a search of new physics. As for high energy physics, search for physics beyond the standard model (SM) is one of the main subjects in Large Hadron Collider (LHC). For both theoretical and numerical analyses to search new physics, we stress again that it is important to obtain accurate irrelevant operators because they include the hints of new physics.

There have been some works of listing higher dimensional operators in the field content of the SM (or also including right-handed neutrinos.) Allowed irrelevant operators in an effective theory should be determined by symmetries existed in the theory. Let us focus in this paper, mainly on dimension six operators. Possible dimension six operators within the SM field content were listed in Ref. [1]. Meanwhile, this set was not irreducible, and Refs. [2,3] have obtained a complete set systematically. However, Ref. [4,5,6] insisted that 80 numbers of complete set can be diminishable to 59 by using equation of motions (EOMs).

In this paper, we show a general method of calculating accurate irrelevant effective operators in a scattering process without use of EOMs. By using this method, we will represent dimension six operators induced from supersymmetry (SUSY) [7], universal extra dimension (UED) [8], and little Higgs model (LH) [9]. We will show coefficients of dimension six operators in QCD by integrating out sparticles (KK particles) in SUSY (UED). We will also show coefficients of 4-Fermi operators originating from anomalous interactions in LH. They are promising candidates of beyond the SM, and our analyses will shed lights on the search of beyond the SM. Our method can apply to other quantum field theories in any dimensions, so that we believe this technique is very useful in a lot of researches in physics.

2. Method of obtaining effective operators

Let us show a general method of obtaining accurate irrelevant operators which are useful for calculating a scattering processes in an effective theory. The effects of beyond the SM must contain in higher order irrelevant operators in general. Thus, when we integrate out new particles and high momentum (physics) above an energy scale of Λ , the effective theory $\mathcal{L}_{\text{eff}}$ can be expanded as a power of Λ^{-1} as

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_0 + \frac{1}{\Lambda} \mathcal{L}_1 + \frac{1}{\Lambda^2} \mathcal{L}_2 + \cdots, \quad (2.1)$$

where $\mathcal{L}_0$ is the SM Lagrangian, and $\mathcal{L}_1$ ($\mathcal{L}_2$) represents dimension five (six) operators. As for dimension five operator, there is only one operator written within the SM field contents, which induces Majorana neutrino masses. Thus, let us mainly focus on a non-trivial next lowest operator, *i.e.*, dimension six operators

$$\mathcal{L}_2 = \sum_i c_i \mathcal{O}_i^{(6)}, \quad (2.2)$$

where c_i is a coefficient, and i is the index of all possible dimension six operators allowed by the SM gauge symmetry. How can we calculate dimension

six operators in the effective Lagrangian by integrating out high energy degrees of freedom? One correct answer is to take a path integral of the full theory as

$$Z = \int \mathcal{D}\phi_{\text{SM}} \mathcal{D}\phi_h e^{iS[\phi_{\text{SM}}, \phi_h]}, \quad (2.3)$$

where ϕ_{SM} and ϕ_h represent the SM fields and heavy fields, respectively. By integrating out ϕ_h as

$$e^{iS_{\text{eff}}[\phi_{\text{SM}}]} = \int \mathcal{D}\phi_h e^{iS[\phi_{\text{SM}}, \phi_h]} \quad (2.4)$$

we can obtain an effective action

$$S_{\text{eff}}[\phi_{\text{SM}}] = S_{\text{SM}} + S_1[\phi_{\text{SM}}] + S_2[\phi_{\text{SM}}] + \cdots, \quad (2.5)$$

where $S_1 = (1/\Lambda) \int \mathcal{L}_1$ and $S_2 = (1/\Lambda^2) \int \mathcal{L}_2$. Coefficients of dimension six operators have been basically calculated in $S_2[\phi_{\text{SM}}]$, by which a S-matrix element in this effective theory could be estimated.

Let us reveal more about the strategy to obtain irrelevant operators. We now consider a Lagrangian such as

$$\mathcal{L}[\phi_l, \phi_h] = \mathcal{L}[\phi_l] + F_i[\phi_l]\phi_{h,i} + \frac{1}{2}K_{ij}[\phi_l]\phi_{h,i}\phi_{h,j}, \quad (2.6)$$

where $\phi_l(\phi_h)$ represent light (heavy) scalar fields, and i denotes a index of scalar fields. The effective Lagrangian can be obtained using Gaussian integral of ϕ_h as

$$e^{iS_{\text{eff}}[\phi_l]} = \exp \left[-\frac{1}{2} \text{Tr} \log K[\phi_l] - \frac{i}{2} \int d^4x d^4y F_i[\phi_l](x) K_{ij}^{-1}[\phi_l](x, y) F_j[\phi_l](y) \right]. \quad (2.7)$$

Note that Eq. (2.7) is exact when the Lagrangian is up to $\mathcal{O}(\phi_h^2)$. Although the operators in Eq. (2.7) are non-local, they can be expanded by infinite number of local operators. Noting mass of $\phi_l(\phi_h)$ as $\mu(\Lambda)$, the non-local operators can be expanded in powers of μ/Λ . Then, $K[\phi_l]$ in Eq. (2.7) is divided into two parts as

$$K[\phi_l] = K^h + K^l[\phi_l], \quad (2.8)$$

where K^h is the kinetic term of ϕ_h , and $K^l[\phi_l]$ is ϕ_l dependent part of $K[\phi_l]$. Expanding momentum of $K^l[\phi_l]$ by Λ , and regarding $K^l[\phi_l]/K^h \sim$

$\mathcal{O}(\mu^2/\Lambda^2) \ll 1$, $\text{Tr} \log K[\phi_l]$ and $K[\phi_l]^{-1}$ are given by

$$\begin{aligned} \text{Tr} \log K[\phi_l] &= \text{Tr} \log \left(K^h \right)^{-1} + \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \text{Tr} \left(\left(K^h \right)^{-1} K^l[\phi_l] \right)^n, \quad (2.9) \\ K[\phi_l]^{-1} &= \left(K^h + K^l[\phi_l] \right)^{-1} \sim \left(K^h \right)^{-1} - \left(K^h \right)^{-1} K^l[\phi_l] \left(K^h \right)^{-1} + \cdots, \quad (2.10) \end{aligned}$$

where $(K^h)^{-1} = (K^h)^{-1}(x, y)$ is the propagator of ϕ_h except for imaginary factor i . Then, the action, S_{eff} , can be written as

$$\begin{aligned} S_{\text{eff}}[\phi_l] &= \int d^4x \left[\mathcal{L}[\phi_l] + \frac{i}{2} \int d^4y \delta^4(y-x) (K^h)^{-1}(x, y) K^l[\phi_l](y) \right. \\ &+ \frac{i}{4} \int d^4y (K^h)^{-1}(x, y) K^l[\phi_l](y) (K^h)^{-1}(y, x) K^l[\phi_l](x) + \cdots \\ &- \frac{1}{2} \int d^4y F_i[\phi_l](x) \left\{ (K^h)^{-1}(x, y) \right. \\ &\left. \left. - \int d^4z (K^h)^{-1}(x, z) K^l[\phi_l](z) (K^h)^{-1}(z, y) K^l[\phi_l](y) + \cdots \right\} F_j[\phi_l](y) \right]. \quad (2.11) \end{aligned}$$

In this expression, we can obtain an effective theory with infinite number of local operators.

We comment that the perturbative expansion in Eqs. (2.9) and (2.10) does not depend on $F[\phi_l]$ in Eq. (2.7). We can also consider a theory with interactions of $\phi_h^3, \phi_h^4, \dots$, where they can be treated perturbatively. For example, if there is an interaction, ϕ_h^3 , with coupling G_{ijk} , the Lagrangian is given by

$$\mathcal{L}'[\phi_l, \phi_h] = \mathcal{L}[\phi_l, \phi_h] + \frac{1}{3!} G_{ijk}[\phi_l] \phi_{h,i} \phi_{h,j} \phi_{h,k}. \quad (2.12)$$

By integrating out ϕ_h , the effective action, $S'_{\text{eff}}[\phi_l]$, is given by

$$\begin{aligned} S'_{\text{eff}}[\phi_l] &= S_{\text{eff}}[\phi_l] + \int d^4x A_{ijk}(x) G_{ijk}[\phi_l](x) \\ &+ \int \int d^4x d^4y B_{ijklmn}(x, y) G_{ijk}[\phi_l](x) G_{lmn}[\phi_l](y) + \cdots \quad (2.13) \end{aligned}$$

with dimensionful couplings A_{ijk} and B_{ijklmn} . They are calculated as

$$A_{ijk}(x) = \frac{\delta}{\delta G_{ijk}(x)} e^{iS'_{\text{eff}}} \Big|_{G_{ijk}=0}, \quad (2.14)$$

$$B_{ijklmn}(x, y) = \frac{\delta^2}{\delta G_{lmn}(y) \delta G_{ijk}(x)} e^{iS'_{\text{eff}}} \Big|_{G_{ijk}=0}. \quad (2.15)$$

Apparently, A_{ijk} is zero. B_{ijklmn} is $\langle (\phi_{h,i} \phi_{h,j} \phi_{h,k})_x (\phi_{h,l} \phi_{h,m} \phi_{h,n})_y \rangle_h$, where $\langle \mathcal{O} \rangle_h$ is defined by

$$\langle \mathcal{O} \rangle_h \equiv \int \mathcal{D}\phi_h \mathcal{O} e^{i \int \mathcal{L}[\phi_l, \phi_h]}. \quad (2.16)$$

In this stage, $S'_{\text{eff}}[\phi_l]$ is explicitly obtained. Before showing a concrete calculation method, we consider a role of EOMs when we calculate irrelevant operators.

We now stand in a position of showing our accurate method of calculating irrelevant operators without use of EOMs. Let us show a concrete calculation by using a toy model. We consider a Lagrangian

$$\begin{aligned} \mathcal{L} = & \bar{\psi}_l (i\partial - m) \psi_l + \bar{\psi}_h (i\partial - M) \psi_h + \frac{1}{2} (\partial_\mu \phi)^2 - \frac{1}{2} M^2 \phi^2 \\ & - g\phi (\bar{\psi}_l \psi_h + \bar{\psi}_h \psi_l), \end{aligned} \quad (2.17)$$

where ψ_l (ψ_h) denotes a light (heavy) Dirac fermion and ϕ is a heavy real scalar with $m \ll M$. We will obtain an effective action of ψ_l after integrating out heavy fields, where irrelevant operators must include traces of heavy particles and their interactions at high energy scale. Let us calculate the effective action by integrate out ψ_h, ϕ , and show dimension six operators by expanding $1/M^n$. The effective action should be given by

$$e^{iS_{\text{eff}}[\psi_l]} = \int \mathcal{D}\phi \mathcal{D}\psi_h \mathcal{D}\bar{\psi}_h e^{iS[\psi_l, \psi_h, \phi]}, \quad (2.18)$$

and firstly, by integrating out H , it becomes

$$\begin{aligned} &= \int \mathcal{D}\phi \mathcal{D}\psi_h \mathcal{D}\bar{\psi}_h \exp i \left\{ S_{\text{free}}[\psi_l, \phi] + (\bar{\psi}_h - \bar{A} K_0^{-1}) \right. \\ &\quad \left. \times K_0 (\psi_h - K_0^{-1} A) - \bar{A} K_0^{-1} A \right\} \\ &= (\text{Det} K_0) \exp i \left\{ S_{\text{free}}[\psi_l, \phi] - \bar{A} K_0^{-1} A \right\}, \end{aligned} \quad (2.19)$$

where

$$\begin{aligned} K_0^{-1} &= \int \frac{d^4 k}{(2\pi)^4} \frac{1}{\not{p} - M} e^{-ip(x-y)} = -i\mathcal{D}^{(\psi_h)}(x-y), \\ A &= g\phi\psi_l, \quad \bar{A} = g\bar{\psi}_l\phi. \end{aligned} \quad (2.20)$$

The second term in Eq. (2.19) is written by

$$- \bar{A} K_0^{-1} A = -\frac{1}{2} \int d^4 x d^4 y \phi(x) \delta \tilde{K}(x, y) \phi(y) \equiv -\frac{1}{2} \phi \delta \tilde{K} \phi, \quad (2.21)$$

$$\delta \tilde{K}(x, y) \equiv -2ig^2 \bar{\psi}_l(x) \mathcal{D}^{(\psi_h)}(x-y) \psi_l(y). \quad (2.22)$$

Next step is an integration of ϕ , which gives

$$\begin{aligned}
 e^{iS_{\text{eff}}[\psi_l]} &= (\text{Det} K_0) \int \mathcal{D}\phi \exp i \left\{ S_{\text{free}}[\psi_l] - \frac{1}{2} \phi \left(\tilde{K}_0 + \delta \tilde{K} \right) \phi \right\} \\
 &= \left(\frac{\text{Det} K_0}{\text{Det}^{\frac{1}{2}} \left(\tilde{K}_0 + \delta \tilde{K} \right)} \right) e^{iS_{\text{free}}[\psi_l]} \\
 &= \left(\frac{\text{Det} K_0}{\text{Det}^{\frac{1}{2}} \tilde{K}_0} \right) \exp \left\{ iS_{\text{free}}[\psi_l] - \frac{1}{2} \text{Tr} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \left(\tilde{K}_0^{-1} \delta \tilde{K} \right)^n \right\}, \quad (2.23)
 \end{aligned}$$

$$\tilde{K}_0^{-1}(x, y) \equiv \int \frac{d^4 p}{(2\pi)^4} \frac{-1}{p^2 - M^2} e^{-ip(x-y)} = i\mathcal{D}^{(\phi)}(x - y). \quad (2.24)$$

Determinant of K_0 and $\tilde{K}_0$ are cancelled by normalization, so that we finally obtain the effective action of ψ_l as

$$S_{\text{eff}}[\psi_l] = S_{\text{free}}[\psi_l] + \frac{i}{2} \text{Tr} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \left(\tilde{K}_0^{-1} \delta \tilde{K} \right)^n. \quad (2.25)$$

Higher dimensional operators are included in the second term of Eq. (2.25), thus dimension six operators are calculated from the second order of $1/M^n$ expansion. A space integration of $\mathcal{O}(1/M^2)$ gives

$$\begin{aligned}
 & - \frac{i}{4} \int d^4 x d^4 y d^4 z d^4 w \tilde{K}_0^{-1}(x, y) \delta \tilde{K}(y, z) \tilde{K}_0^{-1}(z, w) \delta \tilde{K}(w, x) \\
 &= -ig^4 \int \frac{d^4 k_1}{(2\pi)^4} \frac{d^4 k_2}{(2\pi)^4} \frac{d^4 k_3}{(2\pi)^4} \frac{d^4 k_4}{(2\pi)^4} (2\pi)^4 \delta^4(k_1 - k_2 + k_3 - k_4) \\
 &\times \frac{d^4 p}{(2\pi)^4} \frac{1}{p^2 - M^2} \bar{\psi}_l(k_1) \frac{1}{\not{p} + \not{k}_1 - M} \psi_l(k_2) \frac{1}{(p + k_1 - k_2)^2 - M^2} \\
 &\times \bar{\psi}_l(k_2) \frac{1}{\not{p} + \not{k}_4 - M} \psi_l(k_4), \quad (2.26)
 \end{aligned}$$

and integration of all momenta $p, k_i, (i = 1, \dots, 4)$ with $k_i \ll M$ induces 4-Fermi operators

$$\mathcal{O}_{4\text{F}}(x) = \frac{g^4}{192\pi^2} \frac{1}{M^2} \left[- (\bar{\psi}_l \gamma^\mu \psi_l) (\bar{\psi}_l \gamma_\mu \psi_l) + 2 (\bar{\psi}_l \psi_l) (\bar{\psi}_l \psi_l) \right]. \quad (2.27)$$

These are the dimension six operators in this model. Notice that accurate coefficients are automatically obtained without care of symmetric factors. Other higher order operators can be calculated similarly.

3. Dimension six operators induced from new physics

By using the calculation method represented in the previous section, we concretely calculate coefficients of dimension six operators induced from some candidates of new physics, SUSY, UED, and LH. We obtain coefficients of dimension six operators in QCD by integrating out SUSY and UED particles. We also obtain coefficients of 4-Fermi operators originating from anomalous interactions induced from LH.

3.1. SUSY

Let us calculate coefficients of dimension six operators in QCD when beyond the SM is SUSY. In the SUSY with R-parity, SUSY particles propagate only inside the loop diagrams. Lagrangian of the QCD sector in SUSY SM is given by

$$\begin{aligned}
\mathcal{L} = & \mathcal{L}_{\text{SM}} + \frac{1}{2} \bar{g} (i \not{\partial} - m_{\tilde{g}}) \tilde{g} - \tilde{q}_R^\dagger (\partial^2 + m_{\tilde{q}_R}^2) \tilde{q}_R - \tilde{q}_L^\dagger (\partial^2 + m_{\tilde{q}_L}^2) \tilde{q}_L \\
& + \frac{i}{2} g_s f^{abc} \bar{g}^a \gamma^\mu \tilde{g}^b G_\mu^c - i g_s \sum_q \left(\tilde{q}_L^\dagger \frac{\lambda^a}{2} \overleftrightarrow{\partial}^\mu \tilde{q}_L \right) G_\mu^a - i g_s \sum_q \left(\tilde{q}_R^\dagger \frac{\lambda^a}{2} \overleftrightarrow{\partial}^\mu \tilde{q}_R \right) G_\mu^a \\
& + g_s^2 \sum_q \tilde{q}_L^\dagger \frac{\lambda^a}{2} \frac{\lambda^b}{2} \tilde{q}_L G_\mu^a G^{b\mu} + g_s^2 \sum_q \tilde{q}_R^\dagger \frac{\lambda^a}{2} \frac{\lambda^b}{2} \tilde{q}_R G_\mu^a G^{b\mu} \\
& - \sqrt{2} g_s \sum_q \left(\tilde{q}_L^\dagger \bar{g}_R^a \frac{\lambda^a}{2} q_L + \text{h.c.} \right) + \sqrt{2} g_s \sum_q \left(\tilde{q}_R^\dagger \bar{g}_L^a \frac{\lambda^a}{2} q_R + \text{h.c.} \right), \quad (3.1)
\end{aligned}$$

where $\sum_q$ represents a sum over all flavors. Effective action should be obtained by integrating out $\tilde{q}_L$, $\tilde{q}_R$, and $\tilde{g}$ as

$$e^{iS_{\text{eff}}} = \int \mathcal{D}\tilde{g} \mathcal{D}\tilde{q}_L \mathcal{D}\bar{\tilde{q}}_L \mathcal{D}\tilde{q}_R \mathcal{D}\bar{\tilde{q}}_R e^{iS}. \quad (3.2)$$

S_{eff} includes all possible irrelevant operators. Calculating results are listed in Appendix B.1, where coefficients of dimension six operators, 4-Fermi $\mathcal{O}_{qqqq}$, quark–quark–gluon–gluon $\mathcal{O}_{qqGG}$, and quark–quark–gluon $\mathcal{O}_{qqG}$ are represented. They are operators up to $\mathcal{O}(g_s^4)$, which are useful to estimate phenomenology at LHC [10]. We overview explicit technique to calculate them in the following discussions.

The first step is integrating out $\tilde{q}_R$ as

$$\int \mathcal{D}\tilde{q}_R \mathcal{D}\bar{\tilde{q}}_R e^{iS} = \exp i \left[i \text{Tr}(\log K) + B^\dagger K^{-1} B \right], \quad (3.3)$$

$$\begin{aligned}
K &= K_0 + \delta K, \\
K_0 &= (\partial^\mu \partial_\mu + m_R^2), & \delta K &= ig_s [2G_\mu \partial^\mu + (\partial^\mu G_\mu)] - g_s^2 G_\mu G^\mu, \\
B &= -\sqrt{2}g_s \left(\bar{g}^a \frac{\lambda^a}{2} P_R q \right), & B^\dagger &= -\sqrt{2}g_s \left(\bar{q} P_L \frac{\lambda^a}{2} \tilde{g}^a \right),
\end{aligned} \tag{3.4}$$

where $\text{Tr}(\log K)$ includes some loop diagrams which have external gluon lines (*e.g.* Fig. 1). The second term, $B^\dagger K^{-1} B$, can be expanded by right-handed squark propagator as

$$\begin{aligned}
B^\dagger K^{-1} B &= B^\dagger (1 + K_0^{-1} \delta K)^{-1} K_0^{-1} B \\
&= B^\dagger K_0^{-1} B - B^\dagger K_0^{-1} \delta K K_0^{-1} B + B^\dagger K_0^{-1} \delta K K_0^{-1} \delta K K_0^{-1} B + \dots
\end{aligned} \tag{3.5}$$




Fig. 1. Diagrams of gluon external lines.

We take the expansion up to order of g_s^4 in S_{eff} . Similarly, $\tilde{q}_L$ integration can be performed, and after $\tilde{q}_R, \tilde{q}_L$ integrations, an “effective action” in this stage is given by

$$S^{(\tilde{g})} = \frac{1}{2} \int d^4x d^4y \bar{\tilde{g}}(x)_i^a \left[\tilde{K}_0 + \tilde{K}_I \right]_{xy}^{ab} \tilde{g}(y)_j^b, \tag{3.6}$$

where $\tilde{K}_0$ and $\tilde{K}_I$ are

$$\left[\tilde{K}_0 \right]_{xy}^{ab} \equiv \delta^{ab} \delta^4(x - y) (i\partial_y - m_{\tilde{g}})_{ij}, \tag{3.7}$$

$$\begin{aligned}
\left[\tilde{K}_I \right]_{xy}^{ab} &\equiv ig_s f^{abc} \delta^4(x - y) G_{ij}^c - 4ig_s^2 \sum_{q=q_L, q_R} \bar{q}_j(y) \frac{\lambda^b}{2} \mathcal{D}^{(\tilde{q})}(y - x) \frac{\lambda^a}{2} q_i(x) \\
&- 4ig_s^3 \int d^4z \sum_{q=q_L, q_R} \bar{q}_j(y) \frac{\lambda^b}{2} \mathcal{D}^{(\tilde{q})}(y - z) \{2G_\mu \partial^\mu + (\partial^\mu G_\mu)\}_z \mathcal{D}^{(\tilde{q})}(z - x) \frac{\lambda^a}{2} q_i(x) \\
&- 4ig_s^4 \int d^4z d^4w \sum_{q=q_L, q_R} \bar{q}_j(y) \bar{q}_j(y) \frac{\lambda^b}{2} \mathcal{D}^{(\tilde{q})}(y - z) \{2G_\mu \partial^\mu + (\partial^\mu G_\mu)\}_z \mathcal{D}^{(\tilde{q})}(z - w) \\
&\times \{2G_\mu \partial^\mu + (\partial^\mu G_\mu)\}_w \mathcal{D}^{(\tilde{q})}(w - x) \frac{\lambda^a}{2} q_i(x) + \mathcal{O}(g_s^5),
\end{aligned} \tag{3.8}$$

and

$$\mathcal{D}^{(\tilde{g})}(x-y) = -iK_0^{-1} = \int \frac{d^4k}{(2\pi)^4} \frac{i}{k^2 - m_{\tilde{q}}^2 + i\epsilon} e^{-ik(x-y)}. \quad (3.9)$$

Here i, j denote spinor indexes. Next step is integrating out gluino, and the final effective action is obtained as

$$\begin{aligned} S_{\text{eff}} = & \int d^4x d^4y \alpha(x, y)_{ij}^{ab} \tilde{K}_I(x, y)_{ij}^{ab} \\ & + \int d^4x d^4y d^4z d^4w \beta(x, y, z, w)_{ijkl}^{abcd} \tilde{K}_I(x, y)_{ij}^{ab} \tilde{K}_I(z, w)_{kl}^{cd} + \mathcal{O}(\tilde{K}_I^3), \end{aligned} \quad (3.10)$$

where $\alpha(x, y)$, $\beta(x, y, z, w)$ consist of gluino propagator as

$$\alpha(x, y)_{ij}^{ab} = -\frac{1}{2} \delta^{ab} \mathcal{D}^{(\tilde{g})}(y-x)_{ji}, \quad (3.11)$$

$$\begin{aligned} \beta(x, y, z, w)_{ijkl}^{abcd} = & \frac{i}{8} \delta^{ac} \delta^{bd} \left[C^\dagger \mathcal{D}^{(\tilde{g})}(x-z) \right]_{ik} \left[\mathcal{D}^{(\tilde{g})}(w-y) C^T \right]_{lj} \\ & - \frac{i}{8} \delta^{ad} \delta^{bc} \mathcal{D}^{(\tilde{g})}(w-x)_{li} \mathcal{D}^{(\tilde{g})}(y-z)_{lj}. \end{aligned} \quad (3.12)$$

We can know these results by differentiating interacting parts of the effective action as

$$\begin{aligned} \left. \frac{\delta}{\delta \tilde{K}_I(x, y)_{ij}^{ab}} e^{iS_{\text{eff}}} \right|_{\tilde{K}_I=0} &= i\alpha(x, y)_{ij}^{ab}, \quad (3.13) \\ \left. \frac{\delta}{\delta \tilde{K}_I(z, w)_{kl}^{cd}} \frac{\delta}{\delta \tilde{K}_I(x, y)_{ij}^{ab}} e^{iS_{\text{eff}}} \right|_{\tilde{K}_I=0} &= -\alpha(x, y)_{ij}^{ab} \alpha(z, w)_{kl}^{cd} \\ &+ i\beta(x, y, z, w)_{ijkl}^{abcd} + i\beta(z, w, x, y)_{klij}^{cdab}. \end{aligned} \quad (3.14)$$

In this stage, all $\mathcal{O}_{qqqq}$ and $\mathcal{O}_{qqG}$ at 1-loop level are included in Eq. (3.10) up to the second order of $\tilde{K}_I$.

For example, the 4-Fermi operators all $\mathcal{O}_{qqqq}$ are obtained by picking up $\mathcal{O}(g_s^2)$ order terms from each $\tilde{K}_I$ in $\beta\tilde{K}_I\tilde{K}_I$, which is given by

$$\begin{aligned} & \int d^4x d^4y d^4z d^4w \beta(x, y, z, w)_{ijkl}^{abcd} \tilde{K}_I(x, y)_{ij}^{ab} \tilde{K}_I(z, w)_{kl}^{cd} \\ & \supset \int d^4x d^4y d^4z d^4w \left[\frac{i}{8} \delta^{ac} \delta^{bd} \left[C^\dagger \mathcal{D}^{(\tilde{g})}(x-z) \right]_{ik} \left[\mathcal{D}^{(\tilde{g})}(w-y) C^T \right]_{lj} \right. \\ & \quad \left. - \frac{i}{8} \delta^{ad} \delta^{bc} \mathcal{D}^{(\tilde{g})}(w-x)_{li} \mathcal{D}^{(\tilde{g})}(y-z)_{lj} \right] \\ & \times \left[-4ig_s^2 \bar{q}_j(y) \frac{\lambda^b}{2} \mathcal{D}^{(\tilde{q})}(y-x) \frac{\lambda^a}{2} q_i(x) \right] \left[-4ig_s^2 \bar{q}'_l(w) \frac{\lambda^d}{2} \mathcal{D}^{(\tilde{q}')} (w-z) \frac{\lambda^c}{2} q'_k(z) \right]. \end{aligned} \quad (3.15)$$

The first term is given by

$$\begin{aligned} & -2ig_s^4 \int d^4x d^4y d^4z d^4w \int \frac{d^4k_1}{(2\pi)^4} \frac{d^4k_2}{(2\pi)^4} \frac{d^4k_3}{(2\pi)^4} \frac{d^4k_4}{(2\pi)^4} \frac{d^4p_1}{(2\pi)^4} \frac{d^4p_2}{(2\pi)^4} \frac{d^4p_3}{(2\pi)^4} \frac{d^4p_4}{(2\pi)^4} \\ & \times \left[\bar{q}_j(k_2) \frac{\lambda^b}{2} \frac{\lambda^a}{2} q_i(k_1) \right] \left[\bar{q}'_l(k_4) \frac{\lambda^d}{2} \frac{\lambda^c}{2} q'_k(k_3) \right] \\ & \times \frac{i[C^\dagger(p'_1 + m_{\tilde{g}})]_{ik}}{p_1^2 - m_{\tilde{g}}^2} \frac{i[(p'_2 + m_{\tilde{g}})C^T]_{lj}}{p_2^2 - m_{\tilde{g}}^2} \frac{i}{p_3^2 - m_{\tilde{q}}^2} \frac{i}{p_4^2 - m_{\tilde{q}'}^2} \\ & \times e^{ik_2y} e^{-ik_1x} e^{ik_4x} e^{-ik_3z} e^{-ip_1(x-z)} e^{ip_2(w-y)} e^{-ip_3(y-x)} e^{-ip_4(w-z)}, \end{aligned} \quad (3.16)$$

and by integrating out $x, y, z, w, p_2, p_3, p_4$, this term becomes

$$\begin{aligned} & = -2ig_s^4 \int \frac{d^4k_1}{(2\pi)^4} \frac{d^4k_2}{(2\pi)^4} \frac{d^4k_3}{(2\pi)^4} \frac{d^4k_4}{(2\pi)^4} \delta^4(k_1 - k_2 + k_3 - k_4) \\ & \times \left[\bar{q}_j(k_2) \frac{\lambda^b}{2} \frac{\lambda^a}{2} q_i(k_1) \right] \left[\bar{q}'_l(k_4) \frac{\lambda^d}{2} \frac{\lambda^c}{2} q'_k(k_3) \right] \\ & \times \left[A(C^\dagger \gamma^\mu)_{ik} (\gamma^\nu C^T)_{lj} + B(C^\dagger)_{ik} (C^T)_{lj} \right], \end{aligned} \quad (3.17)$$

$$A = \int \frac{d^4p_1}{(2\pi)^4} \frac{p_{1\mu}(p_1 + k_1 - k_2)_\nu}{(p_1^2 - m_{\tilde{g}}^2)[(p_1 + k_1 - k_2)^2 - m_{\tilde{g}}^2][(p_1 + k_1)^2 - m_{\tilde{q}}^2][(p_1 - k_1)^2 - m_{\tilde{q}'}^2]}, \quad (3.18)$$

$$B = \int \frac{d^4p_1}{(2\pi)^4} \frac{m_{\tilde{g}}^2}{(p_1^2 - m_{\tilde{g}}^2)[(p_1 + k_1 - k_2)^2 - m_{\tilde{g}}^2][(p_1 + k_1)^2 - m_{\tilde{q}}^2][(p_1 - k_1)^2 - m_{\tilde{q}'}^2]}. \quad (3.19)$$

Here A, B are Feynman parameter integral, and they become

$$A \rightarrow -i \frac{6}{192\pi^2} f_1(m_{\tilde{q}}, m_{\tilde{q}}), \quad B \rightarrow i \frac{12}{192\pi^2} f_2(m_{\tilde{q}}, m_{\tilde{q}}), \quad (3.20)$$

when k_i , ($i = 1, \dots, 4$) are much smaller than masses of squarks and gluino. f_1, f_2 are shown in Appendix B.1, and the spinor can be rearranged by Fierz transformation of Eqs. (A.1) and (A.2) in Appendix A. Necessary Fierz transformations and color factors are shown in Appendix A.

We can summarize all 4-Fermi operators as separating color singlet $\mathcal{O}_{qqqq}^{(1)}$ or color octet $\mathcal{O}_{qqqq}^{(8)}$, which is shown in Appendix B.1.1. When their chiralities are (LL)(LL) or (RR)(RR) (L: left-handed, R: right-handed), the 4-Fermi operators are given by

$$\mathcal{O}_{qqqq}^{(1)} = \frac{12}{192\pi^2} g_s^4 \left[\frac{2}{9} (f_1 + f_2) \right] (\bar{q}\gamma^\mu q) (\bar{q}'\gamma^\mu q') , \quad (3.21)$$

$$\mathcal{O}_{qqqq}^{(8)} = \frac{12}{192\pi^2} g_s^4 \left[-\frac{1}{3} f_1 - \frac{7}{6} f_2 \right] \left(\bar{q}\gamma^\mu \frac{\lambda^a}{2} q \right) \left(\bar{q}'\gamma^\mu \frac{\lambda^a}{2} q' \right) . \quad (3.22)$$

On the other hand, when their chiralities are (LL)(RR) or (RR)(LL), the 4-Fermi operators are given by

$$\mathcal{O}_{qqqq}^{(1)} = \frac{12}{192\pi^2} g_s^4 \left[\frac{2}{9} (-f_1 + f_2) \right] (\bar{q}\gamma^\mu q) (\bar{q}'\gamma^\mu q') , \quad (3.23)$$

$$\mathcal{O}_{qqqq}^{(8)} = \frac{12}{192\pi^2} g_s^4 \left[-\frac{7}{6} f_1 - \frac{1}{3} f_2 \right] \left(\bar{q}\gamma^\mu \frac{\lambda^a}{2} q \right) \left(\bar{q}'\gamma^\mu \frac{\lambda^a}{2} q' \right) . \quad (3.24)$$

As for $\mathcal{O}_{qqG}$, vertex originates from two parts, $\alpha \tilde{K}_I$ and $\beta \tilde{K}_I \tilde{K}_I$, as

$$\begin{aligned} & \int d^4x d^4y \alpha(x, y)_{ij}^{ab} \tilde{K}_I(x, y)_{ij}^{ab} \supset \int d^4x d^4y \left[-\frac{1}{2} \delta^{ab} \mathcal{D}^{(\bar{g})}(y-x)_{ji} \right] \\ & \times \left[-4ig_s^3 \int d^4z \sum_{q=q_L, q_R} \bar{q}_j(y) \frac{\lambda^b}{2} \mathcal{D}^{(\bar{q})}(y-z) \{2G_\mu \partial^\mu + (\partial^\mu G_\mu)\}_z \right. \\ & \left. \times \mathcal{D}^{(\bar{q})}(z-x) \frac{\lambda^a}{2} q_i(x) \right] , \end{aligned} \quad (3.25)$$

$$\begin{aligned} & \int d^4x d^4y d^4z d^4w \beta(x, y, z, w)_{ijkl}^{abcd} \tilde{K}_I(x, y)_{ij}^{ab} \tilde{K}_I(z, w)_{kl}^{cd} \\ & \supset 2 \int d^4x d^4y d^4z d^4w \left[\frac{i}{8} \delta^{ac} \delta^{bd} \left[C^\dagger \mathcal{D}^{(\bar{g})}(x-z) \right]_{ik} \left[\mathcal{D}^{(\bar{g})}(w-y) C^T \right]_{lj} \right. \\ & \left. - \frac{i}{8} \delta^{ad} \delta^{bc} \mathcal{D}^{(\bar{g})}(w-x)_{li} \mathcal{D}^{(\bar{g})}(y-z)_{lj} \right] \\ & \times \left[ig_s f^{abc} \delta^4(x-y) G_{ij}^c \right] \left[-4ig_s^2 \sum_{q=q_L, q_R} \bar{q}_j(y) \frac{\lambda^b}{2} \mathcal{D}^{(\bar{q})}(y-x) \frac{\lambda^a}{2} q_i(x) \right] . \end{aligned} \quad (3.26)$$

Equations (3.25) and (3.26) include not only dimension four operators but also all higher dimensional operators such as dimension six operator. Higher dimensional operators have been obtained by expanding the full operator by $k^2 \ll \Lambda^2$, where k^μ and Λ denote the momentum of the SM particles and SUSY particles, respectively. Anyhow, we can obtain all $\mathcal{O}_{qqG}$ in the similar calculations as 4-Fermi operators, which is shown Appendix B.1.2.

For $\mathcal{O}_{qqGG}$, they can be also obtained in the same manner. $\mathcal{O}_{qqGG}$ contains in Eq.(3.10), and there are two contributions in the first order of $\tilde{K}_I$ as

$$\begin{aligned} & \int d^4x d^4y \alpha(x, y)_{ij}^{ab} \tilde{K}_I(x, y)_{ij}^{ab} \\ & \supset \int d^4x d^4y \alpha(x, y)_{ij}^{ab} 4g_s^2 \int d^4z \bar{q}_j(y) \frac{\lambda^a}{2} i\mathcal{D}^{(\tilde{q})}(y-z) [-g_s^2 G_\mu G^\mu]_z \\ & \times i\mathcal{D}^{(\tilde{q})}(z-x) \frac{\lambda^a}{2} q_i(x), \end{aligned} \quad (3.27)$$

$$\begin{aligned} & \int d^4x d^4y \alpha(x, y)_{ij}^{ab} \tilde{K}_I(x, y)_{ij}^{ab} \\ & \supset \int d^4x d^4y d^4z d^4w \alpha(x, y)_{ij}^{ab} (-4g_s^2) \bar{q}_j(y) \frac{\lambda^a}{2} i\mathcal{D}^{(\tilde{q})}(y-z) i g_s [2G^\mu \partial_\mu + (\partial_\mu G^\mu)]_z \\ & \times i\mathcal{D}^{(\tilde{q})}(z-w) i g_s [2G^\mu \partial_\mu + (\partial_\mu G^\mu)]_w i\mathcal{D}^{(\tilde{q})}(w-x) \frac{\lambda^a}{2} q_i(x). \end{aligned} \quad (3.28)$$

Similarly, there is one contribution in the second order of $\tilde{K}_I$, which is shown as

$$\begin{aligned} & \int d^4x d^4y d^4z d^4w \beta(x, y, z, w)_{ijkl}^{abcd} \tilde{K}_I(x, y)_{ij}^{ab} \tilde{K}_I(z, w)_{kl}^{cd} \\ & \supset 2 \int d^4x d^4y d^4z d^4w \beta(x, y, z, w)_{ijkl}^{abcd} \left[i g_s f^{abe} \tilde{G}_{ij}^e(x) \delta^4(x-y) \right] \\ & \times \int d^4z' 4g_s^2 \bar{q}_l(z) \frac{\lambda^d}{2} i\mathcal{D}^{(\tilde{q})}(z-z') \{ i g_s (2G^\mu \partial_\mu + (\partial_\mu G^\mu)) \}_{z'} \\ & \times i\mathcal{D}^{(\tilde{q})}(z'-w) \frac{\lambda^c}{2} q_k(w). \end{aligned} \quad (3.29)$$

There is one contribution in the third order in Eq. (3.10), $\gamma \tilde{K}_I \tilde{K}_I \tilde{K}_I$, where γ is given by

$$\begin{aligned}
\gamma(x, y, z, w, u, v)_{ijklmn}^{abcdef} = & \frac{1}{8} \left\{ \delta^{ad} \delta^{be} \delta^{cf} \mathcal{D}^{(\tilde{g})}(w-x)_{li} \right. \\
& \times \mathcal{D}^{(\tilde{g})}(y-u)_{jm} \mathcal{D}^{(\tilde{g})}(v-z)_{nk} \\
& + \delta^{af} \delta^{bc} \delta^{ed} \mathcal{D}^{(\tilde{g})}(v-x)_{ni} \mathcal{D}^{(\tilde{g})}(y-x)_{jk} \mathcal{D}^{(\tilde{g})}(w-u)_{lm} \\
& - \delta^{ac} \delta^{be} \delta^{df} [C^\dagger \mathcal{D}^{(\tilde{g})}(z-x)]_{ki} \mathcal{D}^{(\tilde{g})}(y-u)_{jm} [\mathcal{D}^{(\tilde{g})}(v-w) C^T]_{nl} \\
& - \delta^{ac} \delta^{bf} \delta^{de} \left[C^\dagger \mathcal{D}^{(\tilde{g})}(x-z) \right]_{ik} \left[\mathcal{D}^{(\tilde{g})}(v-y) C^T \right]_{nj} \mathcal{D}^{(\tilde{g})}(w-u)_{lm} \\
& - \delta^{ad} \delta^{bf} \delta^{ce} \mathcal{D}^{(\tilde{g})}(w-x)_{li} \left[\mathcal{D}^{(\tilde{g})}(y-v) C^T \right]_{jn} \left[C^\dagger \mathcal{D}^{(\tilde{g})}(u-z) \right]_{mk} \\
& - \delta^{ae} \delta^{bc} \delta^{fd} \left[C^\dagger \mathcal{D}^{(\tilde{g})}(u-x) \right]_{mi} \mathcal{D}^{(\tilde{g})}(y-z)_{jk} \left[\mathcal{D}^{(\tilde{g})}(w-v) C^T \right]_{ln} \\
& - \delta^{ae} \delta^{bd} \delta^{fc} \left[C^\dagger \mathcal{D}^{(\tilde{g})}(x-u) \right]_{im} \left[\mathcal{D}^{(\tilde{g})}(y-w) C^T \right]_{lj} \mathcal{D}^{(\tilde{g})}(v-z)_{nk} \\
& \left. - \delta^{af} \delta^{bd} \delta^{ec} \mathcal{D}^{(\tilde{g})}(v-x)_{ni} \left[\mathcal{D}^{(\tilde{g})}(y-w) C^T \right]_{jl} \left[C^\dagger \mathcal{D}^{(\tilde{g})}(z-u) \right]_{km} \right\}, \quad (3.30)
\end{aligned}$$

and the third order is shown as

$$\begin{aligned}
& \int d^4x \cdots d^4v \gamma(x, y, z, w, u, v)_{ijklmn}^{abcdef} \tilde{K}_I(x, y)_{ij}^{ab} \tilde{K}_I(z, w)_{kl}^{cd} \tilde{K}_I(u, v)_{mn}^{ef} \\
& \supset \int d^4x \cdots d^4v \gamma(x, y, z, w, u, v)_{ijklmn}^{abcdef} \left[(-4g_s^2) \bar{q}_j(y) \frac{\lambda^b}{2} i \mathcal{D}^{(\tilde{q})}(y-x) \frac{\lambda^a}{2} q_i(x) \right] \\
& \times \left[ig_s f^{cdg} G^g(z)_{kl} \delta^4(z-w) \right] \left[ig_s f^{efh} G^h(u)_{mn} \delta^4(u-v) \right]. \quad (3.31)
\end{aligned}$$

Although γ has eight terms in total, they are all the same in Eq. (3.31) since each term of γ corresponds to statistic factor in Feynman diagram. Notice again that we do not care about a statistic factor in each operator since it is automatically included in α, β, γ . We can obtain all $\mathcal{O}_{qqGG}$ induced from SUSY which is shown in Appendix B.1.3.

3.2. UED

Next, we estimate QCD dimension six operators induced from UED. The UED has KK-parity so that KK particles can propagate only inside loop processes. As the SUSY case, we can calculate dimension six operators by integrating out KK particles.

After dimensional reduction of the fifth dimensional compactified space of S^1/Z_2 , UED Lagrangian in the 4-dimensional space-time is given by

$$\begin{aligned}
\mathcal{L} &= \mathcal{L}_{\text{SM}} + \mathcal{L}_{q^{(n)}} + \mathcal{L}_{G^{(n)}} + \mathcal{L}_{G_5^{(n)}} , \\
\mathcal{L}_{q^{(n)}} &= \sum_n \left[\overline{q_L^{(n)}} i \left(\not{\partial} + i g_s \not{G} - m_L^{(n)} \right) q_L^{(n)} + \overline{q_R^{(n)}} i \left(\not{\partial} + i g_s \not{G} + m_R^{(n)} \right) q_R^{(n)} \right. \\
&\quad \left. - g_s \left(\overline{q} G^{(n)} P_L q^{(n)} + \overline{q} i \gamma^5 G_5^{(n)} P_R q^{(n)} \right) - (L \longleftrightarrow R) \right] + \dots , \\
\mathcal{L}_{G^{(n)}} &= \sum_n \left[-\frac{1}{4} \left(\partial_\mu G_\nu^{(n)a} - \partial_\nu G_\mu^{(n)a} \right)^2 + \frac{1}{2} m_g^{(n)2} G_\mu^{(n)a} G^{(n)a\mu} \right. \\
&\quad - \frac{1}{2} g_s f^{abc} \left(\partial_\mu G_\nu^a - \partial_\nu G_\mu^a \right) G^{(n)b\mu} G^{(n)c\nu} - \frac{1}{2} g_s f^{abc} \left(\partial_\mu G_\nu^{(n)a} - \partial_\nu G_\mu^{(n)a} \right) G^{(n)b\mu} G^{(n)c\nu} \\
&\quad - \frac{1}{2} g_s f^{abc} \partial_\mu G_\nu^{(n)a} - \partial_\nu G_\mu^{(n)a} G^{(n)b\mu} G^{(n)c\nu} - \frac{1}{4} f^{abc} f^{ade} \left\{ 2 G_\mu^{(0)b} G_\nu^{(0)c} G^{(n)d\mu} G^{(n)e\mu} \right. \\
&\quad \left. + \left(G_\mu^{(0)b} G_\nu^{(n)c} + G_\mu^{(n)b} G_\nu^{(0)c} \right) \left(G^{(0)d\mu} G^{(n)e\mu} + G^{(n)d\mu} G^{(0)e\mu} \right) \right\} \dots , \\
\mathcal{L}_{G_5^{(n)}} &= \sum_n \left[\frac{1}{2} \partial_\mu G_5^{(n)a} \partial^\mu G_5^{(n)a} - \frac{1}{2} m_5^{(n)2} G_5^{(n)a} G_5^{(n)a} \right. \\
&\quad \left. + g_s f^{abc} \left(m_g G_\nu^{(n)a} + \partial_\nu G_5^{(n)a} \right) G^{(n)b5} G^{(n)c\nu} - \frac{1}{2} g_s^2 f^{abc} f^{ade} G_5^{(n)b} G_\nu^c G^{(n)d5} G^{(n)e\nu} \right] + \dots ,
\end{aligned}$$

where

$$\begin{aligned}
m_L^{(n)} &= \frac{n}{R} + \delta m_L , & m_R^{(n)} &= \frac{n}{R} + \delta m_R , \\
m_g^{(n)} &= \frac{n}{R} + \delta m_g , & m_5^{(n)} &= \frac{n}{R} + \delta m_5 .
\end{aligned}$$

Here we take a 't Hooft–Feynman gauge fixing, and $m_L^{(n)}$, $m_R^{(n)}$, $m_g^{(n)}$, $m_5^{(n)}$ are SU(2) doublet KK quark mass, SU(2) singlet KK quark mass, KK gluon mass, and KK scalar (fifth dimensional component of KK gluon) mass with each radiative correction, respectively. At a tree level, these KK particles are degenerate in a minimal UED, but there is a slight difference between $m_L^{(n)}$ and $m_R^{(n)}$ when we consider radiative corrections. So here we show general effective operators by using these general parameters. The effective operators of S_{eff} in UED can be calculated by the similar technique of the previous subsection, and the results are shown in Appendix B.2.

Here we overview this calculation. By integrating out KK quarks, KK scalars, and KK gluons, the effective action becomes

$$S_{\text{eff}} = \tilde{S} + \int d^4x d^4y \alpha(x, y)_{\mu\nu}^{ab} K_I(x, y)_{\mu\nu}^{ab} \\ + \int d^4x d^4y d^4z d^4w \beta(x, y, z, w)_{\mu\nu\rho\sigma}^{abcd} K_I(x, y)_{\mu\nu}^{ab} K_I(z, w)_{\rho\sigma}^{cd} + \cdots, \quad (3.32)$$

where $\tilde{S}$ does not include KK gluons that is given by

$$\begin{aligned} \tilde{S} = & -g_s^3 \int d^4x d^4y d^4z \mathcal{D}^{(s)}(x - y) \\ & \times \left[\bar{q}_L(x) \frac{\lambda^a}{2} \mathcal{D}^{(L)}(x - z) G(z) \mathcal{D}^{(L)}(z - y) \frac{\lambda^a}{2} q_L(y) + (L \leftrightarrow R) \right] \\ & + i \int d^4x d^4y d^4z_1 d^4z_2 d^4z_3 \delta^4(x - y) \\ & \times \left[g_s f^{bcf} G^{f\mu}(z_1) \partial_{z_1\mu} \delta^4(z_1 - z_2) + i g_s^2 \right. \\ & \times \left. \left\{ \bar{q}_L(z_1) \frac{\lambda^b}{2} \mathcal{D}^{(L)}(z_1 - z_2) \frac{\lambda^c}{2} q_L(z) + (L \leftrightarrow R) \right\} \right] \\ & \times \left[\mathcal{D}^{(s)}(x - z_1) \delta^{ab} \right] \\ & \times \left[g_s f^{deg} G^{g\nu}(z_3) \partial_{z_3\nu} \delta^4(z_3 - y) + i g_s^2 \right. \\ & \times \left. \left\{ \bar{q}_L(z_3) \frac{\lambda^b}{2} \mathcal{D}^{(L)}(z_3 - y) \frac{\lambda^c}{2} q_L(y) + (L \leftrightarrow R) \right\} \right] \\ & \times \left[\mathcal{D}^{(s)}(z_2 - z_3) \delta^{cd} \right]. \end{aligned} \quad (3.33)$$

Here $\mathcal{D}^{(L)}$, $\mathcal{D}^{(R)}$, and $\mathcal{D}^{(s)}$ are propagators of KK quarks and KK scalars as

$$\mathcal{D}^{(L)} = \int \frac{d^4p}{(2\pi)^4} \frac{i}{\not{p} - m_L^{(n)}} e^{-ip(x-y)}, \quad (3.34)$$

$$\mathcal{D}^{(R)} = \int \frac{d^4p}{(2\pi)^4} \frac{i}{\not{p} + m_R^{(n)}} e^{-ip(x-y)}, \quad (3.35)$$

$$\delta^{ab} \mathcal{D}^{(s)} = \delta^{ab} \int \frac{d^4p}{(2\pi)^4} \frac{i}{p^2 - m_5^{(n)2}} e^{-ip(x-y)}, \quad (3.36)$$

respectively. And α, β, K_I are given by

$$\alpha(x, y)_{\mu\nu}^{ab} = \frac{1}{2} \mathcal{D}^{(g)}(x - y) \delta^{ab} g_{\mu\nu} \equiv \frac{1}{2} \delta^{ab} g_{\mu\nu} \int \frac{d^4 p}{(2\pi)^4} \frac{-i}{p^2 - m_g^2} e^{-i(x-y)}, \quad (3.37)$$

$$\begin{aligned} \beta(x, y, z, w)_{\mu\nu\rho\sigma}^{abcd} &= \frac{i}{8} \left[\mathcal{D}^{(g)}(x - z) \mathcal{D}^{(g)}(y - w) \delta^{ac} \delta^{bd} g_{\mu\rho} g_{\nu\sigma} \right. \\ &\quad \left. + \mathcal{D}^{(g)}(x - w) \mathcal{D}^{(g)}(y - z) \delta^{ad} \delta^{bc} g_{\mu\sigma} g_{\nu\rho} \right], \end{aligned} \quad (3.38)$$

$$\begin{aligned} K_I(x, y)_{\mu\nu}^{ab} &= -2g_s f^{abc} [\partial^\mu G^{c\nu}(x) + G^{c\mu}(x) \partial_x^\nu - g^{\mu\nu} G^{c\sigma}(x) \partial_{x\sigma}] \delta^4(x - y) \\ &+ 2ig_s^2 \left[\bar{q}_L(x) \gamma^\mu \frac{\lambda^a}{2} \mathcal{D}^{(L)}(x - y) \gamma^\nu \frac{\lambda^b}{2} q_L(y) + (L \leftrightarrow R) \right] \\ &+ 2g_s^3 \int d^4 z \left[\bar{q}_L(x) \gamma^\mu \frac{\lambda^a}{2} \mathcal{D}^{(L)}(x - z) G(z) \mathcal{D}^{(L)}(z - y) \gamma^\nu \frac{\lambda^b}{2} q_L(y) + (L \leftrightarrow R) \right] \\ &- ig_s^4 \int d^4 z d^4 w \\ &\times \left[\left(\bar{q}_L(x) \gamma^\mu \frac{\lambda^a}{2} \mathcal{D}^{(L)}(x - z) \frac{\lambda^c}{2} q_L(z) \right) \mathcal{D}^{(s)}(z - w) \left(\bar{q}_L(y) \gamma^\nu \frac{\lambda^b}{2} \mathcal{D}^{(L)}(y - w) \frac{\lambda^c}{2} q_L(w) \right) \right. \\ &+ \left(\bar{q}_L(z) \frac{\lambda^c}{2} \mathcal{D}^{(L)}(z - x) \gamma^\mu \frac{\lambda^a}{2} q_L(x) \right) \mathcal{D}^{(s)}(z - w) \left(\bar{q}_L(w) \frac{\lambda^c}{2} \mathcal{D}^{(L)}(w - y) \gamma^\nu \frac{\lambda^b}{2} q_L(y) \right) \\ &- 2 \left(\bar{q}_L(x) \gamma^\mu \frac{\lambda^a}{2} \mathcal{D}^{(L)}(x - z) \frac{\lambda^c}{2} q_L(z) \right) \mathcal{D}^{(s)}(z - w) \left(\bar{q}_L(w) \frac{\lambda^c}{2} \mathcal{D}^{(L)}(w - y) \gamma^\nu \frac{\lambda^b}{2} q_L(y) \right) \\ &- \left(\bar{q}_L(x) \gamma^\mu \frac{\lambda^a}{2} \mathcal{D}^{(L)}(x - z) \frac{\lambda^c}{2} q_L(z) \right) \mathcal{D}^{(s)}(z - w) \left(\bar{q}_R(y) \gamma^\nu \frac{\lambda^b}{2} \mathcal{D}^{(R)}(y - w) \frac{\lambda^c}{2} q_R(w) \right) \\ &- \left(\bar{q}_L(z) \frac{\lambda^c}{2} \mathcal{D}^{(L)}(z - x) \gamma^\mu \frac{\lambda^a}{2} q_L(x) \right) \mathcal{D}^{(s)}(z - w) \left(\bar{q}_R(w) \frac{\lambda^c}{2} \mathcal{D}^{(R)}(w - y) \gamma^\nu \frac{\lambda^b}{2} q_R(y) \right) \\ &+ \left(\bar{q}_L(x) \gamma^\mu \frac{\lambda^a}{2} \mathcal{D}^{(L)}(x - z) \frac{\lambda^c}{2} q_L(z) \right) \mathcal{D}^{(s)}(z - w) \left(\bar{q}_R(w) \frac{\lambda^c}{2} \mathcal{D}^{(R)}(w - y) \gamma^\nu \frac{\lambda^b}{2} q_R(y) \right) \\ &+ \left(\bar{q}_L(w) \frac{\lambda^c}{2} \mathcal{D}^{(L)}(w - y) \gamma^\nu \frac{\lambda^b}{2} q_L(y) \right) \mathcal{D}^{(s)}(z - w) \left(\bar{q}_R(x) \gamma^\mu \frac{\lambda^a}{2} \mathcal{D}^{(R)}(x - z) \frac{\lambda^c}{2} q_R(z) \right) \\ &\left. + (L \leftrightarrow R) \right], \end{aligned} \quad (3.39)$$

respectively. Here $\partial_x^\nu, \partial_{x\sigma}$ in the first line of Eq. (3.39) means derivatives of KK gluons.

Dimension six operators induced from UED can be also obtained in the same manner as in SUSY, which is discussed in Sec. 3.1. In the UED case, the analysis is also made up to the order of $\mathcal{O}(g_s^4)$. We show that $\mathcal{O}_{qqqq}$ comes from the terms $(K_I)^0$, $(K_I)^1$, and $(K_I)^2$ in Eq. (3.32). Similarly, we can obtain $\mathcal{O}_{qqG}$ from the terms $(K_I)^1$, $(K_I)^2$, and $(K_I)^3$. As for $\mathcal{O}_{qqGG}$, it comes from the terms $(K_I)^1$, $(K_I)^2$, $(K_I)^3$, and $(K_I)^4$. Explicit forms of $\mathcal{O}_{qqqq}$, $\mathcal{O}_{qqG}$, and $\mathcal{O}_{qqGG}$ are shown in Appendix B.2.

3.3. Little Higgs

In this subsection, we calculate dimension six operators with anomalous coupling in LH model. The Lagrangian is given by

$$\begin{aligned}\mathcal{L}_{\text{int}} = & -g_s \bar{t} G t - \frac{2}{3} e \bar{t} A t - \frac{g}{\sqrt{2}} \cos \beta (\bar{b} W P_L t + \text{h.c.}) \\ & - \frac{g}{\cos \theta_W} \bar{t} Z \left(-\frac{2}{3} \sin^2 \theta_W + \frac{1}{2} \cos^2 \beta P_L \right) t, \quad (3.40)\end{aligned}$$

where β and m_t are $\tan^{-1} \frac{\lambda_1^2}{\lambda_1^2 + \lambda_2^2} \frac{v}{f}$ and $\frac{\sqrt{\lambda_1 \lambda_2}}{\lambda_1^2 + \lambda_2^2}$, respectively. The f denotes the VEV of the Little Higgs. When we take $\lambda_1 \simeq \lambda_2 \simeq 1$, we can estimate $\cos \beta \simeq 1 - \frac{v^2}{2f^2}$. Only 4-Fermi operators have non-standard effects originated from W , Z -boson exchanges in the electroweak interaction. Then, below an energy scale of M_W , we can obtain $\mathcal{O}_{qqqq}$ as

$$\begin{aligned}\mathcal{O}^{\text{LH}} = & \frac{g^2}{2} \cos^2 \beta \frac{1}{k^2 - M_W^2} (\bar{t} \gamma^\mu P_L b) (\bar{b} \gamma^\mu P_L t) \\ & + \frac{g^2}{3} \tan^2 \theta_W \cos^2 \beta \frac{1}{k^2 - M_Z^2} (\bar{q} \gamma^\mu P_L q) (\bar{t} \gamma^\mu P_L t). \quad (3.41)\end{aligned}$$

They are useful for estimating an evidence of the LH.

4. Summary

It is important to obtain effective operators by integrating out high energy degrees of freedom in physics. In a quantum field theory, we can obtain effective Lagrangian by integrating out high energy momentum and heavy particles. In this paper, we have shown a general method of calculating accurate irrelevant effective operators in a scattering process without use of EOMs. By using this method, for example, we have represented coefficients of dimension six operators induced from SUSY, UED, and LH. We have shown coefficients of dimension six operators in QCD by integrating out sparticles (KK particles) in SUSY (UED). We have also shown coefficients of 4-Fermi operators originating from anomalous interactions in LH. They are promising candidates of beyond the SM, and our analyses will shed lights on the search of beyond the SM.

Our method can also give an effective action by integrating out high momentum degrees of freedom of massless particles. In this case, an effective Lagrangian have, in general, non-local interactions. Anyhow, our method can apply to other quantum field theories in any dimensions, so that we believe this technique is very useful in a lot of researches in physics.

This work is partially supported by the scientific grants of the Ministry of Education and Science, Japan, Nos. 20540272, 20039006, and 20025004.

Appendix A

Fierz transformation and color factor

Here let us summarize some formulas which are useful for calculations. Fierz transformation of γ -matrix in Eq. (3.17) shows

$$\left(C^\dagger \gamma^\mu\right)_{ik} (\gamma^\nu C^T)_{lj} = -\frac{1}{2} (\gamma^\mu)_{ji} (\gamma_\mu)_{lk} - \frac{1}{2} (\gamma^\mu \gamma^5)_{ji} (\gamma_\mu \gamma^5)_{lk}, \quad (\text{A.1})$$

$$\left(C^\dagger\right)_{ik} (C^T)_{lj} = \frac{1}{4} (\gamma^\mu)_{ji} (\gamma_\mu)_{lk} - \frac{1}{4} (\gamma^\mu \gamma^5)_{ji} (\gamma_\mu \gamma^5)_{lk}. \quad (\text{A.2})$$

We should notice that neither scalar nor pseudo-scalar components appear by the Fierz transformation, since spinor components, (i, j, k, l) , always have the same chirality in each set of (i, j) and (k, l) .

The color factor becomes

$$\left(\frac{\lambda^b}{2}\right)_{jn} \left(\frac{\lambda^a}{2}\right)_{ni} \left(\frac{\lambda^b}{2}\right)_{lm} \left(\frac{\lambda^a}{2}\right)_{mk} = \frac{2}{9} \delta_{lk} \delta_{jk} - \frac{1}{3} \left(\frac{\lambda^a}{2}\right)_{lk} \left(\frac{\lambda^a}{2}\right)_{jk} \quad (\text{A.3})$$

from Fierz transformation of spinors. Next formulas are useful for the second term of Eq. (3.15), whose spinor and color factor are different from those of the first term. The spinor is given by

$$(\gamma^\mu)_{li} (\gamma_\mu)_{jk} = -\frac{1}{2} (\gamma^\mu)_{ji} (\gamma_\mu)_{lk} + \frac{1}{2} (\gamma^\mu \gamma^5)_{ji} (\gamma_\mu \gamma^5)_{lk}, \quad (\text{A.4})$$

$$\delta_{li} \delta_{jk} = \frac{1}{4} (\gamma^\mu)_{ji} (\gamma_\mu)_{lk} + \frac{1}{4} (\gamma^\mu \gamma^5)_{ji} (\gamma_\mu \gamma^5)_{lk}, \quad (\text{A.5})$$

and the color factor becomes

$$\left(\frac{\lambda^b}{2}\right)_{jn} \left(\frac{\lambda^a}{2}\right)_{ni} \left(\frac{\lambda^a}{2}\right)_{lm} \left(\frac{\lambda^b}{2}\right)_{mk} = \frac{2}{9} \delta_{lk} \delta_{jk} + \frac{7}{6} \left(\frac{\lambda^a}{2}\right)_{lk} \left(\frac{\lambda^a}{2}\right)_{jk}, \quad (\text{A.6})$$

while A, B are the same as the first term.

Appendix B

Explicit coefficients of dimension six operators

We define the following functions to write coefficients in terms of linear combinations of them

$$\begin{aligned}
 \alpha(a, b, m, n) &\equiv \int_0^1 dx \frac{x^n}{(x(a^2 - b^2) + b^2)^m}, \\
 \beta(a, b, c, m, n, l) &\equiv \int_0^1 dx \int_x^1 dy \frac{x^n y^l}{(x(c^2 - b^2) + y(b^2 - a^2) + a^2)^m}, \\
 \gamma(a, b, c, m, n, l, r) &\equiv \int_0^1 dx \int_x^1 dy \int_y^1 dz \\
 &\quad \times \frac{x^n y^l z^r}{(x(c^2 - a^2) + y(a^2 - c^2) + z(c^2 - b^2) + b^2)^m}, \\
 \delta(a, b, c, m, n, l, r) &\equiv \int_0^1 dx \int_x^1 dy \int_y^1 dz \frac{x^n y^l z^r}{(x(c^2 - a^2) + z(a^2 - b^2) + b^2)^m}.
 \end{aligned} \tag{B.1}$$

Appendix B.1. SUSY

In SUSY SM, dimension six operators are written as follows,

$$\begin{aligned}
 \int d^4x \mathcal{O}_{qqqq}^{(1)}(x) &= \frac{12g_s^4}{192\pi^2} \int \frac{d^4k_1}{(2\pi)^4} \frac{d^4k_2}{(2\pi)^4} \frac{d^4k_3}{(2\pi)^4} \frac{d^4k_4}{(2\pi)^4} (2\pi)^4 \delta^4(-k_1 + k_2 - k_3 + k_4) \\
 &\times \sum_{i,j=L,R} C_{ij} (\bar{q}(k_1) \gamma^\mu P_i q(k_2)) (\bar{q}'(k_3) \gamma_\mu P_j q'(k_4)), \tag{B.2}
 \end{aligned}$$

$$\begin{aligned}
 \int d^4x \mathcal{O}_{qqqq}^{(8)}(x) &= \frac{12g_s^4}{192\pi^2} \int \frac{d^4k_1}{(2\pi)^4} \frac{d^4k_2}{(2\pi)^4} \frac{d^4k_3}{(2\pi)^4} \frac{d^4k_4}{(2\pi)^4} (2\pi)^4 \delta^4(-k_1 + k_2 - k_3 + k_4) \\
 &\times \sum_{i,j=L,R} D_{ij} (\bar{q} T^a \gamma^\mu P_i q) (\bar{q}' T^a \gamma_\mu P_j q'), \tag{B.3}
 \end{aligned}$$

$$\begin{aligned}
 \int d^4x \mathcal{O}_{qqG}(x) &= \frac{g_s^3}{96\pi^2} \int \frac{d^4k_1}{(2\pi)^4} \frac{d^4k_2}{(2\pi)^4} \frac{d^4k_3}{(2\pi)^4} (2\pi)^4 \delta^4(-k_1 + k_2 + k_3) \bar{q}(k_2) \\
 &\times T^a E_{L,R}^\mu G_\mu^a(k_3) P_{L,R} q(k_1), \tag{B.4}
 \end{aligned}$$

$$\int d^4x \mathcal{O}_{qqGG}(x) = \frac{g_s^4}{192\pi^2} \int \frac{d^4k_1}{(2\pi)^4} \frac{d^4k_2}{(2\pi)^4} \frac{d^4k_3}{(2\pi)^4} \frac{d^4k_4}{(2\pi)^4} (2\pi)^4 \delta^4(-k_1+k_2+k_3+k_4) \\ \times \bar{q}(k_1) \left[F_{L,R}^{\mu\nu} \delta^{ab} + H_{L,R}^{\mu\nu} T^a T^b \right] G_\mu^a(k_2) G_\nu^b(k_3) P_{L,R} q(k_4), \quad (\text{B.5})$$

where E_i^μ , $F_i^{\mu\nu}$, $H_i^{\mu\nu}$, ($i = L, R$) are

$$E_i^\mu = \{e_{1i}k_1 + e_{2i}k_2\} k_1^\mu + \{e_{1i}k_2 + e_{2i}k_1\} k_2^\mu \\ + \{e_{3i}(k_1^2 + k_2^2) - e_{4i}k_1 k_2\} \gamma^\mu - e_{5i}i\epsilon^{\alpha\beta\mu\nu} \gamma_5 \gamma_\nu k_{1\alpha} k_{2\beta}, \quad (\text{B.6})$$

$$F_i^{\mu\nu} = f_{1i\alpha} i\epsilon^{\alpha\mu\nu\beta} \gamma_5 \gamma_\beta + f_{2i\alpha} g^{\mu\nu} \gamma^\alpha + f_{3i\alpha} g^{\alpha\mu} \gamma^\nu + f_{4i\alpha} g^{\alpha\nu} \gamma^\mu, \quad (\text{B.7})$$

$$H_i^{\mu\nu} = h_{1i\alpha} i\epsilon^{\alpha\mu\nu\beta} \gamma_5 \gamma_\beta + h_{2i\alpha} g^{\mu\nu} \gamma^\alpha + h_{3i\alpha} g^{\alpha\mu} \gamma^\nu + h_{4i\alpha} g^{\alpha\nu} \gamma^\mu. \quad (\text{B.8})$$

Appendix B.1.1. Coefficients in $\mathcal{O}_{qqqq}$

The coefficients of 4-Fermi operator are given as

$$C_{LL}^{\text{SUSY}} = \frac{2}{9} \left[f_1(m_{\tilde{q}_L}, m_{\tilde{q}'_L}) + f_2(m_{\tilde{q}_L}, m_{\tilde{q}'_L}) \right], \quad (\text{B.9})$$

$$C_{RR}^{\text{SUSY}} = \frac{2}{9} \left[f_1(m_{\tilde{q}_R}, m_{\tilde{q}'_R}) + f_2(m_{\tilde{q}_R}, m_{\tilde{q}'_R}) \right], \quad (\text{B.10})$$

$$C_{LR}^{\text{SUSY}} = -\frac{2}{9} \left[f_1(m_{\tilde{q}_R}, m_{\tilde{q}'_L}) - f_2(m_{\tilde{q}_L}, m_{\tilde{q}'_R}) \right], \quad (\text{B.11})$$

$$C_{RL}^{\text{SUSY}} = -\frac{2}{9} \left[f_1(m_{\tilde{q}_L}, m_{\tilde{q}'_R}) + f_2(m_{\tilde{q}_L}, m_{\tilde{q}'_R}) \right], \quad (\text{B.12})$$

$$D_{LL}^{\text{SUSY}} = -\frac{1}{3} f_1(m_{\tilde{q}_L}, m_{\tilde{q}'_L}) - \frac{7}{6} f_2(m_{\tilde{q}_L}, m_{\tilde{q}'_L}), \quad (\text{B.13})$$

$$D_{RR}^{\text{SUSY}} = -\frac{1}{3} f_1(m_{\tilde{q}_R}, m_{\tilde{q}'_R}) - \frac{7}{6} f_2(m_{\tilde{q}_R}, m_{\tilde{q}'_R}), \quad (\text{B.14})$$

$$D_{LR}^{\text{SUSY}} = -\frac{7}{6} f_1(m_{\tilde{q}_L}, m_{\tilde{q}'_R}) - \frac{1}{3} f_2(m_{\tilde{q}_L}, m_{\tilde{q}'_R}), \quad (\text{B.15})$$

$$D_{RL}^{\text{SUSY}} = -\frac{7}{6} f_2(m_{\tilde{q}_R}, m_{\tilde{q}'_L}) - \frac{1}{3} f_1(m_{\tilde{q}_R}, m_{\tilde{q}'_L}), \quad (\text{B.16})$$

where f_1 and f_2 are given by

$$f_1(m_{\tilde{q}}, m_{\tilde{q}'}) = \beta(m_{\tilde{q}'}, m_{\tilde{q}}, m_{\tilde{g}}, 1, 1, 0), \quad (\text{B.17})$$

$$f_2(m_{\tilde{q}}, m_{\tilde{q}'}) = m_{\tilde{g}}^2 \beta(m_{\tilde{q}'}, m_{\tilde{q}}, m_{\tilde{g}}, 2, 1, 0). \quad (\text{B.18})$$

Appendix B.1.2. Coefficients in $\mathcal{O}_{qqG}$

The coefficients of q - q - G operator are given as

$$E_L^\mu \equiv E^\mu(m_{\tilde{q}} = m_{\tilde{q}_L}) \quad (\text{B.19})$$

$$\begin{aligned} &= \{e_1(m_{\tilde{q}_L})\not{k}_1 + e_2(m_{\tilde{q}_L})\not{k}_2\}k_1^\mu + \{e_1(m_{\tilde{q}_L})\not{k}_2 + e_2(m_{\tilde{q}_L})\not{k}_1\}k_2^\mu \\ &\quad + \{e_3(m_{\tilde{q}_L})(k_1^2 + k_2^2) - e_4(m_{\tilde{q}_L})k_1 k_2\}\gamma^\mu \\ &\quad - e_5(m_{\tilde{q}_L})i\epsilon^{\alpha\beta\mu\nu}\gamma_5\gamma_\nu k_{1\alpha}k_{2\beta}, \end{aligned} \quad (\text{B.20})$$

$$E_R^\mu = E^\mu(m_{\tilde{q}_R}), \quad (\text{B.21})$$

$$\begin{aligned} e_1(m_{\tilde{q}}) &= \frac{107m_{\tilde{g}}^6 - 495m_{\tilde{g}}^4m_{\tilde{q}}^2 + 477m_{\tilde{g}}^2m_{\tilde{q}}^4 - 89m_{\tilde{q}}^6 - 6\left(m_{\tilde{g}}^6 + 3m_{\tilde{g}}^4m_{\tilde{q}}^2 - 54m_{\tilde{g}}^2m_{\tilde{q}}^4 + 18m_{\tilde{q}}^6\right)\log\left(m_{\tilde{g}}^2/m_{\tilde{q}}^2\right)}{18\left(m_{\tilde{g}}^2 - m_{\tilde{q}}^2\right)^4}, \end{aligned} \quad (\text{B.22})$$

$$\begin{aligned} e_2(m_{\tilde{q}}) &= \frac{-203m_{\tilde{g}}^6 + 351m_{\tilde{g}}^4m_{\tilde{q}}^2 - 189m_{\tilde{g}}^2m_{\tilde{q}}^4 + 41m_{\tilde{q}}^6 + 6\left(m_{\tilde{g}}^6 + 51m_{\tilde{g}}^4m_{\tilde{q}}^2 - 54m_{\tilde{g}}^2m_{\tilde{q}}^4 + 18m_{\tilde{q}}^6\right)\log\left(m_{\tilde{g}}^2/m_{\tilde{q}}^2\right)}{18\left(m_{\tilde{g}}^2 - m_{\tilde{q}}^2\right)^4}, \end{aligned} \quad (\text{B.23})$$

$$e_3(m_{\tilde{q}}) = e_2(m_{\tilde{q}}), \quad (\text{B.24})$$

$$\begin{aligned} e_4(m_{\tilde{q}}) &= \frac{-155m_{\tilde{g}}^6 + 423m_{\tilde{g}}^4m_{\tilde{q}}^2 - 333m_{\tilde{g}}^2m_{\tilde{q}}^4 + 65m_{\tilde{q}}^6 + 6\left(m_{\tilde{g}}^6 + 27m_{\tilde{g}}^4m_{\tilde{q}}^2 - 54m_{\tilde{g}}^2m_{\tilde{q}}^4 + 18m_{\tilde{q}}^6\right)\log\left(m_{\tilde{g}}^2/m_{\tilde{q}}^2\right)}{9\left(m_{\tilde{g}}^2 - m_{\tilde{q}}^2\right)^4}, \end{aligned} \quad (\text{B.25})$$

$$e_5(m_{\tilde{q}}) = \frac{9\left(m_{\tilde{g}}^4 - m_{\tilde{q}}^4 - 2m_{\tilde{g}}^2m_{\tilde{q}}^2\log\left(m_{\tilde{g}}^2/m_{\tilde{q}}^2\right)\right)}{\left(m_{\tilde{g}}^2 - m_{\tilde{q}}^2\right)^3}. \quad (\text{B.26})$$

Appendix B.1.3. Coefficients in $\mathcal{O}_{qqGG}$

The coefficients of q - q - G - G operator are given as

$$F^{\mu\nu} = f_{1\alpha}i\epsilon^{\alpha\mu\nu\beta}\gamma_5\gamma_\beta + f_{2\alpha}g^{\mu\nu}\gamma^\alpha + f_{3\alpha}g^{\alpha\mu}\gamma^\nu + f_{4\alpha}g^{\alpha\nu}\gamma^\mu, \quad (\text{B.27})$$

$$H^{\mu\nu} = h_{1\alpha}i\epsilon^{\alpha\mu\nu\beta}\gamma_5\gamma_\beta + h_{2\alpha}g^{\mu\nu}\gamma^\alpha + h_{3\alpha}g^{\alpha\mu}\gamma^\nu + h_{4\alpha}g^{\alpha\nu}\gamma^\mu, \quad (\text{B.28})$$

$$f_{1\alpha} = \left[\frac{1}{2}S_{0\alpha} - 2(P_0 + Q_0)_\alpha\right](m_a = m_{\tilde{g}}, m_b = m_{\tilde{q}_i}), \quad (\text{B.29})$$

$$f_{2\alpha} = \left[-3K_\alpha + \frac{1}{2}R_{3\alpha} + \frac{1}{2}S_{1\alpha} + 2(P_1 + Q_1)_\alpha\right](m_a = m_{\tilde{g}}, m_b = m_{\tilde{q}_i}), \quad (\text{B.30})$$

$$f_{3\alpha} = \left[\frac{1}{2}R_{2\alpha} + \frac{1}{2}S_{2\alpha} + 2(P_3 + Q_3)_\alpha\right](m_a = m_{\tilde{g}}, m_b = m_{\tilde{q}_i}), \quad (\text{B.31})$$

$$f_{4\alpha} = \left[\frac{1}{2}R_{1\alpha} + \frac{1}{2}S_{3\alpha} + 2(P_2 + Q_2)_\alpha \right] (m_a = m_{\tilde{g}}, m_b = m_{\tilde{q}_i}), \quad (\text{B.32})$$

$$h_{1\alpha} = [12(P_0 + Q_0)_\alpha] (m_a = m_{\tilde{g}}, m_b = m_{\tilde{q}_i}), \quad (\text{B.33})$$

$$h_{2\alpha} = \left[2K_\alpha - \frac{1}{3}R_{3\alpha} + 12(P_1 + Q_1)_\alpha \right] (m_a = m_{\tilde{g}}, m_b = m_{\tilde{q}_i}), \quad (\text{B.34})$$

$$h_{3\alpha} = \left[-\frac{1}{3}R_{2\alpha} + 12(P_3 + Q_3)_\alpha \right] (m_a = m_{\tilde{g}}, m_b = m_{\tilde{q}_i}), \quad (\text{B.35})$$

$$h_{4\alpha} = \left[-\frac{1}{3}R_{1\alpha} + 12(P_2 + Q_2)_\alpha \right] (m_a = m_{\tilde{g}}, m_b = m_{\tilde{q}_i}), \quad (\text{B.36})$$

where $K(m_a, m_b)$, $P(m_a, m_b)$, $Q(m_a, m_b)$, $R(m_a, m_b)$, $S(m_a, m_b)$ are written in terms of a linear combination of the momentums;

$$K_\alpha = a_1 k_{1\alpha} + a_2 k_{4\alpha}, \quad (\text{B.37})$$

$$P_{0\alpha} = (i_{11} + i_{12} + i_{13})k_{1\alpha} + (i_{21} + i_{22} + i_{23})k_{2\alpha} + (i_{31} + i_{32} + i_{33})k_{4\alpha}, \quad (\text{B.38})$$

$$P_{1\alpha} = (i_{11} + i_{12} - i_{13})k_{1\alpha} + (i_{21} + i_{22} - i_{23})k_{2\alpha} + (i_{31} + i_{32} - i_{33})k_{4\alpha}, \quad (\text{B.39})$$

$$P_{2\alpha} = (i_{11} - i_{12} + i_{13})k_{1\alpha} + (i_{21} - i_{22} + i_{23})k_{2\alpha} + (i_{31} - i_{32} + i_{33})k_{4\alpha}, \quad (\text{B.40})$$

$$P_{3\alpha} = (-i_{11} + i_{12} + i_{13})k_{1\alpha} + (-i_{21} + i_{22} + i_{23})k_{2\alpha} + (-i_{31} + i_{32} + i_{33})k_{4\alpha}, \quad (\text{B.41})$$

$$Q_{0\alpha} = (j_{11} - j_{12} + j_{13})k_{1\alpha} + (j_{21} - j_{22} + j_{23})k_{2\alpha} + (j_{31} - j_{32} + j_{33})k_{4\alpha}, \quad (\text{B.42})$$

$$Q_{1\alpha} = Q_{0\alpha}, \quad (\text{B.43})$$

$$Q_{2\alpha} = (-j_{11} + j_{12} + j_{13})k_{1\alpha} + (-j_{21} + j_{22} + j_{23})k_{2\alpha} + v(-j_{31} + j_{32} + j_{33})k_{4\alpha}, \quad (\text{B.44})$$

$$Q_{3\alpha} = (j_{11} + j_{12} - j_{13})k_{1\alpha} + (j_{21} + j_{22} - j_{23})k_{2\alpha} + (j_{31} + j_{32} - j_{33})k_{4\alpha}, \quad (\text{B.45})$$

$$R_{1\alpha} = b_{11}k_{1\alpha} + b_{21}k_{2\alpha} + b_{31}k_{4\alpha}, \quad (\text{B.46})$$

$$R_{2\alpha} = b_{12}k_{1\alpha} + b_{22}k_{2\alpha} + b_{32}k_{4\alpha}, \quad (\text{B.47})$$

$$R_{3\alpha} = b_{13}k_{1\alpha} + b_{23}k_{2\alpha} + b_{33}k_{4\alpha}, \quad (\text{B.48})$$

$$S_{0\alpha} = (-f_{12} + f_{13})k_{1\alpha} + (-f_{22} + f_{23})k_{2\alpha} + (-f_{32} + f_{33})k_{4\alpha}, \quad (\text{B.49})$$

$$S_{1\alpha} = (f_{12} + f_{13})k_{1\alpha} + (f_{22} + f_{23})k_{2\alpha} + (f_{32} + f_{33})k_{4\alpha}, \quad (\text{B.50})$$

$$S_{2\alpha} = S_{1\alpha}, \quad (\text{B.51})$$

$$S_{3\alpha} = (h_1 - 2f_{11} - f_{12} - f_{13})k_{1\alpha} + (h_2 - 2f_{21} - f_{22} - f_{23})k_{2\alpha} \\ + (h_3 - 2f_{31} - f_{32} - f_{33})k_{4\alpha}. \quad (\text{B.52})$$

These coefficients, $a_1, a_2, \dots$, are given in terms of Feynman parameter integral as

$$\begin{aligned} a_1(m_a, m_b) &= a_1(m_a, m_b) = a_2(m_a, m_b) \\ &= -\alpha(m_a, m_b, 1, 2) + 2\alpha(m_a, m_b, 1, 1) - \alpha(m_a, m_b, 1, 0), \end{aligned} \quad (\text{B.53})$$

$$\begin{aligned} i_{11}(m_a, m_b) &= -2i_{21}(m_a, m_b) = -2i_{22}(m_a, m_b) = 2i_{32}(m_a, m_b) \\ &= 2i_{33}(m_a, m_b) = -2\alpha(m_a, m_b, 1, 3), \end{aligned} \quad (\text{B.54})$$

$$\begin{aligned} i_{12}(m_a, m_b) &= i_{13}(m_a, m_b) \\ &= -2\alpha(m_a, m_b, 1, 3) + 3\alpha(m_a, m_b, 1, 2), \end{aligned} \quad (\text{B.55})$$

$$\begin{aligned} i_{23}(m_a, m_b) &= -i_{31}(m_a, m_b) \\ &= \alpha(m_a, m_b, 1, 3) - 3\alpha(m_a, m_b, 1, 2), \end{aligned} \quad (\text{B.56})$$

$$\begin{aligned} j_{11}(m_a, m_b) &= j_{12}(m_a, m_b) \\ &= -2m_a^2\alpha(m_a, m_b, 2, 3) + 3m_a^2\alpha(m_a, m_b, 2, 2) \end{aligned} \quad (\text{B.57})$$

$$\begin{aligned} j_{13}(m_a, m_b) &= 2j_{21}(m_a, m_b) = 2j_{23}(m_a, m_b) = -2j_{31}(m_a, m_b) \\ &= -2j_{32}(m_a, m_b) = 2m_a^2\alpha(m_a, m_b, 2, 3), \end{aligned} \quad (\text{B.58})$$

$$\begin{aligned} j_{22}(m_a, m_b) &= -j_{33}(m_a, m_b) \\ &= m_a^2\alpha(m_a, m_b, 2, 3) - 3m_a^2\alpha(m_a, m_b, 2, 2), \end{aligned} \quad (\text{B.59})$$

$$\begin{aligned} b_{11}(m_a, m_b) &= 2b_{12}(m_a, m_b) = b_{13}(m_a, m_b) = -2b_{21}(m_a, m_b) \\ &= -2b_{22}(m_a, m_b) = -2b_{23}(m_a, m_b) \\ &= -2b_{31}(m_a, m_b) = 2b_{32}(m_a, m_b) = 2b_{33}(m_a, m_b) \\ &= 8\alpha(m_a, m_b, 1, 3) - 9\alpha(m_a, m_b, 1, 2) + 6\alpha(m_a, m_b, 1, 1) \\ &\quad - \alpha(m_a, m_b, 1, 0), \end{aligned} \quad (\text{B.60})$$

$$f_{11}(m_a, m_b) = -6\alpha(m_a, m_b, 1, 3) + 6\alpha(m_a, m_b, 1, 2), \quad (\text{B.61})$$

$$\begin{aligned} f_{12}(m_a, m_b) &= f_{13}(m_a, m_b) = f_{22}(m_a, m_b) \\ &= 12\alpha(m_a, m_b, 1, 1) - 12\alpha(m_a, m_b, 1, 0), \end{aligned} \quad (\text{B.62})$$

$$f_{23}(m_a, m_b) = 12\alpha(m_a, m_b, 1, 3) + 6\alpha(m_a, m_b, 1, 2) - 6\alpha(m_a, m_b, 1, 1), \quad (\text{B.63})$$

$$f_{31}(m_a, m_b) = -12\alpha(m_a, m_b, 1, 3) + 6\alpha(m_a, m_b, 1, 2) + 6\alpha(m_a, m_b, 1, 1), \quad (\text{B.64})$$

$$\begin{aligned} f_{32}(m_a, m_b) &= f_{33}(m_a, m_b) \\ &= 3\alpha(m_a, m_b, 1, 4) + 4\alpha(m_a, m_b, 1, 3) - \alpha(m_a, m_b, 1, 0), \end{aligned} \quad (\text{B.65})$$

$$\begin{aligned} h_1(m_a, m_b) &= m_a^2\alpha(m_a, m_b, 2, 4) - 2m_a^2\alpha(m_a, m_b, 2, 3) \\ &\quad + 4m_a^2\alpha(m_a, m_b, 2, 1) - m_a^2\alpha(m_a, m_b, 2, 0), \end{aligned} \quad (\text{B.66})$$

$$\begin{aligned} h_3(m_a, m_b) &= -12m_a^2\alpha(m_a, m_b, 2, 3) \\ &\quad + 6m_a^2\alpha(m_a, m_b, 2, 2) + 6m_a^2\alpha(m_a, m_b, 2, 1), \end{aligned} \quad (\text{B.67})$$

and other coefficients are zero.

Appendix B.2. UED

Let us list the dimension six operators of QCD in UED.

Appendix B.2.1. Coefficients in $\mathcal{O}_{qqqq}$

When $(\bar{q}q)(\bar{q}'q')$ chirality is (LL)(LL) or (RR)(RR), 4-Fermi operator is

$$\mathcal{O}_{qqqq}(x) = \frac{g_s^4}{192\pi^2} \left[f_1 (\bar{q}\gamma^\mu q)(\bar{q}'\gamma_\mu q') + f_2 \left(\bar{q}\gamma^\mu \frac{\lambda^a}{2} q \right) \left(\bar{q}'\gamma^\mu \frac{\lambda^a}{2} q' \right) \right], \quad (\text{B.68})$$

and the coefficients f_1, f_2 are

$$f_{1\text{L}} = \frac{4 \left(m_g^4 + 4m_g^2 m_{\text{L}}^2 \log \left(\frac{m_{\text{L}}}{m_g} \right) - m_{\text{L}}^4 \right)}{(m_g^2 - m_{\text{L}}^2)^3}, \quad (\text{B.69})$$

$$f_{1\text{R}} = \frac{4 \left(m_g^4 + 4m_g^2 m_{\text{R}}^2 \log \left(\frac{m_{\text{R}}}{m_g} \right) - m_{\text{R}}^4 \right)}{(m_g^2 - m_{\text{R}}^2)^3}, \quad (\text{B.70})$$

$$\begin{aligned} f_{2\text{L}} = & \frac{1}{(m_g^2 - m_5^2)(m_5 - m_{\text{L}})^2(m_5 + m_{\text{L}})^2(m_g - m_{\text{L}})^2(m_g + m_{\text{L}})^2} \\ & \times \left[18m_{\text{L}}^2 \left(m_g^2 \left(m_5^4 \log \left(\frac{m_{\text{L}}^2}{m_g^2} \right) - m_{\text{L}}^4 \log \left(\frac{m_g^2}{m_{\text{L}}^2} \right) \right) + m_5^2 (m_g^4 + m_{\text{L}}^4) \log \left(\frac{m_5^2}{m_{\text{L}}^2} \right) \right. \right. \\ & + 2m_5^2 m_g^2 m_{\text{L}}^2 \log \left(\frac{m_g^2}{m_5^2} \right) \\ & \left. \left. + (m_5^2 - m_g^2)(m_5^2 - m_{\text{L}}^2)(m_g^2 - m_{\text{L}}^2) \right) + \frac{7 \left(-m_5^4 + 2m_5^2 m_{\text{L}}^2 \log \left(\frac{m_5^2}{m_{\text{L}}^2} \right) + m_{\text{L}}^4 \right)}{2(m_5^2 - m_{\text{L}}^2)^3} \right. \\ & \left. \left. + \frac{30 \left(-m_g^4 + 2m_g^2 m_{\text{L}}^2 \log \left(\frac{m_g^2}{m_{\text{L}}^2} \right) + m_{\text{L}}^4 \right)}{(m_g^2 - m_{\text{L}}^2)^3} \right] \right], \quad (\text{B.71}) \end{aligned}$$

$$\begin{aligned}
f_{2R} = & \frac{1}{(m_g^2 - m_5^2)(m_5 - m_R)^2(m_5 + m_R)^2(m_g - m_R)^2(m_g + m_R)^2} \\
& \times \left[18m_R^2 \left(m_g^2 \left(m_5^4 \log \left(\frac{m_R^2}{m_g^2} \right) - m_R^4 \log \left(\frac{m_g^2}{m_R^2} \right) \right) + m_5^2 (m_g^4 + m_R^4) \log \left(\frac{m_5^2}{m_R^2} \right) \right. \right. \\
& + 2m_5^2 m_g^2 m_R^2 \log \left(\frac{m_g^2}{m_5^2} \right) \\
& \left. \left. + (m_5^2 - m_g^2)(m_5^2 - m_R^2)(m_g^2 - m_R^2) \right) + \frac{7 \left(-m_5^4 + 2m_5^2 m_R^2 \log \left(\frac{m_5^2}{m_R^2} \right) + m_R^4 \right)}{2(m_5^2 - m_R^2)^3} \right. \\
& \left. + \frac{30 \left(-m_g^4 + 2m_g^2 m_R^2 \log \left(\frac{m_g^2}{m_R^2} \right) + m_R^4 \right)}{(m_g^2 - m_R^2)^3} \right]. \quad (B.72)
\end{aligned}$$

When $(\bar{u}u)(\bar{t}t)$ chirality is (LL)(RR) or (RR)(LL), 4-Fermi operator is

$$\mathcal{O}_{qqqq}(x) = \frac{g_s^4}{192\pi^2} \left[f_3 (\bar{q}\gamma^\mu q)(\bar{q}'\gamma_\mu q') + f_4 \left(\bar{q}\gamma^\mu \frac{\lambda^a}{2} q \right) \left(\bar{q}'\gamma^\mu \frac{\lambda^a}{2} q' \right) \right], \quad (B.73)$$

and the coefficients f_3, f_4 are

$$\begin{aligned}
f_3(m_q, m_{q'}) = & -\frac{1}{(m_g^2 - m_q^2)^2(m_g^2 - m_{q'}^2)^2(m_q^2 - m_{q'}^2)} \\
& \times \left[8 \left(2m_g^2 m_{q'}^4 (m_g^2 - 2m_q^2) \log(m_g) - 2m_{q'}^4 (m_g^2 - m_q^2)^2 \log(m_{q'}) \right. \right. \\
& + m_g^2 (m_g^2 - m_q^2)(m_g^2 - m_{q'}^2)(m_q^2 - m_{q'}^2) \\
& \left. \left. + 2m_q^4 \left(m_g^4 \log \left(\frac{m_q}{m_g} \right) + 2m_g^2 m_{q'}^2 \log \left(\frac{m_g}{m_q} \right) + m_{q'}^4 \log(m_q) \right) \right) \right], \quad (B.74)
\end{aligned}$$

$$\begin{aligned}
f_4(m_q, m_{q'}) = & \frac{9}{2(m_5^2 - m_q^2)(m_5^2 - m_{q'}^2)(m_q^2 - m_{q'}^2)} \\
& \times \left(\frac{(m_5^2 - m_q^2)(-m_g^4 + 4m_g^4 \log(m_g) + m_{q'}^4 - 4m_{q'}^4 \log(m_{q'}))}{m_g^2 - m_{q'}^2} \right. \\
& - \frac{(m_5^2 - m_{q'}^2)(-m_g^4 + 4m_g^4 \log(m_g) + m_q^4 - 4m_q^4 \log(m_q))}{m_g^2 - m_q^2} \\
& \left. + \frac{(m_q^2 - m_{q'}^2)(-m_5^4 + 4m_5^4 \log(m_5) + m_g^4 - 4m_g^4 \log(m_g))}{m_5^2 - m_g^2} \right) \\
& + \frac{9}{2(m_5^2 - m_q^2)^2(m_5^2 - m_{q'}^2)(m_{q'}^2 - m_q^2)} \\
& \times \left[2m_5^2 m_{q'}^4 (m_5^2 - 2m_q^2) \log(m_5) - 2m_{q'}^4 (m_5^2 - m_q^2)^2 \log(m_{q'}) \right. \\
& + m_5^2 (m_5^2 - m_q^2)(m_5^2 - m_{q'}^2)(m_q^2 - m_{q'}^2) \\
& \left. + 2m_q^4 \left(m_5^4 \log\left(\frac{m_q}{m_5}\right) + 2m_5^2 m_{q'}^2 \log\left(\frac{m_5}{m_q}\right) + m_{q'}^4 \log(m_q) \right) \right] \\
& - \frac{30}{(m_g^2 - m_q^2)^2(m_g^2 - m_{q'}^2)(m_q^2 - m_{q'}^2)} \\
& \times \left[2m_g^2 m_{q'}^4 (m_g^2 - 2m_q^2) \log(m_g) - 2m_{q'}^4 (m_g^2 - m_q^2)^2 \log(m_{q'}) \right. \\
& + m_g^2 (m_g^2 - m_q^2)(m_g^2 - m_{q'}^2)(m_q^2 - m_{q'}^2) \\
& \left. + 2m_q^4 \left(m_g^4 \log\left(\frac{m_q}{m_g}\right) + 2m_g^2 m_{q'}^2 \log\left(\frac{m_g}{m_q}\right) + m_{q'}^4 \log(m_q) \right) \right], \tag{B.75}
\end{aligned}$$

where $m_q, m_{q'}$ are m_L or m_R ($m_q \neq m_{q'}$).

Appendix B.2.2. Coefficients in $\mathcal{O}_{qqG}$

q - q - G operator in UED takes the form as

$$\begin{aligned}
\int d^4x \mathcal{O}_{qqG}(x) = & -\frac{g_s^3}{192\pi^2} \int \frac{d^4k_1}{(2\pi)^4} \frac{d^4k_2}{(2\pi)^4} \frac{d^4k_3}{(2\pi)^4} (2\pi)^4 \delta^4(-k_1 + k_2 + k_3) \\
& \times \bar{q}(k_1) C^\mu G_\mu(k_3) q(k_2), \tag{B.76}
\end{aligned}$$

$$C^\mu \equiv c_1 i \epsilon^{\alpha\beta\mu\nu} \gamma^5 \gamma_\nu + c_2 g^{\mu\alpha} \gamma^\beta + c_3 g^{\mu\beta} \gamma^\alpha + c_4 g^{\alpha\beta} \gamma^\mu, \quad (\text{B.77})$$

$$c_1 \equiv c_{11} k_{1\alpha} k_{2\beta}, \quad (\text{B.78})$$

$$c_2 \equiv c_{21} k_{1\alpha} k_{1\beta} + c_{22} k_{1\alpha} k_{2\beta} + c_{23} k_{2\alpha} k_{1\beta} + c_{24} k_{2\alpha} k_{2\beta}, \quad (\text{B.79})$$

$$c_3 \equiv c_{31} k_{1\alpha} k_{1\beta} + c_{32} k_{1\alpha} k_{2\beta} + c_{33} k_{2\alpha} k_{1\beta} + c_{34} k_{2\alpha} k_{2\beta}, \quad (\text{B.80})$$

$$c_4 \equiv c_{41} k_{1\alpha} k_{1\beta} + c_{42} k_{1\alpha} k_{2\beta} + c_{43} k_{2\alpha} k_{2\beta}. \quad (\text{B.81})$$

The coefficients are given as

$$c_{11} = 6 \frac{m_g^4 - 4m_g^2 m_q^2 \log\left(\frac{m_g}{m_q}\right) - m_q^4}{(m_g^2 - m_q^2)^3}, \quad (\text{B.82})$$

$$c_{21} = 6 \frac{5m_g^6 - 27m_g^4 m_q^2 + 27m_g^2 m_q^4 + 12m_g^4 (m_g^2 - 3m_q^2) \log\left(\frac{m_g}{m_q}\right) - 5m_q^6}{18(m_g^2 - m_q^2)^4}, \quad (\text{B.83})$$

$$c_{22} = 6 \frac{7m_g^6 - 27m_g^2 m_q^4 + 12(m_g^6 - 6m_g^4 m_q^2 + 6m_g^2 m_q^4) \log\left(\frac{m_g}{m_q}\right) + 20m_q^6}{18(m_g^2 - m_q^2)^4}, \quad (\text{B.84})$$

$$c_{23} = 6 \frac{12m_g^6 \log\left(\frac{m_g}{m_q}\right) - 11m_g^6 + 18m_g^4 m_q^2 - 9m_g^2 m_q^4 + 2m_q^6}{18(m_g^2 - m_q^2)^4}, \quad (\text{B.85})$$

$$c_{24} = 6 \frac{12m_g^6 \log\left(\frac{m_g}{m_q}\right) + 11m_g^6 - 18m_g^4 m_q^2 + 9m_g^2 m_q^4 - 2m_q^6}{18(m_g^2 - m_q^2)^4}, \quad (\text{B.86})$$

$$c_{31} = \frac{1}{3} \left(\frac{5m_g^6 - 27m_g^4 m_q^2 + 27m_g^2 m_q^4 + 12(m_g^6 - 3m_g^4 m_q^2) \log\left(\frac{m_g}{m_q}\right) - 5m_q^6}{(m_g^2 - m_q^2)^4} \right. \\ \left. - \frac{6(2m_g^6 - 9m_g^4 m_q^2 + 18m_g^2 m_q^4 + 12m_g^6 \log\left(\frac{m_g}{m_5}\right) - 11m_q^6)}{(m_5^2 - m_q^2)^4} \right), \quad (\text{B.87})$$

$$c_{32} = \frac{1}{3} \left(\frac{3(7m_g^6 - 36m_g^4 m_q^2 + 45m_g^2 m_q^4 + 12(3m_g^2 m_q^4 - 2m_q^6) \log\left(\frac{m_5}{m_q}\right) - 16m_q^6)}{(m_5^2 - m_q^2)^4} \right. \\ \left. + \frac{7m_g^6 - 27m_g^2 m_q^4 + 12(m_g^6 - 6m_g^4 m_q^2 + 6m_g^2 m_q^4) \log\left(\frac{m_g}{m_q}\right) + 20m_q^6}{(m_g^2 - m_q^2)^4} \right), \quad (\text{B.88})$$

$$c_{33} = \frac{1}{3} \left(\frac{12m_g^6 \log\left(\frac{m_g}{m_q}\right) - 11m_g^6 + 18m_g^4 m_q^2 - 9m_g^2 m_q^4 + 2m_q^6}{(m_g^2 - m_q^2)^4} \right. \\ \left. - \frac{3(2m_g^6 - 9m_g^4 m_q^2 + 18m_g^2 m_q^4 + 12m_q^6 \log\left(\frac{m_q}{m_5}\right) - 11m_q^6)}{(m_5^2 - m_q^2)^4} \right), \quad (\text{B.89})$$

$$c_{34} = \frac{1}{3} \left(\frac{12m_g^6 \log\left(\frac{m_q}{m_g}\right) + 11m_g^6 - 18m_g^4 m_q^2 + 9m_g^2 m_q^4 - 2m_q^6}{(m_g^2 - m_q^2)^4} \right. \\ \left. - \frac{3\left(-5m_5^6 + v27m_5^4 m_q^2 - 27m_5^2 m_q^4 + 12m_q^4 (m_q^2 - 3m_5^2) \log\left(\frac{m_5}{m_q}\right) + 5m_q^6\right)}{(m_5^2 - m_q^2)^4} \right), \quad (\text{B.90})$$

$$c_{41} = \frac{18m_g m_q}{(m_g^2 - m_5^2)^3 (m_5 - m_q)(m_5 + m_q)(m_g - m_q)^3 (m_g + m_q)^3} \\ \times \left[4m_g^2 m_q^2 (m_5^2 - m_g^2)^3 \log(m_q) + 4m_5^2 m_g^2 \log(m_5) (m_g^2 - m_q^2)^3 \right. \\ \left. + (m_5^2 - m_q^2) ((m_5^2 - m_g^2) (m_g^2 - m_q^2) (m_5^2 (m_g^2 + m_q^2) - 3m_g^4 + m_g^2 m_q^2) \right. \\ \left. - 4m_g^2 \log(m_g) (-3m_5^2 m_g^2 m_q^2 + m_5^2 m_q^2 (m_5^2 + m_q^2) + m_g^6)) \right] \\ + \frac{5m_5^6 - 27m_5^4 m_q^2 + 27m_5^2 m_q^4 - 12m_q^4 (m_q^2 - 3m_5^2) \log\left(\frac{m_5}{m_q}\right) - 5m_q^6}{2(m_5^2 - m_q^2)^4} \\ + \frac{5\left(-5m_g^6 + 27m_g^4 m_q^2 - 27m_g^2 m_q^4 + 12(m_g^6 - 3m_g^4 m_q^2) \log\left(\frac{m_g}{m_q}\right) + 5m_q^6\right)}{3(m_g^2 - m_q^2)^4} \\ - \frac{5m_g^6 - 27m_g^4 m_q^2 + 27m_g^2 m_q^4 + 12(m_g^6 - 3m_g^4 m_q^2) \log\left(\frac{m_q}{m_g}\right) - 5m_q^6}{3(m_g^2 - m_q^2)^4}, \quad (\text{B.91})$$

$$c_{42} = \frac{1}{3} \left(- \frac{108m_g m_q}{(m_g^2 - m_5^2)^3 (m_5 - m_q)^2 (m_5 + m_q)^2 (m_g - m_q)^2 (m_g + m_q)^2} \right. \\ \times \left[-2m_q^4 (m_5^2 - m_g^2)^3 \log(m_q) + (m_q^2 - m_5^2) \right. \\ \times \left((m_5^2 - m_g^2) (m_g^2 - m_q^2) (2m_5^2 m_g^2 - m_q^2 (m_5^2 + m_g^2)) \right. \\ \left. + 2m_g^2 (m_5 - m_q) (m_5 + m_q) \log(m_g) (m_5^2 (m_g^2 - 2m_q^2) + m_g^4) \right) \\ \left. + 2m_5^2 \log(m_5) (m_g^2 - m_q^2)^2 (m_5^4 + m_5^2 m_g^2 - 2m_g^2 m_q^2) \right] \\ + \frac{3\left(-2m_5^6 + 9m_5^4 m_q^2 - 18m_5^2 m_q^4 + 12m_q^6 \log\left(\frac{m_5}{m_q}\right) + 11m_q^6\right)}{(m_5^2 - m_q^2)^4} \\ + \frac{-12m_g^6 \log\left(\frac{m_g}{m_q}\right) + 11m_g^6 - 18m_g^4 m_q^2 + 9m_g^2 m_q^4 - 2m_q^6}{(m_g^2 - m_q^2)^4}$$

$$\begin{aligned}
& - \frac{7m_g^6 - 27m_g^2m_q^4 + 12(m_g^6 - 6m_g^4m_q^2 + 6m_g^2m_q^4) \log\left(\frac{m_g}{m_q}\right) + 20m_q^6}{(m_g^2 - m_q^2)^4} \\
& + \frac{10\left(12m_g^6 \log\left(\frac{m_q}{m_g}\right) + 11m_g^6 - 18m_g^4m_q^2 + 9m_g^2m_q^4 - 2m_q^6\right)}{(m_g^2 - m_q^2)^4} \Bigg), \quad (\text{B.92})
\end{aligned}$$

$$\begin{aligned}
c_{43} = & - \frac{18m_gm_q}{(m_g^2 - m_q^2)^3 (m_5 - m_q)^3 (m_5 + m_q)^3 (m_g - m_q)(m_g + m_q)} \\
& \times \left(4m_5^2 \left(m_q^2 (m_g^2 - m_5^2)^3 \log(m_q) + m_g^2 (m_5^2 - m_q^2)^3 \log(m_g) \right. \right. \\
& + \log(m_5)(m_q^2 - m_g^2)(m_5^6 - 3m_5^2m_g^2m_q^2 + m_g^2m_q^2(m_g^2 + m_q^2))) \\
& + (m_5^2 - m_g^2)(m_5^2 - m_q^2)(m_g^2 - m_q^2)(3m_5^4 - m_5^2(m_g^2 + m_q^2) - m_g^2m_q^2)) \\
& \left. + \frac{5m_5^6 - 27m_5^4m_q^2 + 27m_5^2m_q^4 - 12m_q^4(m_q^2 - 3m_5^2) \log\left(\frac{m_5}{m_q}\right) - 5m_q^6}{2(m_5^2 - m_q^2)^4} \right. \\
& + \frac{5\left(-5m_g^6 + 27m_g^4m_q^2 - 27m_g^2m_q^4 + 12(m_g^6 - 3m_g^4m_q^2) \log\left(\frac{m_q}{m_g}\right) + 5m_q^6\right)}{3(m_g^2 - m_q^2)^4} \\
& \left. + \frac{-12m_g^6 \log\left(\frac{m_q}{m_g}\right) - 11m_g^6 + 18m_g^4m_q^2 - 9m_g^2m_q^4 + 2m_q^6}{3(m_g^2 - m_q^2)^4} \right), \quad (\text{B.93})
\end{aligned}$$

Appendix B.2.3. Coefficients in $\mathcal{O}_{qqGG}$

The q - q - G - G operator is written as

$$\begin{aligned}
\int d^4x \mathcal{O}_{qqGG}(x) = & \frac{g_s^4}{192\pi^2} \int \frac{d^4k_1}{(2\pi)^4} \frac{d^4k_2}{(2\pi)^4} \frac{d^4k_3}{(2\pi)^4} \frac{d^4k_4}{(2\pi)^4} (2\pi)^4 \delta^4(-k_1 + k_2 + k_3 + k_4) \\
& \times \bar{q}(k_1) \left[F_{L,R}^{\mu\nu} \delta^{ab} + H_{L,R}^{\mu\nu} T^a T^b \right] G_\mu^a(k_2) G_\nu^b(k_3) P_{L,R} q(k_4), \quad (\text{B.94})
\end{aligned}$$

where E_i^μ , $F_i^{\mu\nu}$, $H_i^{\mu\nu}$, ($i = L, R$) are

$$F_i^{\mu\nu} = f_{1i\alpha} i\epsilon^{\alpha\mu\nu\beta} \gamma_5 \gamma_\beta + f_{2i\alpha} g^{\mu\nu} \gamma^\alpha + f_{3i\alpha} g^{\alpha\mu} \gamma^\nu + f_{4i\alpha} g^{\alpha\nu} \gamma^\mu, \quad (\text{B.95})$$

$$H_i^{\mu\nu} = h_{1i\alpha} i\epsilon^{\alpha\mu\nu\beta} \gamma_5 \gamma_\beta + h_{2i\alpha} g^{\mu\nu} \gamma^\alpha + h_{3i\alpha} g^{\alpha\mu} \gamma^\nu + h_{4i\alpha} g^{\alpha\nu} \gamma^\mu. \quad (\text{B.96})$$

$$\begin{aligned}
& +\frac{1}{4}[i_5(m_5, m_q) - 2j_2(m_g, m_q, m_5) - 2j_2(m_5, m_q, m_g) \\
& +n_3(m_g, m_q) + 2n_{13}(m_g, m_q) + l_2(m_q, m_5) - 2s_3(m_g, m_q)] \\
& -\frac{1}{2}[-r_2(m_g, m_q, m_5) - r_{11}(m_g, m_q, m_5) \\
& -s_8(m_g, m_q, m_5) + s_{12}(m_g, m_q, m_5)]\}k_{3\alpha} \\
& +\{-3g_3(m_g, m_q) + 3g_6(m_g, m_q) + 2g_9(m_g, m_q) \\
& +h_3(m_g, m_q) - h_6(m_g, m_q) - h_9(m_g, m_q) \\
& -\frac{1}{2}[g_3(m_5, m_q) - g_6(m_5, m_q) + g_9(m_5, m_q) \\
& -h_3(m_5, m_q) + h_6(m_5, m_q) - h_9(m_5, m_q)] \\
& +\frac{1}{4}[i_6(m_5, m_q) - 2j_3(m_g, m_q, m_5) - 2j_3(m_5, m_q, m_g) \\
& -3[e_2(m_g, m_q, m_5) - e_2(m_5, m_q, m_g)] \\
& +\frac{1}{4}[n_4(m_g, m_q) + 2n_{14}(m_g, m_q) + l_3(m_q, m_5) - 2s_4(m_g, m_q)] \\
& -\frac{1}{2}[-r_3(m_g, m_q, m_5) - r_{12}(m_g, m_q, m_5) - s_8(m_g, m_q, m_5) \\
& +s_{12}(m_g, m_q, m_5)]\}k_{4\alpha}, \tag{B.98}
\end{aligned}$$

$$\begin{aligned}
f_{3\alpha} = & \{\frac{1}{4}[g_1(m_g, m_q) + 4g_4(m_g, m_q) - 8g_7(m_g, m_q) \\
& +4(h_1(m_g, m_q) + h_4(m_g, m_q) + h_7(m_g, m_q))] \\
& -\frac{1}{2}[g_1(m_5, m_q) + g_4(m_5, m_q) - g_7(m_5, m_q) \\
& -h_1(m_5, m_q) - h_4(m_5, m_q) + h_7(m_5, m_q)] \\
& +\frac{1}{4}(i_7(m_5, m_q) - 2j_1(m_g, m_q, m_5)) - 3e_1(m_g, m_q, m_5) \\
& +\frac{1}{4}(n_5(m_g, m_q) + 2n_{12}(m_g, m_q)) + \frac{1}{4}[l_4(m_q, m_5) - 2s_2(m_g, m_q)] \\
& -\frac{1}{2}[-t_1(m_g, m_q, m_5) - r_4(m_g, m_q, m_5) - r_{13}(m_g, m_q, m_5) \\
& +s_9(m_g, m_q, m_5) - s_{13}(m_g, m_q, m_5)]\}k_{1\alpha} \\
& +\{\frac{1}{4}[g_2(m_g, m_q) + 4g_5(m_g, m_q) - 8g_8(m_g, m_q) \\
& +4(h_2(m_g, m_q) + h_5(m_g, m_q) + h_8(m_g, m_q))] \\
& -\frac{1}{2}[g_2(m_5, m_q) + g_5(m_5, m_q) - g_8(m_5, m_q) \\
& -h_2(m_5, m_q) - h_5(m_5, m_q) + h_8(m_5, m_q)] \\
& +\frac{1}{4}[i_8(m_5, m_q) + 2j_2(m_g, m_q, m_5) + n_6(m_g, m_q) + 2n_{13}(m_g, m_q)] \\
& +\frac{1}{4}l_5(m_q, m_5) - \frac{1}{2}[-s_3(m_g, m_q) - t_2(m_g, m_q, m_5) - r_5(m_g, m_q, m_5) \\
& -r_{14}(m_g, m_q, m_5) + s_{10}(m_g, m_q, m_5) - s_{14}(m_g, m_q, m_5)]\}k_{3\alpha} \\
& +\{\frac{1}{4}[g_3(m_g, m_q) + 4g_6(m_g, m_q) - 8g_9(m_g, m_q) \\
& +4(h_3(m_g, m_q) + h_6(m_g, m_q) + h_9(m_g, m_q))] \\
& -\frac{1}{2}(g_3(m_5, m_q) + g_6(m_5, m_q) - g_9(m_5, m_q)
\end{aligned}$$

$$\begin{aligned}
& -h_3(m_5, m_q) - h_6(m_5, m_q) + h_9(m_5, m_q)) \\
& + \frac{1}{4}[i_9(m_5, m_q) + 2j_3(m_g, m_q, m_5)] - 3e_2(m_5, m_q, m_g) \\
& + \frac{1}{4}[n_7(m_g, m_q) + 2n_{14}(m_g, m_q)] \\
& + \frac{1}{4}l_6(m_q, m_5) - \frac{1}{2}[-s_4(m_g, m_q, m_5) - t_3(m_g, m_q, m_5) - r_6(m_g, m_q, m_5) \\
& - r_{15}(m_g, m_q, m_5) + s_{11}(m_g, m_q, m_5) - s_{15}(m_g, m_q, m_5)]\}k_{4\alpha}, \quad (\text{B.99})
\end{aligned}$$

$$\begin{aligned}
f_{4\alpha} = & \left\{ \frac{1}{4}[12g_1(m_g, m_q) - 12g_4(m_g, m_q) + g_7(m_g, m_q) \right. \\
& + 4(-h_1(m_g, m_q) + h_4(m_g, m_q) + h_7(m_g, m_q))] \\
& - \frac{1}{2}(-g_1(m_5, m_q) + g_4(m_5, m_q) + g_7(m_5, m_q) \\
& + h_1(m_5, m_q) - h_4(m_5, m_q) - h_7(m_5, m_q)) \\
& + \frac{1}{4}[i_{10}(m_5, m_q) - 2j_1(m_g, m_q, m_5)] - 3e_1(m_5, m_q, m_g) \\
& + \frac{1}{4}[n_8(m_g, m_q) + l_7(m_q, m_5)] \\
& + \frac{1}{2}[s_5(m_g, m_q) + s_2(m_g, m_q)] - \frac{1}{2}[-t_4(m_g, m_q, m_5) \\
& - r_7(m_g, m_q, m_5) - r_{16}(m_g, m_q, m_5) \\
& - s_8(m_g, m_q, m_5) + s_{12}(m_g, m_q, m_5)]\}k_{1\alpha} \\
& + \left\{ \frac{1}{4}[12g_2(m_g, m_q) - 12g_5(m_g, m_q) + g_8(m_g, m_q) \right. \\
& + 4(-h_2(m_g, m_q) + h_5(m_g, m_q) + h_8(m_g, m_q))] \\
& - \frac{1}{2}(-g_2(m_5, m_q) + g_5(m_5, m_q) \\
& + g_8(m_5, m_q) + h_2(m_5, m_q) - h_5(m_5, m_q) - h_8(m_5, m_q)) \\
& + \frac{1}{4}[i_{11}(m_5, m_q) + 2j_2(m_g, m_q, m_5) + n_9(m_g, m_q) + l_8(m_q, m_5)] \\
& + \frac{1}{2}[s_6(m_g, m_q) + s_3(m_g, m_q)] - \frac{1}{2}[-t_5(m_g, m_q, m_5) - r_8(m_g, m_q, m_5) \\
& - r_{17}(m_g, m_q, m_5) + s_8(m_g, m_q, m_5) - s_{12}(m_g, m_q, m_5)]\}k_{3\alpha} \\
& + \left\{ \frac{1}{4}[12g_3(m_g, m_q) - 12g_6(m_g, m_q) + g_9(m_g, m_q) \right. \\
& + 4(-h_3(m_g, m_q) + h_6(m_g, m_q) + h_9(m_g, m_q))] \\
& - \frac{1}{2}(-g_3(m_5, m_q) + g_6(m_5, m_q) + g_9(m_5, m_q) \\
& + h_3(m_5, m_q) - h_6(m_5, m_q) - h_9(m_5, m_q)) \\
& + \frac{1}{4}[i_{12}(m_5, m_q) + 2j_3(m_g, m_q, m_5)] - 3e_2(m_5, m_q, m_g) \\
& + \frac{1}{4}[n_{10}(m_g, m_q) + l_9(m_q, m_5)] \\
& + \frac{1}{2}[s_7(m_g, m_q) - s_4(m_g, m_q)] - \frac{1}{2}[-t_6(m_g, m_q) - r_9(m_g, m_q, m_5) \\
& - r_{18}(m_g, m_q, m_5) + s_8(m_g, m_q, m_5) - s_{12}(m_g, m_q, m_5)]\}k_{4\alpha}. \quad (\text{B.100})
\end{aligned}$$

The coefficients of color octet part, $H_i^{\mu\nu}$, are given as

$$\begin{aligned}
 h_{1\alpha} = & \left\{ -2(g_1(m_g, m_q) + g_4(m_g, m_q) + g_7(m_g, m_q)) \right. \\
 & -\frac{2}{3}(h_1(m_g, m_q) - h_4(m_g, m_q) - h_7(m_g, m_q)) \\
 & +\frac{1}{3}(g_1(m_5, m_q) + g_4(m_5, m_q) + g_7(m_5, m_q) \\
 & -h_1(m_5, m_q) + h_4(m_5, m_q) - h_7(m_5, m_q)) \\
 & +\frac{3}{2}[i_1(m_5, m_q) - 2j_1(m_g, m_q, m_5)] - 18e_1(m_5, m_q, m_g) \Big\} k_{1\alpha} \\
 & + \left\{ -2(g_2(m_g, m_q) + g_5(m_g, m_q) + g_8(m_g, m_q)) \right. \\
 & -\frac{2}{3}(h_2(m_g, m_q) - h_5(m_g, m_q) - h_8(m_g, m_q)) \\
 & +\frac{1}{3}(g_2(m_5, m_q) + g_5(m_5, m_q) + g_8(m_5, m_q) \\
 & -h_2(m_5, m_q) + h_5(m_5, m_q) - h_8(m_5, m_q)) \\
 & +\frac{3}{2}[i_2(m_5, m_q) + 2j_2(m_g, m_q, m_5)] \Big\} k_{3\alpha} \\
 & + \left\{ -2(g_3(m_g, m_q) + g_6(m_g, m_q) + g_9(m_g, m_q)) \right. \\
 & -\frac{2}{3}(h_3(m_g, m_q) - h_6(m_g, m_q) - h_9(m_g, m_q)) \\
 & +\frac{1}{3}(g_3(m_5, m_q) + g_6(m_5, m_q) + g_9(m_5, m_q) - h_3(m_5, m_q) \\
 & +h_6(m_5, m_q) - h_9(m_5, m_q)) + \frac{3}{2}[i_3(m_5, m_q) - 2j_3(m_g, m_q, m_5)] \\
 & \left. -18e_2(m_5, m_q, m_g) \right\} k_{4\alpha}, \tag{B.101}
 \end{aligned}$$

$$\begin{aligned}
 h_{2\alpha} = & -\left\{ 2g_1(m_g, m_q) - 2g_4(m_g, m_q) - \frac{4}{3}g_7(m_g, m_q) \right. \\
 & -\frac{2}{3}(h_1(m_g, m_q) - h_4(m_g, m_q) - h_7(m_g, m_q)) \\
 & +\frac{1}{3}(g_1(m_5, m_q) - g_4(m_5, m_q) + g_7(m_5, m_q) \\
 & -h_1(m_5, m_q) + h_4(m_5, m_q) - h_7(m_5, m_q)) \\
 & +\frac{3}{2}[i_4(m_5, m_q) - 2j_1(m_g, m_q, m_5) - 2j_1(m_5, m_q, m_g)] \\
 & -18[e_1(m_g, m_q, m_5) - e_1(m_5, m_q, m_g)] \\
 & \left. +\frac{3}{2}[l_1(m_q, m_5) - 2r_1(m_g, m_q, m_5) - 2r_{10}(m_g, m_q, m_5)] \right\} k_{1\alpha} \\
 & -\left\{ 2g_2(m_g, m_q) - 2g_5(m_g, m_q) - \frac{4}{3}g_8(m_g, m_q) \right. \\
 & -\frac{2}{3}(h_2(m_g, m_q) - h_5(m_g, m_q) - h_8(m_g, m_q)) \\
 & +\frac{1}{3}(g_2(m_5, m_q) - g_5(m_5, m_q) + g_8(m_5, m_q) \\
 & -h_2(m_5, m_q) + h_5(m_5, m_q) - h_8(m_5, m_q)) \\
 & +\frac{3}{2}[i_5(m_5, m_q) - 2j_2(m_g, m_q, m_5) - 2j_2(m_5, m_q, m_g)] \\
 & -3[r_2(m_g, m_q, m_5) + r_{11}(m_g, m_q, m_5)] \Big\} k_{3\alpha} \\
 & -\left\{ 2g_3(m_g, m_q) - 2g_6(m_g, m_q) - \frac{4}{3}g_9(m_g, m_q) \right. \\
 & -\frac{2}{3}(h_2(m_g, m_q) - h_5(m_g, m_q) - h_8(m_g, m_q))
 \end{aligned}$$

$$\begin{aligned}
& +\frac{1}{3}(g_3(m_5, m_q) - g_6(m_5, m_q) + g_9(m_5, m_q) \\
& - h_3(m_5, m_q) + h_6(m_5, m_q) - h_9(m_5, m_q)) \\
& +\frac{3}{2}[i_6(m_5, m_q) - 2j_3(m_g, m_q, m_5) - 2j_3(m_5, m_q, m_g)] \\
& -18[e_3(m_g, m_q, m_5) - e_3(m_5, m_q, m_g)] \\
& +\frac{3}{2}[l_3(m_q, m_5) - 2r_3(m_g, m_q, m_5) - 2r_{12}(m_g, m_q, m_5)]\}k_{4\alpha}, \text{ (B.102)}
\end{aligned}$$

$$\begin{aligned}
h_{3\alpha} = & -\frac{1}{6}(g_1(m_g, m_q) + 4g_4(m_g, m_q) - 8g_7(m_g, m_q) \\
& +4(h_1(m_g, m_q) + h_4(m_g, m_q) + h_7(m_g, m_q))) \\
& +\{\frac{1}{3}g_1(m_5, m_q) + g_4(m_5, m_q) - g_7(m_5, m_q) \\
& -h_1(m_5, m_q) - h_4(m_5, m_q) + h_7(m_5, m_q) \\
& +\frac{3}{2}[i_7(m_5, m_q) - 2j_1(m_g, m_q, m_5)] - 18e_1(m_g, m_q, m_5) \\
& +\frac{3}{2}l_4(m_q, m_5) - 3(t_1(m_5, m_q) + r_4(m_g, m_q, m_5) + r_{13}(m_g, m_q, m_5))\}k_{1\alpha} \\
& -\frac{1}{6}(g_2(m_g, m_q) + 4g_5(m_g, m_q) - 8g_8(m_g, m_q) + 4(h_2(m_g, m_q) \\
& +h_5(m_g, m_q) + h_8(m_g, m_q))) \\
& +\{\frac{1}{3}g_2(m_5, m_q) + g_5(m_5, m_q) - g_8(m_5, m_q) - h_2(m_5, m_q) \\
& -h_5(m_5, m_q) + h_8(m_5, m_q) \\
& +\frac{3}{2}[i_8(m_5, m_q) + 2j_2(m_g, m_q, m_5)] + \frac{3}{2}l_5(m_q, m_5) \\
& -3(t_2(m_5, m_q) + r_5(m_g, m_q, m_5) + r_{14}(m_g, m_q, m_5))\}k_{3\alpha} \\
& -\frac{1}{6}(g_3(m_g, m_q) + 4g_6(m_g, m_q) - 8g_9(m_g, m_q) + 4(h_3(m_g, m_q) \\
& +h_6(m_g, m_q) + h_9(m_g, m_q))) \\
& +\{\frac{1}{3}g_3(m_5, m_q) + g_6(m_5, m_q) - g_9(m_5, m_q) - h_4(m_5, m_q) \\
& -h_6(m_5, m_q) + h_9(m_5, m_q) \\
& +\frac{3}{2}[i_9(m_5, m_q) - 2j_3(m_g, m_q, m_5)] - 18e_3(m_g, m_q, m_5) + \frac{3}{2}l_6(m_q, m_5) \\
& -3(t_3(m_5, m_q) + r_6(m_g, m_q, m_5) + r_{15}(m_g, m_q, m_5))\}k_{4\alpha}, \text{ (B.103)}
\end{aligned}$$

$$\begin{aligned}
h_{4\alpha} = & -\{\frac{1}{6}(12g_1(m_g, m_q) - 12g_4(m_g, m_q) + g_7(m_g, m_q) \\
& +4(-h_1(m_g, m_q) + h_4(m_g, m_q) + h_7(m_g, m_q))) \\
& +\frac{1}{3}(-g_1(m_5, m_q) + g_4(m_5, m_q) + g_7(m_5, m_q) \\
& +h_1(m_5, m_q) - h_4(m_5, m_q) - h_7(m_5, m_q)) \\
& +\frac{3}{2}[i_{10}(m_5, m_q) - 2j_1(m_g, m_q, m_5)] - 18e_1(m_g, m_q, m_5) \\
& +\frac{3}{2}l_7(m_q, m_5) - 3(t_4(m_5, m_q) + r_7(m_g, m_q, m_5) + r_{16}(m_g, m_q, m_5))\}k_{1\alpha} \\
& -\{\frac{1}{6}(12g_2(m_g, m_q) - 12g_5(m_g, m_q) + g_8(m_g, m_q) \\
& +4(-h_2(m_g, m_q) + h_5(m_g, m_q) + h_8(m_g, m_q)))
\end{aligned}$$

$$\begin{aligned}
& +\frac{1}{3}(-g_2(m_5, m_q) + g_5(m_5, m_q) + g_8(m_5, m_q) \\
& + h_2(m_5, m_q) - h_5(m_5, m_q) - h_8(m_5, m_q)) \\
& +\frac{3}{2}[i_{11}(m_5, m_q) - 2j_2(m_g, m_q, m_5)] + \frac{3}{2}l_8(m_q, m_5) \\
& -3(t_5(m_5, m_q) + r_8(m_g, m_q, m_5) + r_{17}(m_g, m_q, m_5))\}k_{3\alpha} \\
& -\left\{\frac{1}{6}(12g_3(m_g, m_q) - 12g_6(m_g, m_q) + g_9(m_g, m_q) \right. \\
& +4(-h_3(m_g, m_q) + h_6(m_g, m_q) + h_9(m_g, m_q))) \\
& +\frac{1}{3}(-g_3(m_5, m_q) + g_6(m_5, m_q) + g_9(m_5, m_q) + h_3(m_5, m_q) \\
& -h_6(m_5, m_q) - h_9(m_5, m_q)) \\
& +\frac{3}{2}[i_{12}(m_5, m_q) - 2j_3(m_g, m_q, m_5)] - 18e_3(m_g, m_q, m_5) + \frac{3}{2}l_9(m_q, m_5) \\
& \left. -3(t_6(m_5, m_q) + r_9(m_g, m_q, m_5) + r_{18}(m_g, m_q, m_5))\right\}k_{4\alpha}. \quad (B.104)
\end{aligned}$$

These coefficients, $g(m_a, m_b), h(m_a, m_b), \dots$, are written in terms of Feynman integral as follows:

$$g_1(m_a, m_b) = -\alpha(m_b, m_a, 1, 3) + 3\alpha(m_b, m_a, 1, 2), \quad (B.105)$$

$$\begin{aligned}
g_2(m_a, m_b) &= g_8(m_a, m_b) = \frac{1}{2}g_3(m_a, m_b) = g_4(m_a, m_b) = g_7(m_a, m_b)) \\
&= -\alpha(m_b, m_a, 1, 3), \quad (B.106)
\end{aligned}$$

$$\begin{aligned}
g_5(m_a, m_b) &= g_6(m_a, m_b) = g_9(m_a, m_b) \\
&= -\alpha(m_b, m_a, 1, 3) + 3\alpha(m_b, m_a, 1, 2), \quad (B.107)
\end{aligned}$$

$$\begin{aligned}
h_1(m_a, m_b) &= h_2(m_a, m_b) = h_8(m_a, m_b)) \\
&= -m_b^2\alpha(m_b, m_a, 2, 3) + 3m_b^2\alpha(m_b, m_a, 2, 2), \quad (B.108)
\end{aligned}$$

$$h_3(m_a, m_b) = -2m_b^2\alpha(m_b, m_a, 2, 3), \quad (B.109)$$

$$\begin{aligned}
h_4(m_a, m_b) &= h_7(m_a, m_b) \\
&= -m_b^2\alpha(m_b, m_a, 2, 3), \quad (B.110)
\end{aligned}$$

$$\begin{aligned}
h_5(m_a, m_b) &= h_6(m_a, m_b) = h_9(m_a, m_b) \\
&= -m_b^2\alpha(m_b, m_a, 2, 3) + 3m_b^2\alpha(m_b, m_a, 2, 2), \quad (B.111)
\end{aligned}$$

$$\begin{aligned}
i_1(m_a, m_b) &= i_4(m_a, m_b) \\
&= 6\alpha(m_a, m_b, 1, 3) + f9\alpha(m_a, m_b, 1, 2), \quad (B.112)
\end{aligned}$$

$$\begin{aligned}
i_2(m_a, m_b) &= i_5(m_a, m_b) \\
&= 6\alpha(m_a, m_b, 1, 3) - 24\alpha(m_a, m_b, 1, 2), \quad (B.113)
\end{aligned}$$

$$\begin{aligned}
i_3(m_a, m_b) &= i_6(m_a, m_b) \\
&= 12\alpha(m_a, m_b, 1, 3) - 12\alpha(m_a, m_b, 1, 2), \quad (B.114)
\end{aligned}$$

$$i_7(m_a, m_b) = -12\alpha(m_a, m_b, 1, 3) + 3\alpha(m_a, m_b, 1, 2), \quad (B.115)$$

$$i_8(m_a, m_b) = 12\alpha(m_a, m_b, 1, 3) + 6\alpha(m_a, m_b, 1, 2), \quad (\text{B.116})$$

$$i_9(m_a, m_b) = -24\alpha(m_a, m_b, 1, 3), \quad (\text{B.117})$$

$$i_{10}(m_a, m_b) = -35\alpha(m_a, m_b, 1, 3), \quad (\text{B.118})$$

$$i_{11}(m_a, m_b) = -26\alpha(m_a, m_b, 1, 3) + 42\alpha(m_a, m_b, 1, 2), \quad (\text{B.119})$$

$$i_{12}(m_a, m_b) = -52\alpha(m_a, m_b, 1, 3) + 66\alpha(m_a, m_b, 1, 2), \quad (\text{B.120})$$

$$j_1(m_a, m_b, m_c) = 3m_b^2\beta(m_a, m_c, m_b, 2, 1, 3), \quad (\text{B.121})$$

$$j_2(m_a, m_b, m_c) = 6m_b^2\beta(m_a, m_c, m_b, 2, 1, 1) - 6m_b^2\beta(m_a, m_c, m_b, 2, 2, 0), \quad (\text{B.122})$$

$$j_3(m_a, m_b, m_c) = 6m_b^2\beta(m_a, m_c, m_b, 2, 0, 3) - 3m_b^2\beta(m_a, m_c, m_b, 2, 1, 3), \quad (\text{B.123})$$

$$e_1(m_a, m_b) = e_2(m_a, m_b) = \alpha(a, b, 1, 2), \quad (\text{B.124})$$

$$n_1(m_a, m_b) = 6\alpha(m_a, m_b, 1, 1) - 6\alpha(m_a, m_b, 1, 2), \quad (\text{B.125})$$

$$n_2(m_a, m_b) = n_8(m_a, m_b) = n_4(m_a, m_b) = \frac{1}{2}n_7(m_a, m_b) = n_{10}(m_a, m_b) = 12\alpha(m_a, m_b, 1, 3) - 24\alpha(m_a, m_b, 1, 2) + 12\alpha(m_a, m_b, 1, 1), \quad (\text{B.126})$$

$$n_5(m_a, m_b) = -24\alpha(m_a, m_b, 1, 2) - 24\alpha(m_a, m_b, 1, 1) + 48\alpha(m_a, m_b, 1, 0), \quad (\text{B.127})$$

$$n_6(m_a, m_b) = -24\alpha(m_a, m_b, 1, 3) + 84\alpha(m_a, m_b, 1, 2) - 27\alpha(a, b, 1, 1), \quad (\text{B.128})$$

$$n_{11}(m_a, m_b) = 6m_b^2\alpha(m_a, m_b, 2, 2) - 6m_b^2\alpha(m_a, m_b, 2, 1), \quad (\text{B.129})$$

$$n_{12}(m_a, m_b) = 6m_b^2\alpha(m_a, m_b, 2, 3) - 12m_b^2\alpha(m_a, m_b, 2, 2) + 6m_b^2\alpha(m_a, m_b, 2, 1), \quad (\text{B.130})$$

$$n_{14}(m_a, m_b) = 6m_b^2\alpha(m_a, m_b, 2, 1) - 12m_b^2\alpha(m_a, m_b, 2, 1) + 6m_b^2\alpha(m_a, m_b, 2, 1), \quad (\text{B.131})$$

$$t_1(m_a, m_b, m_c) = t_2(m_a, m_b, m_c) = -12m_a m_b \gamma(m_a, m_b, m_c, 2, 1, 0, 0) + 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 0), \quad (\text{B.132})$$

$$t_3(m_a, m_b, m_c) = 12m_a m_b \gamma(m_a, m_b, m_c, 2, 1, 0, 0) - 12m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 1) + 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 0), \quad (\text{B.133})$$

$$t_4(m_a, m_b, m_c) = 12m_a m_b \gamma(m_a, m_b, m_c, 2, 1, 0, 0), \quad (\text{B.134})$$

$$\begin{aligned}
t_5(m_a, m_b, m_c) = & 12m_a m_b \gamma(m_c, m_b, m_a, 2, 0, 1, 0) \\
& - 12m_a m_b \gamma(m_c, m_b, m_a, 2, 0, 1, 0) \\
& + 6m_a m_b \gamma(m_c, m_b, m_a, 2, 0, 0, 0), \quad (B.135)
\end{aligned}$$

$$\begin{aligned}
t_6(m_a, m_b, m_c) = & 12m_a m_b \gamma(m_c, m_b, m_a, 2, 1, 0, 0) \\
& - 12m_a m_b \gamma(m_c, m_b, m_a, 2, 0, 0, 1) \\
& + 12m_a m_b \gamma(m_c, m_b, m_a, 2, 0, 0, 0), \quad (B.136)
\end{aligned}$$

$$\begin{aligned}
l_1(m_b, m_c) = l_2(m_b, m_c) = \frac{1}{2}l_3(m_b, m_c) = l_7(m_b, m_c) \\
= 4\alpha(m_c, m_b, 1, 3), \quad (B.137)
\end{aligned}$$

$$\begin{aligned}
l_4(m_b, m_c) = l_5(m_b, m_c) = l_8(m_b, m_c) \\
= 4\alpha(m_c, m_b, 1, 3) - 6\alpha(m_c, m_b, 1, 2), \quad (B.138)
\end{aligned}$$

$$l_6(m_b, m_c) = 8\alpha(m_c, m_b, 1, 3) - 6\alpha(m_c, m_b, 1, 2), \quad (B.139)$$

$$l_9(m_b, m_c) = 8\alpha(m_c, m_b, 1, 3) - 12\alpha(m_c, m_b, 1, 2), \quad (B.140)$$

$$\begin{aligned}
r_1(m_a, m_b, m_c) = r_4(m_a, m_b, m_c) = -\frac{1}{2}r_7(m_a, m_b, m_c) \\
= -6m_a m_b \delta(m_a, m_b, m_c, 2, 1, 0, 0), \quad (B.141)
\end{aligned}$$

$$\begin{aligned}
r_2(m_a, m_b, m_c) = & 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 1, 0) \\
& - 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 1) \\
& + 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.142)
\end{aligned}$$

$$\begin{aligned}
r_3(m_a, m_b, m_c) = & 6m_a m_b \delta(m_a, m_b, m_c, 2, 1, 0, 0) \\
& - 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 1) \\
& + 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.143)
\end{aligned}$$

$$\begin{aligned}
r_5(m_a, m_b, m_c) = & 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 1, 0) \\
& - 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 1), \quad (B.144)
\end{aligned}$$

$$\begin{aligned}
r_6(m_a, m_b, m_c) = & 6m_a m_b \delta(m_a, m_b, m_c, 2, 1, 0, 0) \\
& - 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 1) \\
& + 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.145)
\end{aligned}$$

$$\begin{aligned}
r_8(m_a, m_b, m_c) = & 12m_a m_b \delta(m_a, m_b, m_c, 2, 0, 1, 0) \\
& - 12m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 1) \\
& - 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.146)
\end{aligned}$$

$$\begin{aligned}
r_9(m_a, m_b, m_c) = & 12m_a m_b \delta(m_a, m_b, m_c, 2, 1, 0, 0) \\
& - 12m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 1) \\
& - 12m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.147)
\end{aligned}$$

$$r_{10}(m_a, m_b, m_c) = 6m_a m_b \delta(m_a, m_b, m_c, 2, 1, 0, 0), \quad (B.148)$$

$$\begin{aligned}
 r_{11}(m_a, m_b, m_c) = & 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 1, 0) \\
 & - 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 1) \\
 & + 6m_a m_b \delta(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.149)
 \end{aligned}$$

$$\begin{aligned}
 r_{12}(m_a, m_b, m_c) = & 6m_a m_b \gamma(m_a, m_b, m_c, 2, 1, 0, 0) \\
 & - 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 1) \\
 & - 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.150)
 \end{aligned}$$

$$\begin{aligned}
 r_{13}(m_a, m_b, m_c) = & -12m_a m_b \gamma(m_a, m_b, m_c, 2, 1, 0, 0) \\
 & + 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.151)
 \end{aligned}$$

$$\begin{aligned}
 r_{14}(m_a, m_b, m_c) = & 12m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 1, 0) \\
 & - 12m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 1) \\
 & + 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.152)
 \end{aligned}$$

$$\begin{aligned}
 r_{15}(m_a, m_b, m_c) = & 12m_a m_b \gamma(m_a, m_b, m_c, 2, 1, 0, 0) \\
 & - 12m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 1) \\
 & + 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.153)
 \end{aligned}$$

$$\begin{aligned}
 r_{16}(m_a, m_b, m_c) = & 6m_a m_b \gamma(m_a, m_b, m_c, 2, 1, 0, 0) \\
 & - 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 0), \quad (B.154)
 \end{aligned}$$

$$\begin{aligned}
 r_{17}(m_a, m_b, m_c) = & 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 1, 0) \\
 & - 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 1), \quad (B.155)
 \end{aligned}$$

$$\begin{aligned}
 r_{18}(m_a, m_b, m_c) = & 6m_a m_b \gamma(m_a, m_b, m_c, 2, 1, 0, 0) \\
 & - 6m_a m_b \gamma(m_a, m_b, m_c, 2, 0, 0, 1), \quad (B.156)
 \end{aligned}$$

$$s_1(m_b, m_c) = 6\alpha(m_c, m_b, 1, 1) + 6\alpha(m_c, m_b, 1, 2), \quad (B.157)$$

$$\begin{aligned}
 s_2(m_b, m_c) = & 18\alpha(m_c, m_b, 1, 3) - 30\alpha(m_c, m_b, 1, 2) \\
 & + 6\alpha(m_c, m_b, 1, 1), \quad (B.158)
 \end{aligned}$$

$$\begin{aligned}
 s_3(m_b, m_c) = & -3\alpha(m_c, m_b, 1, 4) + 2\alpha(m_c, m_b, 1, 3) \\
 & + 6\alpha(m_c, m_b, 1, 1) - 5\alpha(m_c, m_b, 1, 0), \quad (B.159)
 \end{aligned}$$

$$\begin{aligned}
 s_4(m_b, m_c) = & 6\alpha(m_c, m_b, 1, 3) + 18\alpha(m_c, m_b, 1, 2) \\
 & - 12\alpha(m_c, m_b, 1, 1), \quad (B.160)
 \end{aligned}$$

$$\begin{aligned}
 s_5(m_b, m_c) = & 18m_b^2\alpha(m_c, m_b, 2, 3) - 36m_b^2\alpha(m_c, m_b, 2, 2) \\
 & + 18m_b^2\alpha(m_c, m_b, 2, 1) + 18m\alpha(m_c, m_b, 1, 3) \\
 & - 36\alpha(m_c, m_b, 1, 2) + 18\alpha(m_c, m_b, 1, 1), \quad (B.161)
 \end{aligned}$$

$$\begin{aligned}
s_6(m_b, m_c) = & -3m_b^2\alpha(m_c, m_b, 2, 4) + 2m_b^2\alpha(m_c, m_b, 2, 3) \\
& + 6m_b^2\alpha(m_c, m_b, 2, 1) - 5m_b^2\alpha(m_c, m_b, 2, 0) \\
& - 3\alpha(m_c, m_b, 1, 4) + 2\alpha(m_c, m_b, 1, 3) \\
& + 6\alpha(m_c, m_b, 1, 1) - 5\alpha(m_c, m_b, 1, 0), \quad (B.162)
\end{aligned}$$

$$\begin{aligned}
s_7(m_b, m_c) = & -6m_b^2\alpha(m_c, m_b, 2, 2) + 12m_b^2\alpha(m_c, m_b, 2, 1) \\
& - 6m_b^2\alpha(m_c, m_b, 2, 0) - 6\alpha(m_c, m_b, 1, 2) \\
& + 12\alpha(m_c, m_b, 1, 1) - 6\alpha(m_c, m_b, 1, 0), \quad (B.163)
\end{aligned}$$

$$s_8(m_a, m_b, m_c) = 6m_a m_b \beta(m_b, m_c, m_a, 2, 0, 0), \quad (B.164)$$

$$\begin{aligned}
s_9(m_a, m_b, m_c) = & -3m_a m_b \beta(m_b, m_c, m_a, 2, 0, 2) \\
& + 3m_a m_b \beta(m_b, m_c, m_a, 2, 0, 0), \quad (B.165)
\end{aligned}$$

$$\begin{aligned}
s_{10}(m_a, m_b, m_c) = & -12m_a m_b \beta(m_b, m_c, m_a, 2, 0, 2) \\
& + 6m_a m_b \beta(m_b, m_c, m_a, 2, 0, 1) \\
& + 12m_a m_b \beta(m_b, m_c, m_a, 2, 1, 1) \\
& - 12m_a m_b \beta(m_b, m_c, m_a, 2, 1, 0) \\
& + 6m_a m_b \beta(m_b, m_c, m_a, 2, 0, 0), \quad (B.166)
\end{aligned}$$

$$\begin{aligned}
s_{11}(m_a, m_b, m_c) = & -6m_a m_b \beta(m_b, m_c, m_a, 2, 0, 2) \\
& + 6m_a m_b \beta(m_b, m_c, m_a, 2, 0, 1), \quad (B.167)
\end{aligned}$$

$$\begin{aligned}
s_{12}(m_a, m_b, m_c) = & -6m_a m_b \beta(m_b, m_a, m_c, 2, 0, 1) \\
& + 6m_a m_b \beta(m_b, m_a, m_c, 2, 0, 0), \quad (B.168)
\end{aligned}$$

$$\begin{aligned}
s_{13}(m_a, m_b, m_c) = & -3m_a m_b \beta(m_b, m_a, m_c, 2, 0, 2) \\
& + 3m_a m_b \beta(m_b, m_a, m_c, 2, 0, 0), \quad (B.169)
\end{aligned}$$

$$\begin{aligned}
s_{14}(m_a, m_b, m_c) = & -12m_a m_b \beta(m_b, m_a, m_c, 2, 0, 2) \\
& + 6m_a m_b \beta(m_b, m_a, m_c, 2, 0, 1) \\
& + 12m_a m_b \beta(m_b, m_a, m_c, 2, 1, 1) \\
& - 12m_a m_b \beta(m_b, m_a, m_c, 2, 1, 0) \\
& + 6m_a m_b \beta(m_b, m_a, m_c, 2, 0, 0), \quad (B.170)
\end{aligned}$$

$$\begin{aligned}
s_{15}(m_a, m_b, m_c) = & -6m_a m_b \beta(m_b, m_a, m_c, 2, 0, 2) \\
& + 5m_a m_b \beta(m_b, m_a, m_c, 2, 0, 1), \quad (B.171)
\end{aligned}$$

and other coefficients are zero.

REFERENCES

- [1] C.J.C. Burges, H.J. Schnitzer, *Nucl. Phys.* **B228**, 464 (1983).
- [2] W. Buchmuller, D. Wyler, *Nucl. Phys.* **B268**, 621 (1986).
- [3] C. Arzt, M.B. Einhorn, J. Wudka, *Nucl. Phys.* **B433**, 41 (1995).
- [4] J.A. Aguilar-Saavedra, *Nucl. Phys.* **B812**, 181 (2009) [[arXiv:0811.3842](#) [hep-ph]].
- [5] J.A. Aguilar-Saavedra, *Nucl. Phys.* **B843**, 638 (2011) [[arXiv:1008.3562](#) [hep-ph]].
- [6] B. Grzadkowski, M. Iskrzynski, M. Misiak, J. Rosiek, *J. High Energy Phys.* **1010**, 085 (2010) [[arXiv:1008.4884](#) [hep-ph]].
- [7] H.P. Nilles, *Phys. Rep.* **110**, 1 (1984).
- [8] T. Appelquist, H.-C. Cheng, B.A. Dobrescu, *Phys. Rev.* **D64**, 035002 (2001) [[arXiv:hep-ph/0012100](#)].
- [9] M. Schmaltz, D. Tucker-Smith, *Annu. Rev. Nucl. Part. Sci.* **55**, 229 (2005) [[arXiv:hep-ph/0502182](#)].
- [10] N. Haba, K. Kaneta, S. Matsumoto, T. Nabeshima, [arXiv:1106.6106](#) [hep-ph].