

# TESTING NEW STRONGLY INTENSIVE MEASURES OF TRANSVERSE MOMENTUM FLUCTUATIONS

KATARZYNA GREBIESZKOW

Faculty of Physics, Warsaw University of Technology  
Koszykowa 75, 00-662 Warszawa, Poland  
kper1@if.pw.edu.pl

*(Received February 27, 2012; revised version received April 5, 2012)*

Recently, the new strongly intensive measures of fluctuations  $\Delta$  and  $\Sigma$  have been proposed. In this publication their properties are tested using an example of event-by-event transverse momentum fluctuations. The obtained values are compared to the long used  $\Phi$  measure of  $p_T$  fluctuations. Several tests are performed within data produced by fast generators, as well as by the UrQMD model. The UrQMD calculations are presented for the systems and energies which are planned to be studied in the NA61/SHINE experiment at the CERN Super Proton Synchrotron.

DOI:10.5506/APhysPolB.43.1333

PACS numbers: 25.75.Gz, 25.75.-q

## 1. Introduction

The main motivation of colliding relativistic heavy ions is to create and study the properties of the system composed by deconfined quarks and gluons (QGP). The data suggest that the energy threshold for deconfinement (onset of deconfinement) is located at low SPS energies [1, 2]. The phase diagram of strongly interacting matter can be presented in terms of temperature  $T$  and baryochemical potential  $\mu_B$ . The bulk of theoretical calculations suggests that the phase boundary between hadrons and QGP is of first order at large values of  $\mu_B$ , ending in a critical point of second order and then turning into a continuous rapid transition at low  $\mu_B$ . Lattice QCD calculations indicate that the critical point (CP) can be located in the SPS energy range, *i.e.*  $T^{\text{CP}} = 162 \pm 2$  MeV,  $\mu_B^{\text{CP}} = 360 \pm 40$  MeV [3] or  $(T^{\text{CP}}, \mu_B^{\text{CP}}) = (0.927(5)T_c, 2.60(8)T_c) = (\sim 157, \sim 441)$  MeV [4], where  $T_c$  is the critical temperature of hadron gas  $\leftrightarrow$  QGP transition at vanishing baryochemical potential.

The analysis of dynamical fluctuations can be an important tool for localizing the phase boundary and the critical point. In particular, significant transverse momentum and multiplicity fluctuations are expected to appear for systems freezing-out close to CP [5]. The position of the freeze-out point in the phase diagram can be moved by varying the collision energy and the size of the colliding nuclei [6]. A non-monotonic evolution of fluctuations with such parameters can serve as a signature for the phase transition and the critical point.

In fact, these considerations motivated an extensive program of fluctuation studies at the SPS and RHIC accelerators. The NA49 [7] experiment reported non-monotonic behaviour of average  $p_T$  and multiplicity fluctuations at the top SPS energy [8]. This intriguing result might be a first hint of the critical point. Therefore, the efforts to look for CP will be continued within the NA61/SHINE [9] project, where a 2D (energy and system size) scan of the phase diagram will be performed. Figure 1 shows the hypothetical chemical freeze-out points in the NA61 experiment.




Fig. 1. Positions of chemical freeze-out points obtained within hadron gas model by fitting NA49 data [6] (blue squares). Circles are those expected in NA61. Taken from [10].

Dynamical fluctuations can be measured by use of event-by-event methods. However, while measuring event-by-event fluctuations in nucleus + nucleus ( $A+A$ ) collisions, one should remember about a trivial source of fluctuations caused by event-by-event changes of the collision geometry. Therefore, a suitable choice of statistical tools for the study of event-by-event fluctuations is really important. In Ref. [11] a strongly intensive measure  $\Phi$  was introduced. In a superposition model,  $\Phi$  does not depend on the number of “sources” (*e.g.* wounded nucleons in the Wounded Nucleon Model [12]) composing  $A+A$  collision (intensive measure) and on the fluctuations of this number of sources (strongly intensive measure). In addition, in thermody-

namical models  $\Phi$  does not depend on volume and volume fluctuations provided that temperature and chemical potential are constant. The  $\Phi$  measure was already used by the NA49 experiment to calculate transverse momentum fluctuations ( $\Phi_{p_T}$ ) [13, 14]. Recently, two new classes of strongly intensive measures have been proposed:  $\Delta$  and  $\Sigma$  [15]. In fact, previously proposed  $\Phi$  belongs to  $\Sigma$ -family measures. In heavy ion experiments, the use of strongly intensive measures of fluctuations, such as  $\Phi$ ,  $\Delta$  and  $\Sigma$ , can be a remedy for an imperfect centrality selection of  $A + A$  collisions. Therefore, the NA49 experiment was able to use relatively wide centrality bins while studying  $\Phi_{p_T}$  measure (up to 0–15% centrality), whereas the analysis of multiplicity fluctuations had to be limited to 1% most central interactions only [8].

In this paper, first basic tests of the newly proposed  $\Delta$  and  $\Sigma$  measures will be presented for transverse momentum fluctuations. The obtained values will be compared to the long used  $\Phi$  measure of  $p_T$  fluctuations. Several effects have been studied for events generated by use of so-called fast generators. Moreover, the analysis within a much more complex UrQMD3.3 model [16] will be shown. The UrQMD calculations have been done for the systems and energies which are planned to be studied in the CERN NA61/SHINE experiment (see Fig. 1).

## 2. Strongly intensive measures

By intensive, we call quantities which do *not* depend on the volume of the system. In contrary, extensive quantities (for example mean multiplicity or variance of multiplicity distribution) are proportional to the system volume. Note, that it is useful to extend the notion of intensive and extensive quantities to the Wounded Nucleon Model. Namely, the intensive can be called quantities which are independent of the number of wounded nucleons, and extensive those which are proportional to the number of wounded nucleons. The ratio of two extensive quantities is an intensive quantity [15]. Therefore, the ratio of mean multiplicities, as well as the scaled variance of multiplicity distribution  $\omega = (\langle N^2 \rangle - \langle N \rangle^2) / \langle N \rangle$ , are intensive measures. In fact, due to its intensity property,  $\omega$  measure is quite commonly used to determine multiplicity fluctuations in heavy ion experiments.

There is one more important problem which one should not forget about. In high energy heavy ion collisions the volume of the produced matter cannot be fixed. In fact, the system volume changes significantly from event to event. Therefore, it is very important to be able to measure the properties of the created matter independently of its volume fluctuations. The quantities which allow this are called strongly intensive measures. They do not depend on the volume and on volume fluctuations.

Mean multiplicities ratios are both intensive and strongly intensive measures. The situation is, however, more difficult for fluctuation analysis. The scaled variance of multiplicity distribution is an intensive measure but not strongly intensive. Quite long ago, a strongly intensive measure  $\Phi$  was first introduced [11]. In the recent paper [15] it was shown that there are at least two families of strongly intensive measures:  $\Delta$  and  $\Sigma$ . The previously known  $\Phi$  measure belongs to  $\Sigma$ -type family. They can be calculated for any two extensive quantities. In this paper,  $\Delta$ ,  $\Sigma$ , and  $\Phi$  measures calculated for particle multiplicity,  $N$ , and sum of their transverse momenta modules will be tested.

### 2.1. $\Phi_{p_T}$ measure

The  $\Phi$  measure [11] was already successfully used by NA49 to determine transverse momentum fluctuations ( $\Phi_{p_T}$ ) [13, 14]. Following the authors of [11] one defines the single-particle variable  $z_{p_T} = p_T - \overline{p_T}$  with the bar denoting averaging over the single-particle inclusive distribution. As seen  $\overline{z_{p_T}} = 0$ . Further, one introduces the event variable  $Z_{p_T}$ , which is a multi-particle analog of  $z_{p_T}$ , defined as  $Z_{p_T} = \sum_{i=1}^N (p_{T_i} - \overline{p_T})$ , where the summation runs over particles in a given event. Note, that  $\langle Z_{p_T} \rangle = 0$ , where  $\langle \dots \rangle$  represents averaging over events. Finally, the  $\Phi_{p_T}$  measure is defined as

$$\Phi_{p_T} = \sqrt{\frac{\langle Z_{p_T}^2 \rangle}{\langle N \rangle}} - \sqrt{z_{p_T}^2}. \quad (1)$$

$\Phi_{p_T}$  is a strongly intensive measure and, therefore, if  $A + A$  collision is represented by an incoherent superposition of independent nucleon+nucleon ( $N + N$ ) interactions (superposition model), then  $\Phi_{p_T}$  has a constant value, the same for  $A + A$  and  $N + N$  interactions. This implies that, in particular,  $\Phi_{p_T}$  does not depend on the impact parameter (centrality), if the  $A + A$  collision is a simple superposition of  $N + N$  interactions.

Another property of this measure is that  $\Phi_{p_T}$  vanishes when the system consists of particles that are emitted independently (no inter-particle correlations) and the single particle momentum spectrum is independent of multiplicity. In contrary,  $\Delta$  and  $\Sigma$  measures, which will be shown below, do not assume zero values for independent particle production.

### 2.2. $\Delta^{XN}$ and $\Sigma^{XN}$ measures

Let  $A$  and  $B$  be two fluctuating extensive quantities. Then  $\Delta^{AB}$  and  $\Sigma^{AB}$  can be defined [15]:

$$\Delta^{AB} = \langle C \rangle^{-1} [\langle B \rangle \omega_A - \langle A \rangle \omega_B], \quad (2)$$

$$\Sigma^{AB} = \langle C \rangle^{-1} [\langle B \rangle \omega_A + \langle A \rangle \omega_B - 2(\langle AB \rangle - \langle A \rangle \langle B \rangle)] , \quad (3)$$

where

$$\omega_A = \frac{\langle A^2 \rangle - \langle A \rangle^2}{\langle A \rangle} , \quad (4)$$

and

$$\omega_B = \frac{\langle B^2 \rangle - \langle B \rangle^2}{\langle B \rangle} \quad (5)$$

are the scaled variances of two fluctuating extensive quantities  $A$  and  $B$ .  $\langle C \rangle$  is the average of any extensive quantity *e.g.*,  $\langle A \rangle$  or  $\langle B \rangle$ .

There is an important difference between  $\Delta^{AB}$  and  $\Sigma^{AB}$ . Only the first two moments:  $\langle A \rangle$ ,  $\langle B \rangle$ , and  $\langle A^2 \rangle$ ,  $\langle B^2 \rangle$  are required to calculate  $\Delta^{AB}$ , whereas  $\Sigma^{AB}$  includes the correlation term  $\langle AB \rangle - \langle A \rangle \langle B \rangle$ . Thus  $\Delta^{AB}$  and  $\Sigma^{AB}$  measures can be sensitive to several physics effects in different ways. In publication [15], all strongly intensive quantities including correlation term are named the  $\Sigma$  family, and those including only mean values and variances, the  $\Delta$  family. The already used  $\Phi$  measure belongs to  $\Sigma$ -type family. The definitions of new quantities  $\Delta^{AB}$  and  $\Sigma^{AB}$  are, however, more general because one can use here *any* two extensive measures  $A$  and  $B$ , whereas in the original definition of  $\Phi$ , one of them was fixed to be particle multiplicity. Nevertheless, in this paper one of the extensive measures will be again multiplicity, in order to compare the results with known measure  $\Phi$ . Finally, one should mention that  $\Delta^{AB}$  and  $\Sigma^{AB}$  have also different properties with respect to exchange  $A$  and  $B$ :  $\Sigma^{AB} = \Sigma^{BA}$  and  $\Delta^{AB} = -\Delta^{BA}$ .

For the analysis of transverse momentum fluctuations one can use [15]:  $A \equiv X = \sum_{i=1}^N x_i$  (where  $x_i \equiv p_{T,i}$  and the summation runs over all *accepted* particles in a given event),  $B \equiv N$ ,  $C \equiv N$ , and  $X_2 = \sum_{i=1}^N x_i^2$ . Then we obtain

$$\begin{aligned} \Delta^{XN} &= \frac{1}{\langle N \rangle} [\langle N \rangle \omega_X - \langle X \rangle \omega_N] \\ &= \frac{1}{\langle N \rangle} \left[ \langle N \rangle \left( \frac{\langle X^2 \rangle - \langle X \rangle^2}{\langle X \rangle} \right) - \langle X \rangle \left( \frac{\langle N^2 \rangle - \langle N \rangle^2}{\langle N \rangle} \right) \right] \end{aligned} \quad (6)$$

and

$$\Sigma^{XN} = \frac{1}{\langle N \rangle} [\langle N \rangle \omega_X + \langle X \rangle \omega_N - 2(\langle XN \rangle - \langle X \rangle \langle N \rangle)] . \quad (7)$$

The  $\Sigma^{XN}$  measure can be also expressed [15] using known  $\Phi_{p_T}$  quantity

$$\Sigma^{XN} = \frac{\Phi_{p_T}^2 \langle N \rangle}{\langle X \rangle} + \frac{\langle X_2 \rangle}{\langle X \rangle} - \frac{\langle X \rangle}{\langle N \rangle} . \quad (8)$$

In this paper,  $\Sigma^{XN}$  is calculated using Eqs. (8). The statistical errors on  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  were estimated as follows. The whole sample of events was divided into 30 independent sub-samples. The values of  $\Phi_{p_T}$ ,  $\Delta^{XN}$ , and  $\Sigma^{XN}$  were evaluated for each sub-sample and the dispersions ( $D_\Phi$ ,  $D_\Delta$ , and  $D_\Sigma$ ) of the results were then calculated. The statistical error of  $\Phi_{p_T}$  ( $\Delta^{XN}$  or  $\Sigma^{XN}$ ) is taken to be equal to  $D_\Phi/\sqrt{30}$  ( $D_\Delta/\sqrt{30}$  or  $D_\Sigma/\sqrt{30}$ ).

### 3. Results of fast generators

#### 3.1. Independent particle production

The basic properties of  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  were tested by use of Monte Carlo models (so-called fast generators). Each interaction (event) was composed by a given number of identical single sources. For each source, the number of particles was generated from the Poisson distribution with a mean value of 5. The particle transverse momentum was generated from exponential transverse mass spectrum with inverse slope parameter (“temperature”)  $T = 150$  MeV. The number of sources composing an event was either constant (circles in Fig. 2) or selected from Poisson (triangles) or from Negative Binomial distribution (squares). For Negative Binomial distribution its dispersion  $\sqrt{\text{Var}(N_S)}$  was large and taken to be equal  $\langle N_S \rangle / 2$ .




Fig. 2.  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  versus number or mean number of sources composing one event. Example of independent particle production.

Figure 2 presents  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  measures versus the number or the mean number of sources composing one event. As the simulation performed here represents independent particle production,  $\Phi_{p_T}$  measure is consistent with zero. In contrary,  $\Delta^{XN}$  and  $\Sigma^{XN}$  have non-zero values because their definitions do not assume zero for independent particle production. The circles nicely confirm the intensity property of all three measures, whereas the triangles and the squares show that these quantities are also strongly intensive (do not depend on the system size and the system size fluctuations). One can also mention here that for a constant number of sources per event (circles in Fig. 2) the scaled variance of multiplicity distribution

$\omega = (\langle N^2 \rangle - \langle N \rangle^2) / \langle N \rangle$  is close to 1 in the whole range of the horizontal axis. For Poisson number of sources distribution  $\omega$  is approximately 6 also in the whole range of the mean number of sources per event. For Negative Binomial distribution of the number of sources  $\omega$  increases from about 7 (for on average 5 sources per event), through 126 (for on average 100 sources per event) up to approximately 1000 (for on average 800 sources per event). Figure 2 shows that within statistical errors (which can be sometimes large) all three measures are strongly intensive even if the multiplicity distribution is extremely and abnormally wide.

### 3.2. “Temperature” fluctuations

In the next simulation, for each single source the number of particles was again selected from the Poisson distribution with a mean value of 5. The particle transverse momentum was generated from exponential transverse mass spectrum with average inverse slope parameter  $\langle T \rangle = 150$  MeV. The  $T$  parameter was generated separately for each single source (source-by-source  $T$  fluctuations) from Gaussian shape with dispersion  $\sigma_T = 25$  MeV. Finally, the number of sources composing an event was generated from the Poisson distribution. The results are presented in Fig. 3. The effect of “temperature” fluctuations results in positive  $\Phi_{p_T}$  values and in higher (then in Fig. 2)  $\Delta^{XN}$  and  $\Sigma^{XN}$  values. As the sources are identical (superposition model)  $\Phi_{p_T}$ ,  $\Delta^{XN}$ , and  $\Sigma^{XN}$  measures do not depend on the mean number of sources composing an event. Therefore, Fig. 3 also confirms that all three measures are strongly intensive.




Fig. 3.  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  versus mean number of sources composing one event. Simulation of the effect of source-by-source inverse slope parameter ( $T$ ) fluctuations.

In the next step, source-by-source  $T$  fluctuations from the previous simulation were replaced by event-by-event  $T$  fluctuations ( $T$  parameter was the same for all sources composing a given event but was varied between events). Here, the  $T$  parameter was generated separately for each event from Gaussian shape with  $\sigma_T = 25$  MeV. Again, the number of sources

composing an event was generated from the Poisson distribution. The results are shown in Fig. 4. In this simulation the values of  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  increase with increasing the number of sources composing an event. It suggests that in real experimental data the effect of event-by-event “temperature” fluctuations should be better detectable for more central (or for heavier  $A$ ) collisions. One should also mention here that the relationship between temperature and multiplicity (or volume) fluctuations was studied in [17, 18, 19] .




Fig. 4.  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  versus mean number of sources composing one event. Simulation of the effect of event-by-event inverse slope parameter ( $T$ ) fluctuations.

In the last simulation, different widths of  $T$  parameter distribution were considered. As usually, for each single source the number of particles was selected from Poisson with a mean value of 5. The particle transverse momentum was generated from exponential transverse mass shape with average inverse slope parameter  $\langle T \rangle = 150$  MeV. The  $T$  parameter was varied from event to event following Gaussian distribution with dispersion  $\sigma_T$ . In order to avoid negative  $T$  values, only events within  $T = 150 \pm 3\sigma_T$  MeV were accepted. Finally, the number of sources composing an event was generated from the Poisson distribution with a mean value of 100. The results are shown in Fig. 5. As expected, the values of all three fluctuation measures increase when event-by-event “temperature” fluctuations are stronger (higher  $\sigma_T$ ).

The influence of “temperature” fluctuations on transverse momentum fluctuations was already studied in [20]. Following the authors of [20], one can easily derive that event-by-event “temperature” fluctuations result in  $\Phi_{p_T}$  equal  $\sqrt{\sigma_T^2(2 + 4\langle N^2 \rangle / \langle N \rangle) + 2\langle T \rangle^2} - \sqrt{6\sigma_T^2 + 2\langle T \rangle^2}$ . For the scaled variance of multiplicity distribution  $\omega \approx 6$  (taken from Monte Carlo simulations in Fig. 5) the formula can be rewritten as:  $\Phi_{p_T} = \sqrt{2\sigma_T^2(13 + 2\langle N \rangle) + 2\langle T \rangle^2} - \sqrt{6\sigma_T^2 + 2\langle T \rangle^2}$ . This function, for  $\langle T \rangle = 150$  MeV and  $\langle N \rangle = 500$ , is drawn as a solid line in Fig. 6, together with results from Monte Carlo






Fig. 5.  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  versus width of  $T$  parameter distribution.

simulations (taken from Fig. 5, left). The Monte Carlo results are in a very good agreement with the analytical formula. In publication [20], it is also suggested that for sufficiently small  $T$  parameter fluctuations, it is when  $\langle N \rangle \langle T \rangle^2 \gg \langle N^2 \rangle (\langle T^2 \rangle - \langle T \rangle^2)$ , and for Poisson ( $\omega = 1$ ) multiplicity distribution  $\Phi_{p_T}$  simplifies to the formula  $\sqrt{2} \langle N \rangle \frac{\langle T^2 \rangle - \langle T \rangle^2}{\langle T \rangle}$ . This simplified function, for  $\langle T \rangle = 150$  MeV and  $\langle N \rangle = 500$ , is drawn as a dashed line in Fig. 6. As seen, the simplified function is indeed valid only for small  $T$  parameter fluctuations and starts to deviate from Monte Carlo simulations for  $\sigma_T / \langle T \rangle$  higher than approximately 3 percent.




Fig. 6.  $\Phi_{p_T}$  versus relative width of  $T$  parameter distribution. See the text for details.

#### 4. Results of the UrQMD model

In order to calculate  $\Delta^{XN}$ ,  $\Sigma^{XN}$  and  $\Phi_{p_T}$ , the UrQMD3.3 model [16] was used. The model was already used to study event-by-event fluctuations of average transverse momentum [21, 22], charged particle ratio [23], and particle number [24]. The UrQMD generator is a microscopic transport model producing hadrons via formation, decay, and rescattering of reso-

nances and strings. The UrQMD approach simulates multiple interactions of both target/beam nucleons and newly produced particles, excitation, and fragmentation of colour strings and the formation and decay of hadronic resonances. The samples of minimum bias events were generated for the systems which will be analysed in the NA61/SHINE experiment, namely Be+Be, Ar+Ca, Xe+La at beam energies 13A, 20A, 30A, 40A, 80A, and 158A GeV (Fig. 1). These beam energies correspond to centre-of-mass energies for nucleon + nucleon pair  $\sqrt{s_{NN}}$  equal 5.12, 6.27, 7.62, 8.76, 12.3, and 17.3 GeV, respectively. Only inelastic  $A + A$  collisions were considered, it is the generated events with the number of collisions equal zero and events with the number of inelastic collisions equal zero were removed from the sample. The remaining events which had the final state multiplicities equal to the sum of the nucleons in both colliding nuclei were also rejected.

From such prepared minimum bias events only 20% of the most central collisions were selected (in agreement to NA61 data taking plans). This centrality selection corresponds to impact parameter cuts in UrQMD:  $b < 1.96$  fm for Be+Be,  $b < 3.81$  fm for Ar+Ca, and  $b < 5.71$  fm for Xe+La.

#### 4.1. Influence of high- $p_T$ particles (at SPS energies)

Figure 7 presents  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  measures obtained for all charged particles produced in UrQMD. Open points are those without any kinematic restrictions. Full points correspond to results with  $0.005 < p_T < 1.5$  GeV/c cut. Such cut was applied by the NA49 experiment [13, 14, 25] in order to eliminate the possible effect originating from hard interactions (the lower cut was due to momentum resolution of the detector). Figure 8 shows mean multiplicities of charged particles with and without  $0.005 < p_T < 1.5$  GeV/c cut. Although at SPS energies there is only a small fraction of particles coming from hard interactions (compare open and full points in Fig. 8) such high- $p_T$  hadrons can increase the values of all considered fluctuation measures (Fig. 7). For Xe+La collisions at the SPS this increase is on the level of 15–19% for all three fluctuation measures.




Fig. 7.  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  measured for all charged particles produced in 20% most central  $A + A$  collisions in UrQMD.




Fig. 8. Mean multiplicities of charged particles produced in 20% most central  $A+A$  collisions in UrQMD.

In Fig. 7 one also observes a slight decrease of  $\Phi_{p_T}$  and  $\Sigma^{XN}$  with increasing energy, while, in contrary,  $\Delta^{XN}$  increases. It is also worth to notice that the values of all three measures are lower for lighter systems (increase when going from Be+Be to Xe+La). In a simple superposition model no system size dependence is expected. However, the increase of  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  for heavier systems may be due to event-by-event impact parameter fluctuations. The sample of 0–20% central collisions may be not narrow enough (especially for the heaviest Xe+La) and the characteristics of more and less central collisions may be significantly different leading to increased event-by-event fluctuations. In the next sections, it will be shown that indeed the values of fluctuation measures are smaller when the centrality is restricted to 7.2% most central data.

#### 4.2. Influence of hadrons different than $\pi$ , $K$ , $\text{anti}(p)$

In the event-by-event analysis of experimental data [13, 14, 25] typically only charged particles originating from the main vertex are used. It practically means that only main vertex pions, protons, kaons and their antiparticles are used in the analysis, because particles coming from the decays of, for example,  $\Lambda$ ,  $\phi$ ,  $\Sigma$ ,  $\Xi$  and  $\Omega$  are believed to be rejected by a set of track selection criteria. The time scale of the simulations performed within the UrQMD model did not allow for weak decays, therefore the UrQMD analysis of charged pions, protons, kaons and their antiparticles can be directly compared to the analysis of experimental data.

Figure 9 compares  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  calculated for all charged particles produced in UrQMD (full symbols) and for only charged pions, protons, kaons and their antiparticles (open symbols). The small difference between these two cases is a reflection of the fact that the majority of final state particles produced in relativistic heavy ion collisions are pions, (anti)protons and kaons.




Fig. 9.  $\Phi_{p_T}$ ,  $\Delta^{XN}$  and  $\Sigma^{XN}$  measured for all charged particles and for charged pions, kaons and (anti)protons produced in 20% most central  $A + A$  collisions in UrQMD.

#### 4.3. $\Phi_{p_T}$ -dependence on energy and charge combination

Figure 10 shows the energy dependence of  $\Phi_{p_T}$  for different charge combinations: all charged particles, negatively charged and positively charged. The open symbols in the right panel represent positively charged particles where protons were removed from the sample. As already discussed in Fig. 7, a decrease of  $\Phi_{p_T}$  with energy can be observed for all charged particles. This decrease is even more pronounced when looking at positively charged particles (Fig. 10, right). On the contrary, negatively charged particles show




Fig. 10. Energy dependence of  $\Phi_{p_T}$  for different charge combinations of particles produced in 20% most central  $A + A$  collisions in UrQMD.

values close to zero and only a very small increase with energy (see inset of Fig. 10, middle) can be observed for heavier systems. The most important difference between positively charged and negatively charged particles is the presence of protons in the sample of positively charged. Therefore, the  $\Phi_{p_T}$  values were calculated also for positively charged particles without protons (open points in the right panel). Their values are consistent with those for negatively charged particles thus confirming that the effect indeed originates from protons. It is also worth to remind that the highest effect from protons is observed for the lowest energies, where the fraction of protons is the

highest. For Xe+La UrQMD data the fraction of protons is about 35% of all charged for 13A GeV and 15% of all charged for 158A GeV. For Ar+Ca data these numbers are 36% and 18% for 13A GeV and 158A GeV, respectively, and for Be+Be collisions 48% and 24%.

In the next set of plots (Fig. 11), the  $\Phi_{p_T}$  values are shown for forward rapidity only ( $1.1 < y_\pi^* < 2.6$ , where  $y_\pi^*$  is the particle rapidity calculated in the centre-of-mass reference system assuming pion mass). The same rapidity cut was used by the NA49 experiment in the analysis of transverse momentum [13, 14] and multiplicity [25] fluctuations. Additionally, only particles with  $y_p^* < y_{\text{beam}}^* - 0.5$  were accepted ( $y_{\text{beam}}^*$  is the rapidity of the beam calculated in the centre-of-mass reference system). This cut allows to get rid of the effect of event-by-event impact parameter fluctuations when restricting the analysis to forward rapidity only. The details of this cut were discussed in [22] and, in particular, it was shown that event-by-event fluctuations of the number of protons in the forward hemisphere and the number of protons that are closer to the production region can lead to increased  $\Phi_{p_T}$  values.




Fig. 11. Energy dependence of  $\Phi_{p_T}$  at forward rapidity for different charge combinations of particles produced in 20% most central  $A + A$  collisions in UrQMD. Additional cut  $y_p^* < y_{\text{beam}}^* - 0.5$  was applied.

Figure 11 shows that  $\Phi_{p_T}$  values measured at forward rapidity are much smaller than those for complete rapidity region (see Fig. 10 for a comparison). Also the mean multiplicities are significantly smaller (Fig. 12). The negatively charged particles show  $\Phi_{p_T}$  consistent with zero for all three studied systems (the slight increase with energy observed for complete rapidity region is not seen here any more). The positively charged particles also result in  $\Phi_{p_T}$  values close to zero and a weak increase with increasing energy can be observed for Xe+La system only.

#### 4.3.1. Centrality restrictions

Figure 10 suggests that for 20% most central interactions the increased  $\Phi_{p_T}$  values at lower SPS energies are mainly due to the significant fraction




Fig. 12. Mean multiplicities of all charged, negatively charged and positively charged particles produced at forward rapidity in 20% most central  $A + A$  collisions in UrQMD. Additional cut  $y_p^* < y_{\text{beam}}^* - 0.5$  was applied.

of protons present in the samples. The centrality bin 0–20% is a relatively wide one and, therefore, the observed increase of  $\Phi_{p_T}$  may be due to event-by-event impact parameter fluctuations (corresponding to event-by-event fluctuations in the number of participating protons). One can suppress this effect by selection of narrower centrality bins. In Fig. 13 the centrality of Xe+La at 13A GeV beam energy is restricted from 0–20% (rightmost points) down to 0–1% most central (leftmost points). As the same (0–20%) event sample was used points may be correlated. The negatively charged particles do not show any dependence on  $\sigma/\sigma_{\text{total}}$ . As expected, the  $\Phi_{p_T}$  values for all charged and positively charged particles decrease when going to more central collisions, reaching the values similar to that for negatively charged at approximately 7% most central interactions.




Fig. 13.  $\Phi_{p_T}$  and mean multiplicities as function of percent of total inelastic cross section for Xe+La collisions at the lowest SPS energy. Note: the values and their errors are correlated.

Figure 13 confirms that the centrality range (0–20%) of the events planned to be recorded by the NA61 experiment may need further restrictions in order to calculate transverse momentum fluctuations. Therefore, in Fig. 14

the same results as in Fig. 10 are presented but additional cut on selection of 7.2% most central collisions was applied<sup>1</sup>. This cut corresponds to centrality selection done by NA49 in the study of the energy dependence of  $\Phi_{p_T}$  for Pb+Pb interactions [14]. In Fig. 14, additional points for UrQMD1.3 Pb+Pb collisions are included. For 7.2% most central interactions  $\Phi_{p_T}$  measure is close to zero but only for negatively charged particles. For positively charged it is close to zero<sup>2</sup> but only for heavier systems (Pb+Pb and Xe+La), however for light systems (especially Be+Be) we still observe a significant increase at lower SPS energies. This increase is not visible any more when protons are removed from the sample. As the centrality is already restricted here, we can guess that there may be another yet source of correlations (conservation laws?) that produces positive  $\Phi_{p_T}$  values for central interactions at lower energies. This hypothesis needs further investigations, however, it is already supported by the recent observation [26] that also in UrQMD  $p+p$  interactions (no centrality restrictions required)  $\Phi_{p_T}$  values for all charged and for positively charged particles are increased at lower SPS energies, whereas the negatively charged ones are consistent with zero in the whole SPS energy range.




Fig. 14. Energy dependence of  $\Phi_{p_T}$  for different charge combinations of particles produced in UrQMD. 7.2% most central interactions are selected.

#### 4.4. $\Delta^{XN}$ -dependence on energy and charge combination

In Fig. 15 the energy dependence of  $\Delta^{XN}$  is shown for 20% most central  $A + A$  interactions. Three different charge combinations are included: all charged particles, negatively charged and positively charged. The open symbols in the right panel represent positively charged particles, where protons were removed from the sample. In contrary to  $\Phi_{p_T}$  measure (see Fig. 10), the  $\Delta^{XN}$  measure shows monotonic increase with increasing energy for all

<sup>1</sup> In UrQMD simulations centrality 0–7.2% corresponds to impact parameter cut  $b < 1.17$  fm for Be+Be,  $b < 2.29$  fm for Ar+Ca, and  $b < 3.42$  fm for Xe+La interactions.

<sup>2</sup> Very weak increase of  $\Phi_{p_T}$  with energy may be observed for Pb+Pb and Xe+La.

three charge combinations and for all studied systems. Similarly to  $\Phi_{pT}$ , the values of  $\Delta^{XN}$  for positively charged particles are comparable with those for negatively charged ones provided that protons are removed from the sample.




Fig. 15. Energy dependence of  $\Delta^{XN}$  for different charge combinations of particles produced in 20% most central  $A + A$  collisions in UrQMD.

In the next set of plots (Fig. 16), the  $\Delta^{XN}$  values are shown for forward rapidity only ( $1.1 < y_{\pi}^* < 2.6$ ). Additionally, only particles with  $y_p^* < y_{\text{beam}}^* - 0.5$  were accepted. As seen  $\Delta^{XN}$  values measured at forward rapidity are much smaller than those for complete rapidity region (see Fig. 15 for a comparison). The negatively charged particles show only a very weak increase and then saturation of  $\Delta^{XN}$  with increasing energy. Positively charged particles at forward rapidity show decrease of  $\Delta^{XN}$  values with energy for heavier systems (Ar+Ca, Xe+La). This is in contrary to  $\Phi_{pT}$  (see Fig. 11), which showed an increase with energy for positively charged particles in Xe+La interactions.




Fig. 16. Energy dependence of  $\Delta^{XN}$  at forward rapidity for different charge combinations of particles produced in 20% most central  $A + A$  collisions in UrQMD. Additional cut  $y_p^* < y_{\text{beam}}^* - 0.5$  was applied.

Similarly to what was done for  $\Phi_{pT}$ , the centrality was also restricted when calculating  $\Delta^{XN}$ . In Fig. 17 the centrality of Xe+La at 13A GeV beam energy is restricted from 0–20% (rightmost points) down to 0–1% most central (leftmost points). The negatively charged particles show a



small increase of  $\Delta^{XN}$  with increasing  $\sigma/\sigma_{\text{total}}$  ( $\Phi_{pT}$  values were constant — see Fig. 13). The  $\Delta^{XN}$  values for all charged particles and for positively charged ones decrease when going to more central collisions, thus confirming that a significant contribution to  $\Delta^{XN}$  may be indeed from event-by-event fluctuations of the number of participating protons.




Fig. 17.  $\Delta^{XN}$  as function of percent of total inelastic cross section for Xe+La collisions at the lowest SPS energy. Note: the values and their errors are correlated.

Figure 18 presents the energy dependence of  $\Delta^{XN}$  for 7.2% most central  $A + A$  collisions (additional points for UrQMD1.3 Pb+Pb collisions are included). In the case of Xe+La and Ar+Ca data, the 7.2% most central interactions result in  $\Delta^{XN}$  values much smaller than those for 20% most central (see Fig. 15 for a comparison). For Be+Be interactions such drop is much less pronounced and, in fact, the magnitudes for 7.2% and 20% most central interactions are comparable. For all systems the values of  $\Delta^{XN}$  for 7.2% most central collisions are higher for positively charged particles than for negatively charged ones and the rejection of protons can bring the magnitudes of positively charged ones close to these for nega-




Fig. 18. Energy dependence of  $\Delta^{XN}$  for different charge combinations of particles produced in UrQMD. 7.2% most central interactions are selected.

tively charged. In contrary to what was observed for  $\Phi_{p_T}$ , for very central collisions  $\Delta^{XN}$  for negatively charged particles show a rudimentary energy dependence. All systems and charge combinations show slight increase of  $\Delta^{XN}$  with energy and only central Pb+Pb simulations seem to exhibit a very weak decrease/plateau of  $\Delta^{XN}$  with energy. The different behaviour of Pb+Pb collisions may be connected with the fact that the earlier version UrQMD1.3 was used for Pb+Pb simulations, whereas UrQMD3.3 was employed to simulate Be+Be, Ar+Ca and Xe+La interactions.

In a simple superposition model all three fluctuation measures  $\Delta^{XN}$ ,  $\Sigma^{XN}$  and  $\Phi_{p_T}$  should have the same value, independent of the system size (the number of sources). Therefore, when cancelling the trivial effect of event-by-event impact parameter fluctuations (restricting to 7.2% most central interactions) we should expect the same values of fluctuation measures for all studied systems. When considering “most save” sample of negatively charged particles one can see that it is indeed true for  $\Phi_{p_T}$  measure (see middle panel of Fig. 14), whereas for  $\Delta^{XN}$  (middle panel of Fig. 18) it is true only for lower SPS energies. As the results from the fast generators in the previous sections showed that  $\Delta^{XN}$ ,  $\Sigma^{XN}$  and  $\Phi_{p_T}$  are indeed strongly intensive measures, it may suggest that the UrQMD model introduces somehow a small but systematic deviation from a simple superposition model beginning from middle SPS energies.  $\Phi_{p_T}$  measure can be less sensitive than  $\Delta^{XN}$  to such deviation and, therefore, its values are the same for all systems in the whole SPS energy range.

In the models the deviation from a simple superposition model (Wounded Nucleon Model) may be due to changes of the shape of the single particle spectrum and/or due to changes of fluctuations. It was suggested [27] that an important difference between  $\Phi$  and  $\Delta$  or  $\Sigma$  is that in case of  $\Phi$  measure the single particle term is subtracted (the second term of  $\Phi$  definition in Eq. (1)), thus making  $\Phi$  sensitive only to changes of fluctuations and insensitive to changes of the shape of the single particle spectrum. Such subtraction is not done for  $\Delta$  and  $\Sigma$ , thus in this respect, they are equivalent to the first (fluctuation) term of  $\Phi$ . As a consequence for an uncorrelated particle production,  $\Phi$  is zero and independent of a shape of the single particle spectrum. On the contrary, the  $\Delta$  and  $\Sigma$  are non-zero and their values depend on the shape of the single particle spectrum. If the UrQMD model introduces deviations from a simple superposition model by modifying the shape of the single particle spectrum only, then the large deviation from a superposition behaviour should be visible for  $\Delta$  and  $\Sigma$  only but not for  $\Phi$ . Therefore, in the analysis of experimental data a simultaneous measurement of all three quantities  $\Phi$ ,  $\Delta$ , and  $\Sigma$  may help to understand the origin of superposition model violation.

4.5.  $\Sigma^{XN}$ -dependence on energy and charge combination

Figure 19 shows the energy dependence of  $\Sigma^{XN}$  for 20% most central  $A + A$  interactions. Three different charge combinations are presented: all charged particles, negatively charged and positively charged. The open symbols in the right panel represent positively charged particles without protons. The samples of all charged and positively charged particles show a monotonic decrease of  $\Sigma^{XN}$  with increasing energy, whereas the negatively charged rise and reach saturation at middle SPS energies. Similarly to  $\Phi_{pT}$  and  $\Delta^{XN}$  the values of  $\Sigma^{XN}$  for positively charged particles became comparable with those for negatively charged ones provided that protons are removed from the sample.




Fig. 19. Energy dependence of  $\Sigma^{XN}$  for different charge combinations of particles produced in 20% most central  $A + A$  collisions in UrQMD.

In Fig. 20 the values of  $\Sigma^{XN}$  are shown for forward rapidity only ( $1.1 < y_{\pi}^* < 2.6$ ). Additionally, only particles with  $y_p^* < y_{\text{beam}}^* - 0.5$  were accepted.  $\Sigma^{XN}$  values measured at forward rapidity are slightly smaller than those for complete rapidity region. All three charge combinations and all systems show an increase and then saturation of  $\Sigma^{XN}$  with increasing energy.




Fig. 20. Energy dependence of  $\Sigma^{XN}$  at forward rapidity for different charge combinations of particles produced in 20% most central  $A + A$  collisions in UrQMD. Additional cut  $y_p^* < y_{\text{beam}}^* - 0.5$  was applied.

Figure 21 presents the values of  $\Sigma^{XN}$  when the centrality of Xe+La at 13A GeV beam energy is restricted from 0–20% (rightmost points) down to 0–1% most central (leftmost points). The negatively charged particles exhibit nearly flat dependence of  $\Sigma^{XN}$  on  $\sigma/\sigma_{\text{total}}$  (similarly to  $\Phi_{pT}$  — see Fig. 13). The values of  $\Sigma^{XN}$  for all charged particles and for positively charged ones decrease when going to more central collisions and reach a plateau for approximately 10% most central collisions. There is, however, one significant difference between the centrality dependence of  $\Phi_{pT}$  or  $\Delta^{XN}$  and  $\Sigma^{XN}$  for Xe+La at 13A GeV. For extremely central collisions, the values of both  $\Phi_{pT}$  and  $\Delta^{XN}$  are similar for positively, negatively, and all charged particles (see Figs. 13 and 17). On the contrary, the values of  $\Sigma^{XN}$  for very central collisions are much different for negatively charged particles than for the positively charged and all charged. It might indicate the presence of additional source of correlations in the sample of positively charged particles, which was not detected by use of  $\Phi_{pT}$  and  $\Delta^{XN}$  measures. The origin of such correlation(s) was not investigated here.




Fig. 21.  $\Sigma^{XN}$  as function of percent of total inelastic cross section for Xe+La collisions at the lowest SPS energy. Note: the values and their errors are correlated.

The last set of plots (Fig. 22) presents the energy dependence of  $\Sigma^{XN}$  for 7.2% most central  $A + A$  collisions. For all systems the values of  $\Sigma^{XN}$  are higher for positively charged particles than for negatively charged ones and, as usually, the rejection of protons brings the magnitudes of positively charged ones close to these for negatively charged.

The 7.2% most central collisions showed a difference between the values of fluctuation measures for positively charged particles and for positively charged without protons. In the case of  $\Phi_{pT}$  (Fig. 14) this difference is significant only for light systems (Be+Be, Ar+Ca) and in the case of  $\Delta^{XN}$  and  $\Sigma^{XN}$  such deviation is observed for all systems. Therefore, the results suggest that either the centrality bin 0–7.2% is still not narrow enough

(it may be the case for  $\Delta^{XN}$ , see Fig. 17) and event-by-event impact parameter (number of protons) fluctuations can be still present, or there is another yet source of correlations visible for positively charge particles only (then protons, of course, would have a significant contribution to such correlation). The  $\Sigma^{XN}$  measure can be much more sensitive to such kind of correlation (conservation laws?) because a difference between  $\Sigma^{XN}$  for positively charged particles and for positively charged without protons is visible for all systems. In the case of  $\Phi_{pT}$ , as already mentioned, only smaller systems show this deviation. The origin of such possible correlation is still under study and, in particular, the  $\Phi_{pT}$  and  $\Sigma^{XN}$  measures will be carefully analysed for the energy scan of UrQMD  $p + p$  collisions [26].

Finally, in Fig. 22, one observes that for 7.2% most central collisions the  $\Sigma^{XN}$  values for negatively charged particles are nearly independent of energy. There is, however, rather significant system size ( $A$ ) dependence suggesting the deviation of UrQMD from a simple superposition model. Such deviation was also observed for  $\Delta^{XN}$  measure (middle panel of Fig. 18), however only at higher SPS energies.




Fig. 22. Energy dependence of  $\Sigma^{XN}$  for different charge combinations of particles produced in UrQMD. 7.2% most central interactions are selected.

## 5. Summary and conclusions

In this publication, the recently proposed  $\Delta^{XN}$  and  $\Sigma^{XN}$  measures of transverse momentum fluctuations are tested and compared to  $\Phi_{pT}$  measure. The fast generator simulations confirm that all three measures are both intensive and strongly intensive. The results suggest that all three measures are also very sensitive to the effect of event-by-event “temperature” fluctuations. Such effect should be better visible for heavier systems.

Some preliminary tests were performed also within the UrQMD model, which showed many interesting effects, different for different measures. The complete understanding of all possible sources of fluctuations and correlations in UrQMD would require much more detailed investigations and it is

out of the scope of this publication. However, a rather consistent picture can be drawn from these basic tests: if one wants to study the exotic effects such as critical point, onset of deconfinement, *etc.* one should find good reference values of the studied measures. The UrQMD model suggests that the best reference values are those for negatively charged particles produced in very central  $A + A$  collisions, however, even for these samples, one should first try to better understand the origin of the rudimentary energy and/or system size dependence of  $\Delta^{XN}$  and  $\Sigma^{XN}$  observed in UrQMD. The simultaneous measurement of all three quantities suggests that the system size ( $A$ ) dependence of  $\Delta^{XN}$  and  $\Sigma^{XN}$  can be due to change of the single particle spectrum mainly. The  $\Phi_{pT}$  values for negatively charged particles and for very central  $A + A$  are consistent with zero and do not depend on the mass of the colliding nuclei. For a sample of positively charged particles, the reference values of all three measures may be masked by some trivial sources of fluctuations and correlations caused by the presence of protons in the sample. The fluctuation measures for a sample of all charged particles is always a non trivial combination of measures for positively and negatively charged particles. Moreover, correlations between positively and negatively charged particles (*i.e.* due to particle decays) contribute here.

Finally, one should mention that in most cases the restriction of the analysis to forward rapidity only reduces the values of fluctuation measures. Therefore, the NA61/SHINE experiment should do its best in order to measure event-by-event fluctuations in a rapidity window as wide as possible.

I am indebted to the authors of the UrQMD model for the permission to use their code in my analysis. I also would like to thank Bartosz Maksiak for a possibility of using his UrQMD3.3 Be+Be, Ar+Ca and Xe+La data sets. I am very grateful to Marek Gaździcki and Zbigniew Włodarczyk for their useful comments concerning this analysis, and to Maja Maćkowiak-Pawłowska for careful reading of the manuscript.

## REFERENCES

- [1] M. Gaździcki, M.I. Gorenstein, *Acta Phys. Pol. B* **30**, 2705 (1999).
- [2] S.V. Afanasiev *et al.* [NA49 Collaboration], *Phys. Rev.* **C66**, 054902 (2002);  
C. Alt *et al.* [NA49 Collaboration], *Phys. Rev.* **C77**, 024903 (2008).
- [3] Z. Fodor, S.D. Katz, *J. High Energy Phys.* **0404**, 050 (2004).
- [4] A. Li *et al.*, RIKEN-BNL Workshop on Fluctuations, Correlations and RHIC Low Energy Runs, Oct. 3–5, 2011, <http://www.bnl.gov/fcrworkshop/>
- [5] M. Stephanov, K. Rajagopal, E.V. Shuryak, *Phys. Rev.* **D60**, 114028 (1999) and private communication.

- [6] F. Beccatini, J. Manninen, M. Gaździcki, *Phys. Rev.* **C73**, 044905 (2006).
- [7] S. Afanasiev *et al.* [NA49 Collaboration], *Nucl. Instrum. Methods* **A430**, 201 (1999).
- [8] K. Grebieszko [NA49 Collaboration], *Nucl. Phys.* **A830**, 547C (2009) and references therein.
- [9] <https://na61.web.cern.ch/na61/xc/index.html>
- [10] K. Grebieszko [NA61 Collaboration], [arXiv:1201.0742v1](https://arxiv.org/abs/1201.0742v1) [nucl-ex].
- [11] M. Gaździcki, S. Mrówczyński, *Z. Phys.* **C54**, 127 (1992).
- [12] A. Białas, M. Bleszyński, W. Czyż, *Nucl. Phys.* **B111**, 461 (1976).
- [13] T. Anticic *et al.* [NA49 Collaboration], *Phys. Rev.* **C70**, 034902 (2004).
- [14] T. Anticic *et al.* [NA49 Collaboration], *Phys. Rev.* **C79**, 044904 (2009).
- [15] M.I. Gorenstein, M. Gaździcki, *Phys. Rev.* **C84**, 014904 (2011).
- [16] S.A. Bass *et al.*, *Prog. Part. Nucl. Phys.* **41**, 225 (1998); M. Bleicher *et al.*, *J. Phys. G* **25**, 1859 (1999).
- [17] G. Wilk, Z. Włodarczyk, *Phys. Rev.* **C79**, 054903 (2009).
- [18] G. Wilk, Z. Włodarczyk, *J. Phys. G* **38**, 065101 (2011).
- [19] G. Wilk, Z. Włodarczyk, [arXiv:1203.4452v1](https://arxiv.org/abs/1203.4452v1) [hep-ph].
- [20] R. Korus *et al.*, *Phys. Rev.* **C64**, 054908 (2001).
- [21] M. Bleicher *et al.*, *Phys. Lett.* **B435**, 9 (1998).
- [22] K. Grebieszko, *Phys. Rev.* **C76**, 064908 (2007).
- [23] M. Bleicher, S. Jeon, V. Koch, *Phys. Rev.* **C62**, 061902 (2000).
- [24] V.P. Konchakovski *et al.*, *Phys. Rev.* **C73**, 034902 (2006).
- [25] C. Alt *et al.* [NA49 Collaboration], *Phys. Rev.* **C75**, 064904 (2007).
- [26] T. Czopowicz, private communication.
- [27] M. Gaździcki, private communication.