LIGHT CLUSTER PRODUCTION AT NICA*

N.-U. BASTIAN^a, D. BLASCHKE^{a,b,c}, G. RÖPKE^{c,d}

^aInstitute of Theoretical Physics, University of Wrocław, Wrocław, Poland ^bBogoliubov Laboratory of Theoretical Physics, JINR, Dubna, Russia ^cNational Research Nuclear University (MEPhI), Moscow, Russia ^dInstitute of Physics, University of Rostock, Rostock, Germany

(Received February 16, 2017)

Measuring abundances and flow of light clusters as "rare probes" in future heavy-ion collision experiments at NICA and FAIR energies would provide unique insights into the QCD phase diagram.

DOI:10.5506/APhysPolBSupp.10.899

In a recent publication [1], we discussed the insights that light nuclear clusters can provide for the QCD phase diagram shown in Fig. 1, in particular within the NICA program, providing fixed target (BM@N) and collider (MPD) experiments. This diagram includes lines for Mott dissociation [2] of deuterons (d), tritons (t) and alpha particles (α), taken from [3], together with the parametrization of the chemical freeze-out line [4]. The chiral and deconfinement lines are from [5] based on a local NJL model and the DD2 equation of state [6]. The nonlocal PNJL extension [7] has a critical endpoint at a location supporting the conjecture of a tricritical point due to a quarkyonic matter phase [8]. The projection of its location to the freezeout line at $E_{\text{lab}} = 15-25 \text{ A MeV} (\sqrt{s_{NN}} = 5.6-7.1 \text{ GeV})$ coincides with a dip in the slope of the directed flow of protons and light clusters [1]. At higher energies, the expectations from the law of mass action [9] are confirmed well by the ALICE experiment [10] whereas at lower energies, the expected strong modifications due to the Mott effect for light clusters [1-3] shall be observed, e.g., in the NICA experiment.

^{*} Presented at the "Critical Point and Onset of Deconfinement" Conference, Wrocław, Poland, May 30–June 4, 2016.



Fig. 1. QCD phase diagram with chemical freezeout [4] and Mott lines for the dissociation of light nuclear clusters [2,3]. For details, see [1,5].

N.-U.B. and D.B. were supported by the National Science Centre, Poland (NCN) under contract No. UMO-2011/02/A/ST2/00306. D.B. and G.R. acknowledge support by the MEPhI Academic Excellence Project under contract No. 02.a03.21.0005.

REFERENCES

- [1] N.-U. Bastian et al., Eur. Phys. J. A 52, 244 (2016).
- [2] G. Röpke et al., Nucl. Phys. A **399**, 587 (1983).
- [3] P. Schuck et al., Few Body Syst. Suppl. 12, 50 (2000).
- [4] J. Cleymans et al., Phys. Rev. C 73, 034905 (2006).
- [5] N.U. Bastian, D. Blaschke, J. Phys.: Conf. Ser. 668, 012042 (2016).
- [6] S. Typel et al., Phys. Rev. C 81, 015803 (2010).
- [7] G.A. Contrera *et al.*, *Eur. Phys. J. A* **52**, 231 (2016).
- [8] A. Andronic et al., Nucl. Phys. A 837, 65 (2010).
- [9] S. Das Gupta *et al.*, *Phys. Rep.* **72**, 131 (1981).
- [10] J. Adam et al. [ALICE Collaboration], Phys. Rev. C 93, 024917 (2016).