### No 3

# ROTATIONAL BANDS IN SUPER-HEAVY NUCLEI WITHIN THE LSD+YF MODEL\*

# BOŻENA NERLO-POMORSKA, KRZYSZTOF POMORSKI

Institute of Physics, Theoretical Physics Department Maria Curie-Skłodowska University, Lublin, Poland

# J. BARTEL

## Institute Pluridisciplinaire Hubert Curien, Université de Strasbourg Strasbourg, France

(Received December 21, 2018)

Rotational energies of heavy and super-heavy nuclei are evaluated in the cranking model which couples the pairing field with the rotational motion. The nuclear ground-state deformation is determined within the macroscopic–microscopic model.

DOI:10.5506/APhysPolBSupp.12.665

# 1. Introduction

An investigation of heavy and super-heavy nuclei requires a proper model to reproduce masses and rotational energies. We had obtained in the past [1] a very good agreement with experimental data for heavy nuclei using the Yukawa folded (YF) single-particle potential [2] and the Lublin–Strasbourg Drop (LSD) [3]. The Strutinsky shell-correction method [4] and the BCS theory [5] were used to evaluate the shell and pairing energy. The equilibrium deformations of the nuclei were determined with the Modified Funny Hills (MFH) [1] shape parametrisation, and the energies of the rotational states obtained using the cranking model [6]. It was shown in Ref. [7] that a good agreement with the experimental data, especially for larger angular momenta, can only be achieved when a coupling of rotation and the pairing mode is taken into account. We are now going to extend our calculations to rotational states of super-heavy nuclei using a new Fourier-type shape

<sup>\*</sup> Presented at the XXV Nuclear Physics Workshop "Structure and Dynamics of Atomic Nuclei", Kazimierz Dolny, Poland, September 25–30, 2018.

parametrization [8] to describe nuclear deformations more accurately. First results of this project as obtained in a 3-dimensional deformation space are presented below.

# 2. Theoretical model

In the macroscopic–microscopic method [9], the total energy of a nucleus at a given deformation (def) is the sum of the macroscopic energy and microscopic corrections due to shell and pairing effects

$$E_{\text{tot}}(\text{def}) = E_{\text{LSD}}(\text{def}) + E_{\text{shell}}(\text{def}) + E_{\text{pair}}(\text{def}).$$
(1)

The shell-correction energy is obtained as the difference between the singleparticle energy sum and the corresponding Strutinsky averaged result [4], while pairing corrections are determined as the difference between the BCS [5] energy and the single-particle level sum, plus an average pairing energy [10]. Energy landscapes were calculated in Ref. [1] on a two-dimensional grid of Modified Funny Hills [11] shapes with the Lublin–Strasbourg Drop and using Yukawa-folded single-particle levels for the quantal corrections. Rotational states were determined within the cranking model in the ground-state minimum and reproduced the experimental data or served as predictions in the case of heavy and super-heavy nuclei as shown in Fig. 1. We are now going to investigate these and some new even–even super-heavy nuclei using the new Fourier-shape parametrisation [8] in a 3D space including elongation  $(q_2)$ , octupole  $(q_3)$  deformation and neck degrees of freedom  $(q_4)$ .



Fig. 1. Nuclei for which the energies of rotational states have been determined in the MFH (× and  $\star$ ) and Fourier ( $\Box$ ) shape parametrizations. The  $\beta$ -stability line is drawn as a guideline.

667

## 3. Results

The calculations were performed for even–even nuclei with charge numbers from Z = 94 to Z = 112, since nuclei with larger Z are probably not living long enough to perform spectroscopy studies [12]. In Fig. 2, we present the difference of theoretical and experimental [13] masses of nuclei between No and Hs. With the single exception of <sup>264</sup>Hs, this deviation never exceeds 0.3 MeV. As an example, the deformation energy sur-



Fig. 2. Difference between theoretical and experimental masses  $M_{\rm th} - M_{\rm exp}$  for No up to Hs even–even isotopes as a function of the neutron number N (see Ref. [1]).

face  $E_{\text{def}} = E_{\text{tot}}(q_2, q_3, q_4^{\text{min}}) - E_{\text{LSD}}^{\text{sph}}(0, 0, 0)$  of <sup>254</sup>No is presented in Fig. 3 as a function of the deformation parameters  $q_2$  (elongation) and  $q_3$  (left-right asymmetry), with a minimisation with respect to  $q_4$  (neck parameter). One observes two pronounced minima corresponding to the ground



Fig. 3. Deformation energy  $E_{\text{def}}$  for <sup>254</sup>No as a function of elongation  $q_2$  and octupole parameter  $q_3$  minimised with respect to  $q_4$  (neck degree of freedom).

and the fission-isomeric state. In Fig. 4, the deformation energy surface  $E_{\text{def}} = E_{\text{tot}}(0.3, q_3, q_4) - E_{\text{LSD}}^{\text{sph}}(0, 0, 0)$  of this nucleus <sup>254</sup>No is displayed for the ground state ( $q_2 = 0.3$ ), which turns out to be octupole and hexade-capole symmetric.



Fig. 4. Deformation energy  $E_{\text{def}}$  for <sup>254</sup>No as a function of  $q_4$  and  $q_3$  in a ground-state elongation  $q_2 = 0.3$ .

To obtain the rotational energies,  $E_L = L(L+1)/2\mathcal{J}$ , the moments of inertia  $\mathcal{J}$  are calculated microscopically within the cranking model [6] for the ground state using a Yukawa-folded single-particle potential. Rotational energies of No isotopes for  $L/\hbar = 2, 4, 6, 8, 10, 12$  are compared with the experimental data [14] in Fig. 5. As seen from the figure, one obtains an almost perfect agreement when a pairing strengths of  $G N_q^{2/3} = 0.32 \hbar \omega_0$  with  $N_q = N, Z$  and  $\hbar \omega_0 = 41 \,\mathrm{MeV}/A^{1/3}$  when the pairing window of  $2\sqrt{15N_q}$ single-particle levels [15] is used. Figure 6 gives a prediction of the  $E_{2+}$  ro-



Fig. 5. Theoretical and experimental rotational energies E for  $^{254}$ No as functions of the angular momentum L.

tational energies for even–even nuclei between No and Cn. In a forthcoming extension of the present work, we are going to predict the rotational energies of heavy and super-heavy nuclei with  $92 \le Z \le 126$  and  $132 \le N \le 200$  within LSD+YF model using the Fourier shape parametrisation [8, 18] in a 4D deformation space.



Fig. 6. Rotational energies  $E_{2+}$  for even-even isotopes of nuclei between No and Cn in their ground state as functions of the neutron number N [16, 17].

This work has been partly supported by the Polish–French COPIN-IN2P3 collaboration agreement under project number 08-131 and by the National Science Centre, Poland (NCN), grant No. 2016/21/B/ST2/01227.

#### REFERENCES

- A. Dobrowolski, B. Nerlo-Pomorska, K. Pomorski, Acta Phys. Pol. B 42, 105 (2011).
- [2] K.T.R. Davies, J.R. Nix, *Phys. Rev. C* 14, 1977 (1976).
- [3] K. Pomorski, J. Dudek, *Phys. Rev. C* 67, 044316 (2003).
- [4] V.M. Strutinsky, Nucl. Phys. A 95, 420 (1967).
- [5] J. Bardeen, L.N. Cooper, J.R. Schrieffer, *Phys. Rev.* 108, 1175 (1957).
- [6] S.G. Nilsson, O. Prior, Mat. Fys. Medd. Dan. Vid. Selsk 32, 16 (1961).
- [7] B. Nerlo-Pomorska, K. Pomorski, J. Bartel, *Phys. Rev. C* 84, 044310 (2011).
- [8] C. Schmitt, K. Pomorski, B. Nerlo-Pomorska, J. Bartel, *Phys. Rev. C* 95, 034612 (2017).

- [9] W.D. Myers, W.J. Świątecki, Nucl. Phys. 81, 1 (1966); Ark. Phys. 36, 343 (1967).
- [10] B. Nerlo-Pomorska, K. Pomorski, Int. J. Mod. Phys. E 16, 328 (2007).
- [11] K. Pomorski, J. Bartel, Int. J. Mod. Phys. E 15, 417 (2006).
- [12] A. Sobiczewski, K. Pomorski, Prog. Part. Nucl. Phys. 58, 292 (2007).
- [13] M.S. Antony, Nuclide Chart, Strasbourg 2002, Impressions Francois, Hagenau 2002.
- [14] http://www.nndc.bnl.gov/nudat2
- [15] S.G. Nilsson et al., Nucl. Phys. A 131, 1 (1969).
- [16] P.T. Greenlees et al., Phys. Rev. Lett. 109, 012501 (2012).
- [17] B. Nerlo-Pomorska, K. Pomorski, J. Bartel, *Phys. Rev. C* 81, 044310 (2011).
- [18] B. Nerlo-Pomorska, K. Pomorski, J. Bartel, C. Schmitt, *Acta Phys. Pol. B* 48, 451 (2017).