

THEORETICAL ASPECTS OF RELATIVISTIC PERFECT-FLUID SPIN HYDRODYNAMIC FRAMEWORK*

RAJEEV SINGH

Institute of Nuclear Physics Polish Academy of Sciences, 31-342 Kraków, Poland

*Received 15 April 2022, accepted 29 April 2022,
published online 9 September 2022*

We review the recently proposed perfect-fluid spin hydrodynamic formalism, which provides a new tool for the description of the spin polarization of $\Lambda(\bar{\Lambda})$ particles. This formalism is based on the de Groot–van Leeuwen–van Weert definitions of the energy-momentum and spin tensors.

DOI:10.5506/APhysPolBSupp.15.3-A35

1. Introduction

Relativistic hydrodynamics has been quite successful in describing phenomena in various areas of physics [1–4], consequently making the path in explaining the interesting properties of quark–gluon plasma [5]. Spin polarization measurements in relativistic heavy-ion collisions [6–10] provided a new area for study [11]. These measurements initiated vast theoretical developments concerning spin-vorticity coupling [12–40]. However, after explaining global spin polarization, spin-thermal-based frameworks have not been able to explain longitudinal spin polarization [7, 20, 41]. This discrepancy instigated a new formalism of spin hydrodynamics, first proposed in Ref. [42], see also Refs. [30, 43–70] for further developments and new approaches. The evolution of spin polarization in spin hydrodynamics [42–44] is controlled by an antisymmetric spin polarization tensor $\omega^{\alpha\beta}$ (independent of thermal vorticity), introducing six spin polarization components which need to be determined along with the background parameters from the conservation laws. The framework of spin hydrodynamics [42, 43] uses the de Groot–van Leeuwen–van Weert (GLW) [71] definitions of energy-momentum and spin tensors. Keeping in mind the smallness of spin polarization magnitude in the experiments, the spin polarization tensor [42, 43] is assumed to be of leading order, hence having contributions only in the spin tensor,

* Presented at the 28th Cracow Epiphany Conference on *Recent Advances in Astroparticle Physics*, Cracow, Poland, 10–14 January, 2022.

making the spin dynamics decouple from the perfect-fluid background dynamics [44, 45, 63]. After obtaining the dynamics of spin, we then move on to calculate the mean spin polarization per particle in the particle rest frame (PRF) to be compared with the experimental data [63].

2. Spin hydrodynamic equations

The formalism of the perfect-fluid spin hydrodynamics for particles with spin-1/2 is based on the conservation of charge, energy-momentum, and angular momentum with the GLW [71] definitions of the energy-momentum tensor, $T_{\text{GLW}}^{\alpha\beta}$, and spin tensor, $S_{\text{GLW}}^{\alpha\beta\gamma}$ ¹, such as [42, 43, 45]

$$\partial_\mu N^\mu = 0, \quad \partial_\mu T_{\text{GLW}}^{\mu\nu} = 0, \quad \partial_\lambda S_{\text{GLW}}^{\lambda,\alpha\beta} = T_{\text{GLW}}^{\beta\alpha} - T_{\text{GLW}}^{\alpha\beta}, \quad (1)$$

where

$$N^\alpha = nU^\alpha, \quad T_{\text{GLW}}^{\alpha\beta} = (\varepsilon + P)U^\alpha U^\beta - Pg^{\alpha\beta}, \quad (2)$$

with N^α being the net baryon charge current while ε , P , and n are the energy density, pressure, and baryon density, respectively. U^β represents the four-vector fluid flow which is time-like, hence $U \cdot U = 1$.

The symmetric nature of GLW energy-momentum tensor (1) implies separate spin conservation [44, 45] with the spin tensor expressed by $S_{\text{GLW}}^{\alpha,\beta\gamma} = \cosh(\xi) \left(n_{(0)}(T) U^\alpha \omega^{\beta\gamma} + S_{\Delta\text{GLW}}^{\alpha,\beta\gamma} \right)$, where [43, 45]

$$S_{\Delta\text{GLW}}^{\alpha,\beta\gamma} = \mathcal{A}_{(0)} U^\alpha U^\delta U^{[\beta} \omega^{\gamma]}_\delta + \mathcal{B}_{(0)} \left(U^{[\beta} \Delta^{\alpha\delta} \omega^{\gamma]}_\delta + U^\alpha \Delta^{\delta[\beta} \omega^{\gamma]}_\delta + U^\delta \Delta^{\alpha[\beta} \omega^{\gamma]}_\delta \right),$$

and the thermodynamic coefficients are $\mathcal{B}_{(0)} = -\frac{2}{\hat{m}^2} s_{(0)}(T)$ and $\mathcal{A}_{(0)} = -3\mathcal{B}_{(0)} + 2n_{(0)}(T)$, with $n_{(0)}(T)$ and $s_{(0)}(T)$ being the number density and entropy density for the case of neutral and massive spin-less Boltzmann particles, $\Delta^{\alpha\beta}$ is the spatial projector operator transverse to U , ξ denotes the ratio between the baryon chemical potential (μ) and temperature (T) whereas $\hat{m}$ denotes the ratio between the mass of the particle and temperature [45, 63]. The quantity $\omega_{\mu\nu}$ is called the spin polarization tensor $\omega_{\mu\nu}$, antisymmetric and rank-two, that can be written in terms of κ^μ and ω^μ four-vectors [42, 63]

$$\omega_{\mu\nu} = \kappa_\mu U_\nu - \kappa_\nu U_\mu + \epsilon_{\mu\nu\alpha\beta} U^\alpha \omega^\beta. \quad (3)$$

κ^μ and ω^μ being parallel to fluid flow have no contribution to the R.H.S. of Eq. (3), thus they fulfill²

$$\kappa \cdot U = 0, \quad \omega \cdot U = 0 \quad (4)$$

¹ Spin polarization is assumed to be small ($|\omega_{\mu\nu}| < 1$) in this formalism.

² κ^μ and ω^μ together form six independent components of $\omega_{\mu\nu}$ [63].

allowing us to write these four-vectors as [44, 45, 63] $\kappa_\mu = \omega_{\mu\alpha} U^\alpha$, $\omega_\mu = \frac{1}{2}\epsilon_{\mu\alpha\beta\gamma}\omega^{\alpha\beta}U^\gamma$, and ω^μ can be decomposed again in terms of the basis vectors X, Y, Z ³ and scalar spin coefficients ($C_{\kappa X}, C_{\kappa Y}, C_{\kappa Z}, C_{\omega X}, C_{\omega Y}, C_{\omega Z}$) after using orthogonality conditions (4) as [63]

$$\kappa^\alpha = C_{\kappa X}X^\alpha + C_{\kappa Y}Y^\alpha + C_{\kappa Z}Z^\alpha, \quad \omega^\alpha = C_{\omega X}X^\alpha + C_{\omega Y}Y^\alpha + C_{\omega Z}Z^\alpha, \quad (5)$$

which then help us to obtain the general form of $\omega_{\alpha\beta}$ [63]

$$\begin{aligned} \omega_{\alpha\beta} = & 2(C_{\kappa X}X_{[\alpha}U_{\beta]} + C_{\kappa Y}Y_{[\alpha}U_{\beta]} + C_{\kappa Z}Z_{[\alpha}U_{\beta]}) \\ & + \epsilon_{\alpha\beta\gamma\delta}U^\gamma(C_{\omega X}X^\delta + C_{\omega Y}Y^\delta + C_{\omega Z}Z^\delta). \end{aligned} \quad (6)$$

3. Spin polarization of particles at the freeze-out

With the information about the evolution of the spin polarization components and background parameters, we can now calculate the mean spin polarization per particle defined as $\langle\pi_\mu\rangle_p = E_p \frac{d\Pi_\mu^*(p)}{d^3p} / E_p \frac{d\mathcal{N}(p)}{d^3p}$, where [44, 63]

$$E_p \frac{d\Pi_\mu^*(p)}{d^3p} = -\frac{1}{(2\pi)^3m} \int \cosh(\xi) \Delta\Sigma_\lambda p^\lambda e^{-\beta \cdot p} \left(\tilde{\omega}_{\mu\beta} p^\beta\right)^*, \quad (7)$$

$$E_p \frac{d\mathcal{N}(p)}{d^3p} = \frac{4}{(2\pi)^3} \int \Delta\Sigma_\lambda p^\lambda \cosh(\xi) e^{-\beta \cdot p}, \quad (8)$$

and $E_p \frac{d\Pi_\mu^*(p)}{d^3p}$ is the total Pauli–Lubański (PL) vector in the PRF (after integrating over the freeze-out hyper-surface element $\Delta\Sigma_\lambda$) for particles having momentum p , while $E_p \frac{d\mathcal{N}(p)}{d^3p}$ represents the momentum density of all particles [45, 63]. The quantity $\langle\pi_\mu\rangle_p$, the average spin polarization per particle, is a function of momentum coordinates p_x, p_y, p_z [63] which will allow us to obtain the local spin polarization (polarization along the beam direction), whereas to calculate the global spin polarization (along the $-y$ axis) we must integrate over the momentum variable such as [63]

$$\langle\pi_\mu\rangle = \frac{\int dP \langle\pi_\mu\rangle_p E_p \frac{d\mathcal{N}(p)}{d^3p}}{\int dP E_p \frac{d\mathcal{N}(p)}{d^3p}} \equiv \frac{\int d^3p \frac{d\Pi_\mu^*(p)}{d^3p}}{\int d^3p \frac{d\mathcal{N}(p)}{d^3p}}. \quad (9)$$

³ X, Y, Z are the space-like four-vectors which span the space transverse to U [63].

4. Summary

In this article, we have reviewed the theoretical aspects of the spin hydrodynamic formalism [42–44]. Considering the spin polarization tensor of the leading order, we find the decoupling nature between the background and the spin dynamics. We then put forward the expressions required to calculate the mean spin polarization of the particles for the comparison with the experimental data [63].

I am extremely grateful to W. Florkowski, R. Ryblewski, and G. Sophys for fruitful collaboration. Supported in part by the National Science Center, Poland (NCN) grants No. 2016/23/B/ST2/00717 and No. 2018/30/E/ST2/00432.

REFERENCES

- [1] N. Andersson, G.L. Comer, «Relativistic Fluid Dynamics: Physics for Many Different Scales», *Living Rev. Rel.* **10**, 1 (2007), [arXiv:gr-qc/0605010](#).
- [2] W. Florkowski, «Phenomenology of Ultra-relativistic Heavy-ion Collisions», *World Scientific*, Singapore 2010, <https://cds.cern.ch/record/1321594>
- [3] C. Gale, S. Jeon, B. Schenke, «Hydrodynamic Modeling of Heavy-Ion Collisions», *Int. J. Mod. Phys. A* **28**, 1340011 (2013), [arXiv:1301.5893 \[nucl-th\]](#).
- [4] S. Jeon, U. Heinz, «Introduction to hydrodynamics», *Int. J. Mod. Phys. E* **24**, 1530010 (2015) [arXiv:1503.03931 \[hep-ph\]](#).
- [5] W. Florkowski, M.P. Heller, M. Spaliński, «New theories of relativistic hydrodynamics in the LHC era», *Rep. Prog. Phys.* **81**, 046001 (2018), [arXiv:1707.02282 \[hep-ph\]](#).
- [6] STAR Collaboration (L. Adamczyk *et al.*), «Global Λ hyperon polarization in nuclear collisions: evidence for the most vortical fluid», *Nature* **548**, 62 (2017), [arXiv:1701.06657 \[nucl-ex\]](#).
- [7] STAR Collaboration (J. Adam *et al.*), «Polarization of Λ ($\bar{\Lambda}$) Hyperons along the Beam Direction in Au+Au Collisions at $\sqrt{s_{NN}} = 200$ GeV», *Phys. Rev. Lett.* **123**, 132301 (2019), [arXiv:1905.11917 \[nucl-ex\]](#).
- [8] ALICE Collaboration (S. Acharya *et al.*), «Evidence of Spin-Orbital Angular Momentum Interactions in Relativistic Heavy-Ion Collisions», *Phys. Rev. Lett.* **125**, 012301 (2020), [arXiv:1910.14408 \[nucl-ex\]](#).
- [9] ALICE Collaboration (S. Acharya *et al.*), «Global polarization of $\Lambda\bar{\Lambda}$ hyperons in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ and 5.02 TeV», *Phys. Rev. C* **101**, 044611 (2020), [arXiv:1909.01281 \[nucl-ex\]](#).
- [10] STAR Collaboration (M.S. Abdallah *et al.*), «Global Λ -hyperon polarization in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV», *Phys. Rev. C* **104**, L061901 (2021), [arXiv:2108.00044 \[nucl-ex\]](#).

- [11] F. Becattini, J. Liao, M. Lisa, «Strongly Interacting Matter Under Rotation: An Introduction», *Lect. Notes Phys.* **987**, 1 (2021), [arXiv:2102.00933 \[nucl-th\]](#).
- [12] F. Becattini, F. Piccinini, J. Rizzo, «Angular momentum conservation in heavy ion collisions at very high energy», *Phys. Rev. C* **77**, 024906 (2008), [arXiv:0711.1253 \[nucl-th\]](#).
- [13] L.-G. Pang, H. Petersen, Q. Wang, X.-N. Wang, «Vortical Fluid and Λ Spin Correlations in High-Energy Heavy-Ion Collisions», *Phys. Rev. Lett.* **117**, 192301 (2016), [arXiv:1605.04024 \[hep-ph\]](#).
- [14] Y. Xie, D. Wang, L.P. Csernai, «Global Λ polarization in high energy collisions», *Phys. Rev. C* **95**, 031901 (2017), [arXiv:1703.03770 \[nucl-th\]](#).
- [15] F. Becattini, V. Chandra, L. Del Zanna, E. Grossi, «Relativistic distribution function for particles with spin at local thermodynamical equilibrium», *Ann. Phys.* **338**, 32 (2013), [arXiv:1303.3431 \[nucl-th\]](#).
- [16] F. Becattini *et al.*, «Global hyperon polarization at local thermodynamic equilibrium with vorticity, magnetic field and feed-down», *Phys. Rev. C* **95**, 054902 (2017), [arXiv:1610.02506 \[nucl-th\]](#).
- [17] Iu. Karpenko, F. Becattini, «Study of Λ polarization in relativistic nuclear collisions at $\sqrt{s_{NN}} = 7.7 - 200$ GeV», *Eur. Phys. J. C* **77**, 213 (2017), [arXiv:1610.04717 \[nucl-th\]](#).
- [18] Y. Sun, C.M. Ko, « Λ hyperon polarization in relativistic heavy ion collisions from a chiral kinetic approach», *Phys. Rev. C* **96**, 024906 (2017), [arXiv:1706.09467 \[nucl-th\]](#).
- [19] H. Li, L.-G. Pang, Q. Wang, X.-L. Xia, «Global Λ polarization in heavy-ion collisions from a transport model», *Phys. Rev. C* **96**, 054908 (2017), [arXiv:1704.01507 \[nucl-th\]](#).
- [20] F. Becattini, Iu. Karpenko, «Collective Longitudinal Polarization in Relativistic Heavy-Ion Collisions at Very High Energy», *Phys. Rev. Lett.* **120**, 012302 (2018), [arXiv:1707.07984 \[nucl-th\]](#).
- [21] D.-X. Wei, W.-T. Deng, X.-G. Huang, «Thermal vorticity and spin polarization in heavy-ion collisions», *Phys. Rev. C* **99**, 014905 (2019), [arXiv:1810.00151 \[nucl-th\]](#).
- [22] X.-L. Xia, H. Li, Z.-B. Tang, Q. Wang, «Probing vorticity structure in heavy-ion collisions by local Λ polarization», *Phys. Rev. C* **98**, 024905 (2018), [arXiv:1803.00867 \[nucl-th\]](#).
- [23] Y.B. Ivanov, V. Toneev, A. Soldatov, «Vorticity and Particle Polarization in Relativistic Heavy-Ion Collisions», *Phys. Atom. Nucl.* **83**, 179 (2020), [arXiv:1910.01332 \[nucl-th\]](#).
- [24] J.I. Kapusta, E. Rrapaj, S. Rudaz, «Hyperon polarization in relativistic heavy ion collisions and axial U(1) symmetry breaking at high temperature», *Phys. Rev. C* **101**, 031901 (2020), [arXiv:1910.12759 \[nucl-th\]](#).
- [25] J.-H. Gao, Z.-T. Liang, Q. Wang, X.-N. Wang, «Global Polarization Effect and Spin–Orbit Coupling in Strong Interaction», *Lect. Notes Phys.* **987**, 195 (2021), [arXiv:2009.04803 \[nucl-th\]](#).

- [26] X.-G. Deng, X.-G. Huang, Y.-G. Ma, S. Zhang, «Vorticity in low-energy heavy-ion collisions», *Phys. Rev. C* **101**, 064908 (2020), [arXiv:2001.01371 \[nucl-th\]](#).
- [27] D. Rindori, L. Tinti, F. Becattini, D. Rischke, «Relativistic quantum fluid with boost invariance», *Phys. Rev. D* **105**, 056003 (2022) [arXiv:2102.09016 \[hep-th\]](#).
- [28] S.Y.F. Liu, Y. Yin, «Spin polarization induced by the hydrodynamic gradients», *J. High Energy Phys.* **2107**, 188 (2021), [arXiv:2103.09200 \[hep-ph\]](#).
- [29] B. Fu *et al.*, «Shear-Induced Spin Polarization in Heavy-Ion Collisions», *Phys. Rev. Lett.* **127**, 142301 (2021), [arXiv:2103.10403 \[hep-ph\]](#).
- [30] D. She, A. Huang, D. Hou, J. Liao, «Relativistic Viscous Hydrodynamics with Angular Momentum», [arXiv:2105.04060 \[nucl-th\]](#).
- [31] J.-H. Gao, «Helicity polarization in relativistic heavy ion collisions», *Phys. Rev. D* **104**, 076016 (2021), [arXiv:2105.08293 \[hep-ph\]](#).
- [32] C. Yi, S. Pu, D.-L. Yang, «Reexamination of local spin polarization beyond global equilibrium in relativistic heavy ion collisions», *Phys. Rev. C* **104**, 064901 (2021), [arXiv:2106.00238 \[hep-ph\]](#).
- [33] H.-H. Peng, J.-J. Zhang, X.-L. Sheng, Q. Wang, «Ideal Spin Hydrodynamics from the Wigner Function Approach», *Chinese Phys. Lett.* **38**, 116701 (2021), [arXiv:2107.00448 \[hep-th\]](#).
- [34] S. Ryu, V. Jovic, C. Shen, «Probing early-time longitudinal dynamics with the Λ hyperon's spin polarization in relativistic heavy-ion collisions», *Phys. Rev. C* **104**, 054908 (2021), [arXiv:2106.08125 \[nucl-th\]](#).
- [35] D.-L. Wang, S. Fang, S. Pu, «Analytic solutions of relativistic dissipative spin hydrodynamics with Bjorken expansion», *Phys. Rev. D* **104**, 114043 (2021), [arXiv:2107.11726 \[nucl-th\]](#).
- [36] M. Hongo *et al.*, «Relativistic spin hydrodynamics with torsion and linear response theory for spin relaxation», *J. High Energy Phys.* **2111**, 150 (2021), [arXiv:2107.14231 \[hep-th\]](#).
- [37] F. Becattini *et al.*, «Local Polarization and Isothermal Local Equilibrium in Relativistic Heavy Ion Collisions», *Phys. Rev. Lett.* **127**, 272302 (2021), [arXiv:2103.14621 \[nucl-th\]](#).
- [38] F. Becattini, M. Buzzegoli, A. Palermo, «Spin-thermal shear coupling in a relativistic fluid», *Phys. Lett. B* **820**, 136519 (2021), [arXiv:2103.10917 \[nucl-th\]](#).
- [39] L. Dong, S. Lin, «Dilepton Helical Production in a Vortical Quark–Gluon Plasma», [arXiv:2112.07153 \[hep-ph\]](#).
- [40] W. Florkowski, A. Kumar, A. Mazeliauskas, R. Ryblewski, «Effect of thermal shear on longitudinal spin polarization in a thermal model», [arXiv:2112.02799 \[hep-ph\]](#).

- [41] W. Florkowski, A. Kumar, R. Ryblewski, A. Mazeliauskas, «Longitudinal spin polarization in a thermal model», *Phys. Rev. C* **100**, 054907 (2019), [arXiv:1904.00002 \[nucl-th\]](#).
- [42] W. Florkowski, B. Friman, A. Jaiswal, E. Speranza, «Relativistic fluid dynamics with spin», *Phys. Rev. C* **97**, 041901 (2018), [arXiv:1705.00587 \[nucl-th\]](#).
- [43] W. Florkowski *et al.*, «Spin-dependent distribution functions for relativistic hydrodynamics of spin-1/2 particles», *Phys. Rev. D* **97**, 116017 (2018), [arXiv:1712.07676 \[nucl-th\]](#).
- [44] W. Florkowski, A. Kumar, R. Ryblewski, «Thermodynamic *versus* kinetic approach to polarization-vorticity coupling», *Phys. Rev. C* **98**, 044906 (2018), [arXiv:1806.02616 \[hep-ph\]](#).
- [45] W. Florkowski, R. Ryblewski, A. Kumar, «Relativistic hydrodynamics for spin-polarized fluids», *Prog. Part. Nucl. Phys.* **108**, 103709 (2019), [arXiv:1811.04409 \[nucl-th\]](#).
- [46] F. Becattini, W. Florkowski, E. Speranza, «Spin tensor and its role in non-equilibrium thermodynamics», *Phys. Lett. B* **789**, 419 (2019), [arXiv:1807.10994 \[hep-th\]](#).
- [47] W. Florkowski, A. Kumar, R. Ryblewski, R. Singh, «Spin polarization evolution in a boost invariant hydrodynamical background», *Phys. Rev. C* **99**, 044910 (2019), [arXiv:1901.09655 \[hep-ph\]](#).
- [48] R. Singh, G. Sophys, R. Ryblewski, «Spin polarization dynamics in the Gubser-expanding background», *Phys. Rev. D* **103**, 074024 (2021), [arXiv:2011.14907 \[hep-ph\]](#).
- [49] S. Bhadury *et al.*, «Relativistic dissipative spin dynamics in the relaxation time approximation», *Phys. Lett. B* **814**, 136096 (2021), [arXiv:2002.03937 \[hep-ph\]](#).
- [50] S. Bhadury *et al.*, «Dissipative spin dynamics in relativistic matter», *Phys. Rev. D* **103**, 014030 (2021), [arXiv:2008.10976 \[nucl-th\]](#).
- [51] E. Speranza, N. Weickgenannt, «Spin tensor and pseudo-gauges: from nuclear collisions to gravitational physics», *Eur. Phys. J. A* **57**, 155 (2021), [arXiv:2007.00138 \[nucl-th\]](#).
- [52] S. Shi, C. Gale, S. Jeon, «From chiral kinetic theory to relativistic viscous spin hydrodynamics», *Phys. Rev. C* **103**, 044906 (2021), [arXiv:2008.08618 \[nucl-th\]](#).
- [53] R. Singh, M. Shokri, R. Ryblewski, «Spin polarization dynamics in the Bjorken-expanding resistive MHD background», *Phys. Rev. D* **103**, 094034 (2021), [arXiv:2103.02592 \[hep-ph\]](#).
- [54] A.D. Gallegos, U. Gürsoy, A. Yarom, «Hydrodynamics of spin currents», *SciPost Phys.* **11**, 041 (2021), [arXiv:2101.04759 \[hep-th\]](#).
- [55] K. Hattori *et al.*, «Fate of spin polarization in a relativistic fluid: An entropy-current analysis», *Phys. Lett. B* **795**, 100 (2019), [arXiv:1901.06615 \[hep-th\]](#).

- [56] K. Fukushima, S. Pu, «Spin hydrodynamics and symmetric energy-momentum tensors – A current induced by the spin vorticity –, *Phys. Lett. B* **817**, 136346 (2021), [arXiv:2010.01608 \[hep-th\]](#).
- [57] S. Li, M.A. Stephanov, H.-U. Yee, «Nondissipative Second-Order Transport, Spin, and Pseudogauge Transformations in Hydrodynamics», *Phys. Rev. Lett.* **127**, 082302 (2021), [arXiv:2011.12318 \[hep-th\]](#).
- [58] A.D. Gallegos, U. Gürsoy, «Holographic spin liquids and Lovelock Chern-Simons gravity», *J. High Energy Phys.* **2011**, 151 (2020), [arXiv:2004.05148 \[hep-th\]](#).
- [59] M. Garbiso, M. Kaminski, «Hydrodynamics of simply spinning black holes and hydrodynamics for spinning quantum fluids», *J. High Energy Phys.* **2012**, 112 (2020), [arXiv:2007.04345 \[hep-th\]](#).
- [60] Y. Hidaka, D.-L. Yang, «Nonequilibrium chiral magnetic/vortical effects in viscous fluids», *Phys. Rev. D* **98**, 016012 (2018), [arXiv:1801.08253 \[hep-th\]](#).
- [61] D.-L. Yang, K. Hattori, Y. Hidaka, «Effective quantum kinetic theory for spin transport of fermions with collisional effects», *J. High Energy Phys.* **2020**, 070 (2020), [arXiv:2002.02612 \[hep-ph\]](#).
- [62] N. Weickgenannt *et al.*, «Generating Spin Polarization from Vorticity through Nonlocal Collisions», *Phys. Rev. Lett.* **127**, 052301 (2021), [arXiv:2005.01506 \[hep-ph\]](#).
- [63] W. Florkowski, R. Ryblewski, R. Singh, G. Sophys, «Spin polarization dynamics in the non-boost-invariant background», *Phys. Rev. D* **105**, 054007 (2022), [arXiv:2112.01856 \[hep-ph\]](#).
- [64] X.-L. Sheng *et al.*, «From Kadanoff–Baym to Boltzmann equations for massive spin-1/2 fermions», *Phys. Rev. D* **104**, 016029 (2021), [arXiv:2103.10636 \[nucl-th\]](#).
- [65] A. Das *et al.*, «Semi-classical kinetic theory for massive spin-half fermions with leading-order spin effects», [arXiv:2203.15562 \[hep-th\]](#).
- [66] R. Singh, M. Shokri, S.M.A.T. Mehr, «Relativistic magnetohydrodynamics with spin», [arXiv:2202.11504 \[hep-ph\]](#).
- [67] V.E. Ambrus, R. Ryblewski, R. Singh, «Spin waves in spin hydrodynamics», [arXiv:2202.03952 \[hep-ph\]](#).
- [68] S. Bhadury *et al.*, «Relativistic spin-magnetohydrodynamics», [arXiv:2204.01357 \[nucl-th\]](#).
- [69] N. Weickgenannt, D. Wagner, E. Speranza, D. Rischke, «Relativistic second-order dissipative spin hydrodynamics from the method of moments», [arXiv:2203.04766 \[nucl-th\]](#).
- [70] A. Daher, A. Das, W. Florkowski, R. Ryblewski, «Equivalence of canonical and phenomenological formulations of spin hydrodynamics», [arXiv:2202.12609 \[nucl-th\]](#).
- [71] S. De Groot, W. Van Leeuwen, C. Van Weert, «Relativistic Kinetic Theory. Principles and Applications», *North Holland*, 1980.