NON-PERTURBATIVE DETERMINATION OF THE COLLISIONAL BROADENING KERNEL AND MEDIUM-INDUCED RADIATION IN QCD PLASMAS*

Sören Schlichting

Fakultät für Physik, Universität Bielefeld 33615 Bielefeld, Germany

Ismail Soudi

Department of Physics and Astronomy, Wayne State University Detroit, MI 48201, USA

Received 25 July 2022, accepted 14 August 2022, published online 14 December 2022

Collisional broadening in QCD plasmas leads to the emission of mediuminduced radiation, which governs the energy loss of highly energetic particles or jets. While recent studies have obtained non-perturbative contributions to the collisional broadening kernel $C(b_{\perp})$ using lattice simulation of the dimensionally reduced long-distance effective theory of QCD, Electrostatic QCD (EQCD), so far all phenomenological calculations of jet quenching rely on perturbative determinations of the collisional broadening kernel. By matching the short-distance behavior of the lattice extracted EQCD broadening kernel, we determine the fully matched QCD broadening kernel non-perturbatively. We present results for the collisional broadening kernel in impact parameter $(C_{\text{QCD}}(b_{\perp}))$ and momentum space $(C_{\text{QCD}}(q_{\perp}))$, and employ them to determine the rates of medium-induced radiation in infinite and finite size QCD plasmas. By contrasting our results with leading and next-to-leading order perturbative determinations as well as various approximations of the splitting rates employed in the literature, we investigate the effect of the non-perturbative determination of $C_{\text{QCD}}(q_{\perp})$ on medium-induced radiation rates.

DOI:10.5506/APhysPolBSupp.16.1-A121

1. Introduction

During ultra-relativistic heavy-ion collisions, high energetic particles originating from the initial hard partonic collisions must traverse and interact

^{*} Presented by I. Soudi at the 29th International Conference on Ultrarelativistic Nucleus–Nucleus Collisions: Quark Matter 2022, Kraków, Poland, 4–10 April, 2022.

with the Quark–Gluon Plasma (QGP) before reaching the detector. Interactions of highly energetic parton with the medium have been studied extensively, and it was shown that besides the elastic interactions with the medium constituents, multiple soft scatterings between the hard partons and the medium trigger the partons to radiate. These soft scatterings lead to an infinite number of diagrams, which can be resummed to obtain an effective $1 \rightarrow 2$ in-medium radiation rates. Several formalisms for in-medium radiation have been developed in the literature [1–3], which encapsulate the same physics but differ on the level of approximations. However, all these formalisms rely on a description of the elastic scattering with the medium, obtained using the transverse momentum broadening kernel

$$C(q_{\perp}) \equiv \frac{(2\pi)^2 \mathrm{d}^3 \Gamma}{\mathrm{d}^2 q_{\perp} \mathrm{d} L},\tag{1}$$

which describes the rate at which the hard partons exchange transverse momentum q_{\perp} with the medium. To obtain the broadening kernel, three treatments of the medium are widely used in the literature:

- Many random, static, screened color centers [1]: $C(q_{\perp}) \propto \frac{1}{(q_{\perp}^2 + m_D^2)^2}$.
- Dynamical moving charges at lowest order in perturbation theory [4]: $C(q_{\perp}) \propto \frac{1}{q_{\perp}^2(q_{\perp}^2 + m_{\rm D}^2)}.$
- Many individually small scatterings, leading to transverse momentum diffusion [5]: $C(b_{\perp}) = \hat{q} b_{\perp}^2/4$.

Recently, there has been progress in obtaining non-perturbative contributions to the broadening kernel [6, 7], using a dimensionally reduced longdistance effective theory for QCD, 3D Electrostatic QCD (EQCD) [8]. After a Fourier transform of the broadening kernel to impact parameter space, lattice simulations of EQCD [6, 9, 10] can be performed to obtain the largedistance behavior of the broadening kernel $C_{EQCD}(b_{\perp})$, which was matched to the short-distance behavior of QCD to obtain a non-perturbative determination of $C_{QCD}(b_{\perp})$ in QCD at all scales [7]. In these proceedings, we employ the broadening kernel extracted to obtain the in-medium splitting rates of a highly energetic parton traversing a medium with a finite extent and compare with results obtained using perturbative results of the broadening kernels.

2. Non-perturbative broadening kernel

The non-perturbative (NP) broadening kernel $(C_{\text{QCD}}(b_{\perp}))$ in impact parameter space b_{\perp} was obtained in [7] and shown in the left panel of Fig. 1. However, in order to compute in-medium splitting rates in a QCD medium



Fig. 1. Left: Non-perturbative elastic broadening kernel $C_{\rm QCD}(b_{\perp})$ in impact parameter space. Data points for two different temperatures T = 250,500 MeV are shown alongside the interpolating splines. Right: Elastic broadening kernel $C_{\rm QCD}(\boldsymbol{q}_{\perp})$ in momentum space for T = 250,500 MeV. We also compare the kernel to leading-order (LO) and next-to-leading order (NLO) determinations at T = 500 MeV.

of finite size, it is more preferable to work in momentum space. Therefore, we perform a Fourier transform to obtain the non-perturbative broadening kernel $(C_{\rm QCD}(q_{\perp}))$ in impact parameter space q_{\perp} which is shown for T = 250,500 MeV in the right panel of Fig. 1 (*cf.* [11] for details of the procedure). We find that both data sets at T = 250,500 MeV display very similar behavior when expressing $TC_{\rm QCD}(q_{\perp})/C_{\rm R}$ as a function of $q_{\perp}/g_{\rm s}T$ in momentum space, or $C(b_{\perp})/g_{\rm s}^2 T C_{\rm R}$ as a function of $g_{\rm s} T b_{\perp}$ in impact parameter space. When comparing with leading order (LO) and next-toleading order (NLO) perturbative broadening kernels, we find that while they all follow the same ultraviolet (UV) behavior $\sim 1/q_{\perp}^4$ and, respectively, $\sim b_{\perp}^2 \log(b_{\perp})$ behavior, the infrared (IR) behavior of the NP and NLO kernels is markedly different from the LO. The NP and NLO kernels in UV follow a $\sim 1/q_{\perp}^3$ and, respectively, $\sim b_{\perp}$. However, the slopes of the kernels differ by a prefactor due to the difference in the string tension $\sigma_{\rm EQCD}$.

3. In-medium splitting rates

Employing the different broadening kernels and following [3], we obtain the rate of medium-induced radiation $\frac{d\Gamma(P,z;t)}{dz}$, the rate of quark with momentum P radiating a gluon with momentum zP after a time t spent in the medium. The results for momenta fraction z = 0.05 and z = 0.5 are shown in Fig. 2 as a function of time t, where we used the LO, NLO, and NP broadening kernels to obtain the square, circle and triangular points, and we show the ratio to the LO curve in the lower inset of each panel. Ini-



Fig. 2. (Color online) Splitting rate for the medium-induced emission of a gluon from a parent quark with energy P = 300 T in an equilibrium plasma with the temperature T = 500 MeV as a function of the evolution time t for different gluon momentum fraction z = 0.05, 0.5. (top) Different curves in each panel show the results for the different, leading order (LO), next-to-leading order (NLO), and non-perturbative (NP) momentum broadening kernels in Fig. 1. (bottom) We compare different approximations of the in-medium splitting rate, namely the opacity expansion at N = 1 [1], the resummed opacity rate of [12] (N = X), and the NLO expansion around the Harmonic Oscillator [13] (NLO-HO) to the full result (T = 500 MeV). Note that all results are obtained with the non-perturbative collision kernel. The lower panel of each plot displays the ratio to the full rate.

tially, we observe a linear behavior of the rate which saturates rather quickly at later times and recovers the infinite medium rates which were obtained earlier [7]. We find that the non-perturbative result starts lower than the LO rates before it settles above the LO and below the NLO, which can be explained by the fact that the early times radiation are driven mainly by single hard scatterings with large momentum exchange $(q_{\perp} \gg m_{\rm D})$, where the NP kernel is below the LO. At late times, multiple soft scatterings with small momentum exchange $(q_{\perp} \sim m_{\rm D})$ are more important, and the NP kernel in this IR behaves similarly to the NLO kernel. Overall, the results obtained using the NP kernel do not depart from a band of $\pm 50\%$ around LO, while the NLO results can become over $2 \times$ larger than the LO result.

Additionally, we computed the in-medium radiation rates using several approximations developed in the literature:

- First order opacity expansion, which considers a perturbative expansion in the number of re-scatterings with the medium. Here, we take only a single scattering (N = 1) [1].
- Resummed opacity expansion (N = X), which re-sums the opacity expansion by employing a cut-off to the momentum exchange of the scattering, the collision integral of subsequent scatterings can then be exponentiated (cf. [12]).
- Harmonic oscillator expansion (NLO-HO) where using an opacity expansion, a single hard scattering is considered as a perturbation on top of fully resummed multiple soft scatterings computed in the diffusion approximation [13].

Extensive details of the calculation of each approximation are given in [11]. We employ these approximations to obtain the in-medium splitting rate for the same NP broadening kernel $C_{\rm QCD}(q_{\perp})$ at T = 500 MeV and show a comparison in Fig. 2. While the naive first order opacity expansion reproduces the linear behavior of the rate at early times, it breaks down later on when the time is large enough to allow for subsequent scatterings. Conversely, the resummed (N = X) opacity expansion is able to reproduce the rate even at late times. For the NLO expansion around the HO, we observe that it can be quite accurate throughout the evolution, especially for quasi-democratic splittings $(z \sim 1/2)$.

4. Conclusion

In these proceedings, we presented our results for in-medium splitting rates computed using the non-perturbative collision kernel and compared them to the calculations using leading and next-to-leading order perturbative collision kernel as well as to different approximations of the in-medium splitting rates. We find that approximations to the splitting rate calculation are quite effective in reproducing the rate within their respective range of validity. Conversely, we observed considerable differences between the results for different collision kernels. Since the differences between the nonperturbative and the LO kernel, which is usually used in phenomenological studies of jet quenching, can be of the order of 30%, it will be interesting to further explore the phenomenological consequences. In this context, it is S. Schlichting, I. Soudi

important to point out that the collisional broadening kernel and in-medium splitting rates obtained in this study can be easily integrated into models of jet quenching using a kinetic approach or Monte Carlo simulations [14].

We thank Guy D. Moore and Niels Schlusser for insightful discussions and collaboration on our previous work [7]. This work is supported in part by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the CRC-TR 211 'Strong-interaction matter under extreme conditions' — project number 315477589–TRR 211 and the German Bundesministerium für Bildung und Forschung (BMBF) through grant No. 05P21PBCAA. The authors also gratefully acknowledge the computing time provided by the Paderborn Center for Parallel Computing (PC2).

REFERENCES

- M. Gyulassy, P. Levai, I. Vitev, Nucl. Phys. B 571, 197 (2000), arXiv:hep-ph/9907461.
- [2] P.B. Arnold, G.D. Moore, L.G. Yaffe, J. High Energy Phys. 0206, 030 (2002), arXiv:hep-ph/0204343.
- [3] S. Caron-Huot, C. Gale, *Phys. Rev. C* 82, 064902 (2010), arXiv:1006.2379 [hep-ph].
- [4] P. Aurenche, F. Gelis, H. Zaraket, J. High Energy Phys. 0205, 043 (2002), arXiv:hep-ph/0204146.
- [5] P.B. Arnold, Phys. Rev. D 79, 065025 (2009), arXiv:0808.2767 [hep-ph].
- [6] G.D. Moore, N. Schlusser, *Phys. Rev. D* 101, 014505 (2020), arXiv:1911.13127 [hep-lat]; *Erratum ibid.* 101, 059903 (2020).
- [7] G.D. Moore, S. Schlichting, N. Schlusser, I. Soudi, J. High Energy Phys. 2110, 059 (2021), arXiv:2105.01679 [hep-ph].
- [8] S. Caron-Huot, *Phys. Rev. D* 79, 065039 (2009), arXiv:0811.1603 [hep-ph].
- [9] M. Panero, K. Rummukainen, A. Schäfer, *Phys. Rev. Lett.* 112, 162001 (2014), arXiv:1307.5850 [hep-ph].
- [10] M. D'Onofrio, A. Kurkela, G.D. Moore, J. High Energy Phys. 1403, 125 (2014), arXiv:1401.7951 [hep-lat].
- [11] S. Schlichting, I. Soudi, *Phys. Rev. D* 105, 7 (2022), arXiv:2111.13731 [hep-ph].
- [12] C. Andres, F. Dominguez, M. Gonzalez Martinez, J. High Energy Phys. 2103, 102 (2021), arXiv:2011.06522 [hep-ph].
- [13] Y. Mehtar-Tani, J. High Energy Phys. 1907, 057 (2019), arXiv:1903.00506 [hep-ph].
- [14] R.M. Yazdi, S. Shi, C. Gale, S. Jeon, arXiv:2206.05855 [hep-ph].