

MEASUREMENT OF MEDIUM-INDUCED
MODIFICATION OF JET YIELD AND
ACOPLANARITY USING SEMI-INCLUSIVE $\gamma_{\text{dir}}+\text{JET}$
AND $\pi^0+\text{JET}$ DISTRIBUTIONS IN $p+p$ AND CENTRAL
Au+Au COLLISIONS AT $\sqrt{s_{NN}} = 200$ GeV BY STAR*

DEREK ANDERSON

for the STAR Collaboration

Cyclotron Institute, Texas A&M University, College Station, TX 77843, USA

*Received 11 August 2022, accepted 14 September 2022,
published online 14 December 2022*

The STAR Collaboration presents measurements of semi-inclusive distributions of charged jets recoiling from high transverse energy (E_T) direct photon and π^0 triggers in $p+p$ and central Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. Jets are reconstructed from charged particles using the anti- k_T algorithm with jet resolution parameters $R = 0.2$ and 0.5 . The large uncorrelated background in central Au+Au collisions is corrected using a mixed-event technique. This enables a jet measurement extending to low transverse momentum and large R with well-controlled systematic uncertainties. We present measurements of the jet R dependence of suppression, intra-jet broadening, and acoplanarity of $\pi^0+\text{jet}$ and $\gamma_{\text{dir}}+\text{jet}$ for trigger E_T (E_T^{trig}) between 9–20 GeV.

DOI:10.5506/APhysPolBSupp.16.1-A55

1. Introduction

Heavy-ion collisions at RHIC and the LHC produce a medium of deconfined partons, the Quark–Gluon Plasma (QGP) [1]. Hard (high-momentum transfer, Q^2) interactions of quarks and gluons in such collisions generate energetic scattered partons which propagate through the medium and interact with it. Consequently, the parton showers are modified (jet quenching) [2]. Jet quenching manifests in several observable effects: transport of energy outside of the reconstructed jet cone, modification of the jet substructure, and enhanced acoplanarity ($\Delta\phi = \phi_{\text{trig}} - \phi_{\text{jet}}$) [4]. While the $\Delta\phi$ distribution has a finite width in vacuum due to the Sudakov radiation [3], the

* Presented at the 29th International Conference on Ultrarelativistic Nucleus–Nucleus Collisions: Quark Matter 2022, Kraków, Poland, 4–10 April, 2022.

presence of a medium may further broaden it due to mechanisms such as multiple in-medium soft scatterings [4], the hard scattering of a parton off QGP quasi-particles [5], and medium response [6].

In these proceedings, the STAR Collaboration reports on measurements of the semi-inclusive yields of jets recoiling from *direct photons* (γ_{dir}) and π^0 , together with their acoplanarity distributions in $p + p$ and central Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV. Simultaneous measurements of these different observables in the same analysis promise a discriminating and multi-messenger approach to the study of jet quenching.

Since direct photons are color neutral, they do not interact with the QGP; their measured energy thereby reflects the Q^2 of the hard interaction and provides a constraint on the initial energy of the recoiling jet. Hence, the measurement of jets coincident with a γ_{dir} ($\gamma_{\text{dir}} + \text{jet}$) provides a valuable tool for quantifying the effects of jet quenching [7]. In addition, comparison with jets coincident with π^0 ($\pi^0 + \text{jet}$) may elucidate the color factor and path length dependence of medium-induced energy loss due to differences between the recoil jet populations of the two triggers in their relative quark/gluon fraction and mean path length [8].

STAR has previously reported on the yield suppression of charged hadrons coincident with π^0 and γ_{dir} triggers [9]. Additionally, STAR has measured the yield of reconstructed charged-particle jets coincident with charged-hadron triggers ($h^\pm + \text{jet}$) using a semi-inclusive approach [10]. In this approach, the large uncorrelated jet background in heavy-ion collisions is corrected with a Mixed Event (ME) technique, enabling the measurement of reconstructed jets at low transverse momentum (p_T) and large resolution parameter. In the current analysis, we combine the $\gamma_{\text{dir}}/\pi^0$ identification of [9] with the semi-inclusive and ME approach of [10] to measure the semi-inclusive $\gamma_{\text{dir}} + \text{jet}$ and $\pi^0 + \text{jet}$ yields in $p + p$ and central Au + Au collisions.

2. Analysis

Two STAR datasets of $\sqrt{s_{NN}} = 200$ GeV collisions are analyzed: a 10 nb^{-1} sample of Au + Au collisions recorded in 2014, and a 23 pb^{-1} sample of $p + p$ collisions recorded in 2009. Both were recorded using an online high-tower trigger, *i.e.*, a calorimeter tower above a certain threshold in energy. Two STAR subsystems are used: the Time Projection Chamber (TPC) [11], which provides charged-particle tracks for jet reconstruction, and the Barrel Electromagnetic Calorimeter (BEMC) [12], which is used to identify π^0 and γ_{dir} triggers.

Discrimination of π^0 and γ_{dir} candidates in the BEMC is carried out using the Transverse Shower Profile (TSP) method [9, 13]. Based on the TSP, the data are separated into two samples: a nearly pure sample of identified π^0 , and a sample with an enhanced fraction of γ_{dir} (γ_{rich}).

Triggers are selected offline to satisfy $E_T^{\text{trig}} = 9\text{--}20$ GeV and $|\eta^{\text{trig}}| < 0.9$. The purity of the γ_{rich} sample, *i.e.*, the percentage of γ_{rich} that are actually γ_{dir} , is determined via a data-driven method [9, 13]. The γ_{dir} +jet distribution is then determined from the γ_{rich} sample via a statistical subtraction, which removes contamination due to hadronic decays and fragmentation photons to the extent that their near-side azimuthal correlations are identical to those of the identified π^0 [9, 13].

Jets are reconstructed from the TPC tracks using the anti- k_T algorithm [14, 15] for two resolution parameters, $R = 0.2$ and 0.5 . Reconstructed jets are subjected to the same fiducial cuts as in [10].

In Au + Au collisions, there is a substantial background yield of jet candidates which are not correlated with the trigger. This background yield is removed using the ME technique described in [10]. Uncorrelated jet yield is small in $p + p$ collisions, and no correction for it is applied. The residual jet p_T -smearing is corrected in two steps [10]: first, jets are corrected for an event-wise energy pedestal, and then residual fluctuations caused by detector effects ($p + p$ and Au + Au collisions) and the heavy-ion background (Au + Au collisions only) are corrected using regularized unfolding. We use $p_{T,\text{jet}}^{\text{reco,ch}}$ (where the superscript “ch” denotes “charged jets”) to refer to the jet p_T after the event-wise pedestal correction, and $p_{T,\text{jet}}^{\text{ch}}$ to the jet p_T after unfolding.

The two-dimensional acoplanarity distributions must also be unfolded for both $p_{T,\text{jet}}^{\text{reco,ch}}$ and $\Delta\phi$ fluctuations. Note that the $\Delta\phi$ distributions shown here have however been unfolded for $p_{T,\text{jet}}^{\text{reco,ch}}$ fluctuations only. We estimate that $\Delta\phi$ smearing effects are small.

3. Results

Jet distributions are reported in two ways: the two-dimensional measurement of $\Delta\phi$ *vs.* $p_{T,\text{jet}}^{\text{ch}}$, and the one-dimensional measurement of $p_{T,\text{jet}}^{\text{ch}}$ for recoil jets, which satisfy $|\Delta\phi - \pi| < \pi/4$. The recoil jet $p_{T,\text{jet}}^{\text{ch}}$ distributions in central Au + Au and $p + p$ are compared against PYTHIA 8 with the Monash tune [16]. The PYTHIA 8 distributions are smeared to account for a trigger energy resolution. We report two different ratios of the trigger-normalized recoil jet yields: I_{AA} , the ratio of the semi-inclusive yield of recoil jets in Au + Au over that in $p + p$ for fixed R ; and $\mathfrak{R}^{0.2/0.5}$, the ratio of the semi-inclusive yield for $R = 0.2$ relative to that for $R = 0.5$, for fixed collision system.

Figure 1 shows I_{AA} for $E_T^{\text{trig}} = 11\text{--}15, 15\text{--}20$ GeV π^0 and γ_{dir} triggers. The recoil jet yield for $R = 0.2$ is systematically more suppressed than that for $R = 0.5$. In addition, the value of I_{AA} is observed to be consistent within uncertainties between π^0 and γ_{dir} for both values of R , despite differences

in the recoil jet quark/gluon fraction and mean path length. Note, however, that the $\gamma_{\text{dir}} + \text{jet}$ $p_{T,\text{jet}}^{\text{ch}}$ spectrum is steeper, so a similar magnitude of yield suppression corresponds to smaller medium-induced out-of-cone energy loss.




Fig. 1. (Color online) I_{AA} for $\pi^0 + \text{jet}$ (dark gray/blue) and $\gamma_{\text{dir}} + \text{jet}$ (gray/red). Dark bands indicate statistical errors, and light bands indicate systematic uncertainties.

Figure 2 shows the $\mathfrak{R}^{0.2/0.5}$ for $E_T^{\text{trig}} = 11\text{--}15$ GeV π^0 (upper panel) and $E_T^{\text{trig}} = 15\text{--}20$ GeV γ_{dir} (lower panel). We see that $\mathfrak{R}^{0.2/0.5}$ for $p + p$ is less than unity and that PYTHIA 8 reproduces the ratio well. However, the value of $\mathfrak{R}^{0.2/0.5}$ for central Au + Au is significantly lower than that for $p + p$ and PYTHIA 8.




Fig. 2. (Color online) $\mathfrak{R}^{0.2/0.5}$ for π^0 (upper panel) and γ_{dir} (lower panel) triggers from $p + p$ (light gray/green), Au+Au (dark gray/blue and gray/red), and PYTHIA 8 (black dashed lines).

Figures 1 and 2 show a clear observation of significant medium-induced intra-jet broadening in central Au + Au collisions at RHIC.

Figure 3(a) shows the corrected $\Delta\phi$ correlations in $p + p$ collisions between $E_T^{\text{trig}} = 9\text{--}11$ GeV π^0 triggers and $R = 0.5$ jets (boxes). These distributions are reproduced well by PYTHIA 8 (dotted lines) for all three ranges of $p_{T,\text{jet}}^{\text{ch}}$ (5–10, 10–15, and 15–20 GeV/c).




Fig. 3. Corrected $R = 0.5$ $\Delta\phi$ distributions in $p + p$ (a) and Au + Au (b) collisions for π^0 ($p + p$ and Au + Au) and γ_{dir} (Au + Au only) triggers. Vertical lines indicate statistical errors, and filled and open boxes indicate uncorrelated and correlated systematic uncertainties, respectively (note that the statistical errors are smaller than the marker size for the Au + Au data points). Dotted and dashed lines are PYTHIA 8.

Figure 3(b) then shows the corrected $\Delta\phi$ correlations in Au + Au collisions between $E_T^{\text{trig}} = 11\text{--}15$ GeV π^0 and γ_{dir} triggers and recoil jets of $R = 0.5$ and $p_{T,\text{jet}}^{\text{ch}} = 10\text{--}15$ GeV/c. The dashed lines are the corresponding distributions from PYTHIA 8, that is validated in the left panel. We observe a marked enhancement in yield at wide angles (small $\Delta\phi$) in central Au + Au collisions relative to vacuum fragmentation. This is the first observation of significant medium-induced modification of $\pi^0 + \text{jet}$ and $\gamma_{\text{dir}} + \text{jet}$ acoplanarity at low $p_{T,\text{jet}}^{\text{ch}}$ in central Au + Au collisions at RHIC.

4. Summary

STAR has measured the R dependence of recoil jet yield and acoplanarity using the semi-inclusive distributions of charged-particle jets recoiling from

π^0 and γ_{dir} triggers in central Au + Au and $p + p$ collisions at $\sqrt{s_{NN}} = 200$ GeV. Model calculations based on the PYTHIA 8 event generator are found to be consistent with the measurements in $p + p$ collisions.

We have reported on both the recoil yield in a fixed angular window as a function of $p_{T,\text{jet}}^{\text{ch}}$ and the distribution of acoplanarity at fixed $p_{T,\text{jet}}^{\text{ch}}$. We observe marked medium-induced intra-jet broadening. We also observe clear medium-induced acoplanarity at low jet $p_{T,\text{jet}}^{\text{ch}}$, which may arise from in-medium jet scattering or from the contribution of medium response to the jet signal. To further investigate the medium-induced acoplanarity and disentangle the underlying mechanisms, it will be essential to extend the kinematic range of this measurement in heavy-ion collisions and compare against theoretical calculations.

This work is funded in part by the United States Department of Energy under grant number DE-SC0015636.

REFERENCES

- [1] W. Busza, K. Rajagopal, W. van der Schee, *Annu. Rev. Nucl. Part. Sci.* **68**, 339 (2018).
- [2] L. Cunqueiro, A.M. Sickles, *Prog. Part. Nucl. Phys.* **124**, 103940 (2022), [arXiv:2110.14490 \[nucl-ex\]](#).
- [3] P. Sun, C.-P. Yuan, F. Yuan, *Phys. Rev. D* **92**, 094007 (2015).
- [4] A.H. Mueller, B. Wu, B.-W. Xiao, F. Yuan, *Phys. Lett. B* **763**, 208 (2016).
- [5] F. D’Eramo, M. Lekaveckas, H. Liu, K. Rajagopal, *J. High Energy Phys.* **2013**, 31 (2013).
- [6] G. Milhano, U.A. Wiedemann, K.C. Zapp, *Phys. Lett. B* **779**, 409 (2018).
- [7] X.-N. Wang, Z. Huang, I. Sarcevic, *Phys. Rev. Lett.* **77**, 231 (1996).
- [8] T. Renk, *Phys. Rev. C* **88**, 054902 (2013).
- [9] STAR Collaboration (L. Adamczyk *et al.*), *Phys. Lett. B* **760**, 689 (2016).
- [10] STAR Collaboration (L. Adamczyk *et al.*), *Phys. Rev. C* **96**, 024905 (2017).
- [11] M. Anderson *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **499**, 659 (2003).
- [12] M. Beddo *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **499**, 725 (2003).
- [13] STAR Collaboration (B.I. Abelev *et al.*), *Phys. Rev. C* **82**, 034909 (2010).
- [14] M. Cacciari, G.P. Salam, G. Soyez, *J. High Energy Phys.* **0804**, 063 (2008).
- [15] M. Cacciari, G.P. Salam, G. Soyez, *Eur. Phys. J. C* **72**, 1896 (2012).
- [16] T. Sjöstrand, S. Mrenna, P.Z. Skands, *Comput. Phys. Commun.* **178**, 852 (2008).