

# NEWS FROM THE STRONG INTERACTIONS PROGRAM OF NA61/SHINE\* \*\*

ANGELIKA TEFELSKA

for the NA61/SHINE Collaboration

Warsaw University of Technology, Faculty of Physics, Poland

*Received 17 February 2023, accepted 24 August 2023,  
published online 27 October 2023*

The NA61/SHINE experimental physics program focuses on searching for the critical point and studying the properties of the onset of deconfinement in the strongly interacting matter. A two-dimensional scan is performed by varying the beam momentum (from 13A to 150/158A GeV/c) and the system size (from  $p+p$  to Pb+Pb) of the collided nuclei. This contribution presents the most recent results from the NA61/SHINE strong interactions program and includes future data-taking and analysis plans.

DOI:10.5506/APhysPolBSupp.16.8-A1

## 1. Introduction

The NA61/SHINE is a large acceptance, fixed target experiment, which uses eight Time Projection Chambers and a Time-of-Flight detector to identify the produced particles and measure their momenta. The collision centrality is determined based on measurements from the high-resolution forward calorimeter, the Projectile Spectator Detector (PSD).

The large variety of beams and targets allowed to perform two-dimensional scan by varying the beam momentum (from 13A to 150/158A GeV/c) and the system size (from  $p+p$  to Pb+Pb) of the collided nuclei. The main goals of the strong interactions program are searching for the critical point of strongly interacting matter and studying the properties of the onset of deconfinement.

---

\* Presented at *Excited QCD 2022*, Sicily, Italy, 23–29 October, 2022.

\*\* This work was supported by the Polish Ministry of Science and Higher Education (grant WUT ID-UB), the Norwegian Financial Mechanism 2014–2021 (grant 2019/34/H/ST2/00585), the Polish Minister of Education and Science (contract No. 2021/WK/10).

## 2. Study of the properties of the onset of deconfinement

The Statistical Model of the Early Stage [1] assumes a first-order phase transition from the hadronic matter to the Quark–Gluon Plasma (QGP) between top AGS ( $\sqrt{s_{NN}} \approx 5$  GeV) and top SPS ( $\sqrt{s_{NN}} \approx 17$  GeV) energies.

One of the predicted signatures of the deconfinement transition is horn, a non-monotonic change of  $K^+/\pi^+$  ratio as a function of collision energy. The horn structure was observed by the NA49 experiment in central Pb+Pb collisions [2]. The NA61/SHINE extends the set of experimental data by  $p+p$ , Be+Be, and Ar+Sc collisions (Xe+La and Pb+Pb results are expected soon). The compilation of NA61/SHINE and world data is shown in Fig. 1. The plateau-like structures are visible in  $p+p$  and Be+Be interactions at SPS energies. The  $K^+/\pi^+$  ratio for Ar+Sc collisions is located between  $p+p$  and Pb+Pb data but no horn-like structure is visible.




Fig. 1. The energy dependence of the  $K^+/\pi^+$  particle yields ratio at mid-rapidity (left) and full acceptance (right) for the 20% most central Be+Be, central Pb+Pb, and Au+Au collisions, as well as inelastic  $p+p$  interactions. The NA61/SHINE  $p+p$  and Be+Be results are taken from Refs. [3, 4], whereas Ar+Sc points [5] are preliminary.

The  $K^+/\pi^+$  ratio at mid-rapidity and inverse slope parameter  $T$  fitted to the transverse momentum spectra of  $K^+$  mesons as a function of system size are shown in Fig. 2. All plots show similar, threshold-like behavior, which cannot be reproduced by any of the considered models. The observed rapid change of hadron production properties that starts when moving from Be+Be to Ar+Sc collisions at top SPS energies hints at the beginning of the creation of large clusters of strongly interacting matter called *the onset of fireball*.




Fig. 2. Top: The system-size dependence of the  $K^+/\pi^+$  ratio at mid-rapidity measured in NA61/SHINE  $p+p$ ,  $Be+Be$ ,  $Ar+Sc$ , and NA49  $Pb+Pb$  collisions at 150A ( $Be+Be$ ,  $Ar+Sc$ ) or 158A ( $p+p$ ,  $Pb+Pb$ ) GeV/c compared with dynamical (top left) and statistical (top right) models. Bottom: The inverse slope parameter  $T$ , fitted to  $K^+$  transverse momentum spectra at the similar collision energy [5]. The NA61/SHINE  $p+p$  and  $Be+Be$  results are taken from Refs. [3, 4], whereas  $Ar+Sc$  points [5] are preliminary.

### 3. Search for the critical point

The critical point (CP) signal should be visible as a non-monotonic dependence of various fluctuation/correlation measures. Figure 3 shows the energy dependence of fluctuations for negatively charged hadrons ( $h^-$ ) and net-electric charge ( $h^+ - h^-$ ), measured in  $p+p$ ,  $Be+Be$ , and  $Ar+Sc$  collisions. The comparison of fluctuations in systems of different sizes is possible using intensive quantities. For  $h^-$ , one considers ratios of cumulants  $\kappa_2/\kappa_1[h^-]$  (scaled variance),  $\kappa_3/\kappa_2[h^-]$  (scaled skewness), and  $\kappa_4/\kappa_2[h^-]$  (scaled kurtosis) (reference value of 1 is defined by the Poisson distribution). In the case of net-electric charge, the first two ratios are slightly modified to keep the ref-

erence value (1 for the Skellam distribution):  $\kappa_2[h^+ - h^-]/(\kappa_1[h^+] + \kappa_1[h^-])$ ,  $\kappa_3/\kappa_1[h^+ - h^-]$ . Figure 3 shows that in the case of  $h^-$ , only the scaled variance presents significant differences between heavier and lighter systems. In the case of net-electric charge, the scaled skewness and scaled kurtosis suggest a non-monotonic behavior, within sizeable systematic uncertainties (current subject of analysis).




Fig. 3. Preliminary results on the energy dependence of fluctuations for negatively charged hadrons (top) and net-electric charge (bottom) in  $p+p$ , Be+Be, and Ar+Sc collisions.

Another possible tool for the search for CP is an intermittency analysis. Generally, the scaled factorial moments  $F_r$  are calculated based on the counted particles in cells in transverse momentum space [6]. If the system freezes out in the vicinity of the critical point, the scaled factorial moments should reveal a power-law dependence:  $F_r(M) \sim M^{\phi_r}$ , where  $M$  is the number of cells [7]. In this analysis, the statistically-independent points are

used and instead of the  $(p_x, p_y)$  cells, the cumulative variables are used to remove the dependence of the moments on the momentum distribution [8]. Preliminary results on  $F_2(M)$  of mid-rapidity protons measured in the 0–20% most central Ar+Sc collisions at 150A GeV/c and 0–10% most central Pb+Pb collisions at 13A GeV/c are presented in Fig. 4 (top panels). The negatively charged hadron intermittency in 0–10% central Pb+Pb collisions at 30A GeV/c beam momentum up to the fourth scaled factorial moment is also shown in Fig. 4 (bottom panel). The measured  $F_2(M)$  of protons for Ar+Sc at 150A GeV/c and Pb+Pb at 13A GeV/c, as well as  $F_2(M)$ ,  $F_3(M)$ , and  $F_4(M)$  of negatively charged hadrons in Pb+Pb at 30A GeV/c show no indication for power-law increase with a bin size which could indicate CP.




Fig. 4. Top: Preliminary results on intermittency analysis of mid-rapidity protons for the 0–20% most central Ar+Sc collisions at 150A GeV/c (top left) and 0–10% most central Pb+Pb collisions at 13A GeV/c (top right). Bottom: Preliminary results on intermittency analysis of mid-rapidity negatively charged hadrons for the 0–10% most central Pb+Pb collisions at 30A GeV/c.

The last topic connected with searching for CP are symmetric Lévy HBT correlations for the same-charge pion pairs in central Be+Be and Ar+Sc collisions at 150A GeV/c beam momentum. Usually, the correlation function assumes a Gaussian source but it can be generalized by the Lévy-shaped formula,  $C(q) = 1 + \lambda e^{-(qR)^\alpha}$ , where the parameter  $\alpha$  describes the shape of the source (for  $\alpha = 2$ , the source is Gaussian, for  $\alpha = 1$ , the source is described by the Cauchy distribution). For the critical system, the  $\alpha$  parameter should be equal to 0.5 according to the prediction of the 3D Ising model [9]. The  $\alpha$  parameter as a function of transverse mass is shown in Fig. 5. The results do not indicate the critical point in Be+Be and Ar+Sc collisions. The values of  $\alpha$  parameters are between Gaussian and Cauchy shapes and might be a sign of anomalous diffusion. The  $R$  Lévy scale parameter is also presented in Fig. 5 and describes the length of homogeneity. The visible transverse mass dependence can be a sign of transverse flow.




Fig. 5. Preliminary results on the symmetric Lévy HBT correlations for the same-charge pion pairs in central Be+Be and Ar+Sc collisions at 150A GeV/c beam momentum. The panels show the dependence of the  $\alpha$  (left) and  $R$  (right) parameters on the transverse mass of the pair.

#### 4. New data on hadron spectra in $p + p$ reactions

Finally, NA61/SHINE provides new and unique results on strangeness production in  $p + p$  interactions. The spectra of strange mesons and multi-strange hyperons were measured with unprecedented precision at the SPS energy range. Figure 6 presents the rapidity distributions of  $K^*(892)^0$  [10] and  $K_S^0$  mesons (Ref. [11] and preliminary results) produced in inelastic  $p + p$  interactions at the SPS energies.

The high-statistics data of  $p + p$  collisions at 158 GeV/c allowed the more challenging measurement of  $\Xi(1530)^0$  and  $\bar{\Xi}(1530)^0$  hyperon production [12], which are the only such results at the SPS energy. The first two-dimensional spectra of  $\Xi(1530)^0$  and  $\bar{\Xi}(1530)^0$  [12] in  $y$  and  $p_T$  bins are presented in Fig. 7.




Fig. 6. Rapidity distributions of  $K^*(892)^0$  (top) produced in inelastic  $p + p$  interactions at 40–158 GeV/c [10] and  $K_S^0$  produced in inelastic  $p + p$  collisions at 158 GeV/c [11] (bottom left) and 80 GeV/c (bottom right).  $K_S^0$  results at 80 GeV/c are preliminary.




Fig. 7. Transverse momentum spectra in rapidity intervals of  $\Xi(1530)^0$  and  $\bar{\Xi}(1530)^0$  production in inelastic  $p + p$  collisions at 158 GeV/c [12].

## REFERENCES

- [1] M. Gaździcki, M.I. Gorenstein, *Acta Phys. Pol. B* **30**, 2705 (1999).
- [2] NA49 Collaboration (C. Alt *et al.*), *Phys. Rev. C* **77**, 024903 (2008).
- [3] NA61/SHINE Collaboration (A. Aduszkiewicz *et al.*), *Phys. Rev. C* **102**, 011901 (2020).
- [4] NA61/SHINE Collaboration (A. Acharya *et al.*), *Eur. Phys. J. C* **81**, 73 (2021).
- [5] M.P. Lewicki, [arXiv:2212.06539](https://arxiv.org/abs/2212.06539) [nucl-ex].
- [6] NA61/SHINE Collaboration (H. Adhikary *et al.*), CERN-SPSC-2022-034, 2022.
- [7] N.G. Antoniou, F.K. Diakonos, A.S. Kapoyannis, K.S. Kousouris, *Phys. Rev. Lett.* **97**, 032002 (2006).
- [8] A. Bialas, M. Gazdzicki, *Phys. Lett. B* **252**, 483 (1990).
- [9] T. Csörgő, S. Hegyi, W.A. Zajc, *Eur. Phys. J. C* **36**, 67 (2004).
- [10] NA61/SHINE Collaboration (A. Acharya *et al.*), *Eur. Phys. J. C* **82**, 322 (2022).
- [11] NA61/SHINE Collaboration (A. Acharya *et al.*), *Eur. Phys. J. C* **82**, 96 (2022).
- [12] NA61/SHINE Collaboration (A. Acharya *et al.*), *Eur. Phys. J. C* **81**, 911 (2021).