

## STRONG AND ELECTROMAGNETIC FORCES IN HEAVY ION COLLISIONS\*

MARIOLA KLUSEK-GAWENDA<sup>a</sup>, EWA KOZIK<sup>a</sup>, ANDRZEJ RYBICKI<sup>a</sup>  
IWONA SPUTOWSKA<sup>a</sup>, ANTONI SZCZUREK<sup>a,b</sup>

<sup>a</sup>The Henryk Niewodniczański Institute of Nuclear Physics  
Polish Academy of Sciences

Radzikowskiego 152, 31-342 Kraków, Poland

<sup>b</sup>University of Rzeszów, Rejtana 16, 35-959 Rzeszów, Poland

*(Received January 25, 2013)*

The interplay between the strong and electromagnetic force in high energy nucleus–nucleus collisions was studied experimentally and theoretically in our earlier works. This effect appeared to result in very large distortions in spectra of charged pions produced in the collision. It was also found to bring new, independent information on the space-time evolution of the non-perturbative process of particle production. In this paper, we present our new results on the influence of the spectator-induced electromagnetic force on spectra of charged particles produced in two different Pb-induced reactions. For the first time, we also address the topic of  $p + A$  collisions in view of obtaining information about their centrality and nuclear break-up, both subjects being of importance in the context of the new  $p + A$  data collected at the LHC.

DOI:10.5506/APhysPolBSupp.6.451

PACS numbers: 25.75.-q, 12.38.Mh

### 1. Introduction

This paper addresses a specific class of nucleus–nucleus collisions, where a large “spectator system” survives. Such specific reactions give the opportunity to investigate the interplay between phenomena occurring in the participant and spectator zones. In particular, this is the case for our study which concerns the interplay between the strong and electromagnetic forces in such reactions. This can be used as a new source of information on the collision dynamics.

---

\* Presented at the International Symposium on Multiparticle Dynamics, Kielce, Poland, September 17–21, 2012.

## 2. The data

All the studies summarized in this paper were inspired by experimental observations made in the SPS energy regime (beam energy of 158 GeV/nucleon,  $\sqrt{s_{NN}} = 17.3$  GeV). These were coming from two data sets, both collected by the NA49 experiment [4]. The first was a sample of peripheral Pb+Pb reactions, defined by a cut of 150–300 charged particles measured by the NA49 detector [1]. This corresponded to an average number of  $54 \pm 11$  participating nucleons and a mean impact parameter of  $10.9 \pm 0.5$  fm [2]. The second data set was a sample of “Pb+gas” events, *i.e.* collisions of the Pb beam projectiles with nuclei of gas surrounding the NA49 target [3]. These




Fig. 1. (Color online) Invariant density  $d = E \frac{d^3 n}{dp^3}$  of positive and negative pions produced in (a) peripheral Pb+Pb and (b) Pb+gas collisions, drawn as a function of pion transverse momentum at fixed values of  $x_F$  (listed from the top to the bottom distribution). Note: for clarity, the subsequent distributions are multiplied by (a) 1, 0.1, 0.01,  $10^{-3}$ ,  $10^{-4}$ ,  $2 \times 10^{-6}$ ,  $10^{-7}$ ,  $10^{-8}$ ,  $10^{-9}$ , and (b) 1, 0.1, 0.01,  $10^{-3}$ ,  $10^{-4}$ ,  $10^{-5}$ . The data come from (a) [1, 2] and (b) [3].

collisions (mostly Pb+N, Pb+O) were experimentally isolated by means of proper interaction vertex cuts, together with the same cut on measured multiplicity as given above [5]. It is, therefore, expected that the corresponding mean number of participating nucleons was roughly comparable to that in peripheral Pb+Pb reactions. The double differential spectra of positively and negatively charged pions produced in both reaction samples are shown in Fig. 1. They are drawn as a function of transverse momentum  $p_T$  at fixed values of the Feynman variable<sup>1</sup>  $x_F = \frac{2p_L}{\sqrt{s_{NN}}}$ .

### 3. The ratios $\frac{\pi^+}{\pi^-}$ in Pb+Pb and Pb+gas events

Figures 2(a) and 2(b) show the ratios of positively over negatively charged pions produced in peripheral Pb+Pb and Pb+gas collisions, respectively. The similarity of the evolution of the ratios with  $x_F$  and  $p_T$  in the two collision types is evident. In both cases, the  $\frac{\pi^+}{\pi^-}$  ratio displays a steep, rapidly varying structure, with a deep minimum at low transverse momenta where it comes below 0.2. Taking into account the comparable number of protons and neutrons participating in both reactions (40% over 60% for the Pb nucleus), such a low value of the  $\frac{\pi^+}{\pi^-}$  ratio breaks isospin symmetry. This demonstrates that the strong interaction cannot be the sole responsible for this effect. The position of the minimum ( $x_F \approx 0.15 = \frac{m_\pi}{m_N}$ ) corresponds to pions moving longitudinally with spectator velocity. Indeed, the effect is caused by the *electromagnetic interaction between charged pions and the spectator system*. The repulsion of positive pions results in the depopulation of the region  $x_F \approx 0.15$ , low  $p_T$ , while the attraction of negative pions causes their accumulation in the same region of phase space (see also Fig. 1).

In order to quantify this hypothesis, we constructed an intentionally simplified model of the peripheral Pb+Pb collision, taking the spectator-induced electromagnetic interaction into account [1, 6]. This is illustrated in Fig. 2(d). The two spectator systems are modeled as two Lorentz-contracted, uniformly charged spheres. Charged pions are emitted from a single point in space (the interaction point). The time of pion emission  $t_E$  is a free parameter of the model, defining the initial conditions for the electromagnetic interaction. Initial spectra of pions are assumed to be equal to these in the mixture of nucleon–nucleon ( $p+p$ ,  $n+p$ ,  $p+n$ ,  $n+n$ ) collisions. They are constructed on the basis of NA49  $p+p$  data [8]. Proper account is taken of isospin symmetry and of the proton/neutron ratio in the Pb nucleus (40%/60%); details of this procedure can be found in [9]. Charged pions are then numerically propagated in the electromagnetic field of the spectators. Relativistic effects such as retardation are taken into account.

<sup>1</sup> All the kinematical variables will always be defined in the collision c.m.s.




Fig. 2. (Color online) (a)  $\frac{\pi^+}{\pi^-}$  ratio in peripheral Pb+Pb reactions, (b) the same ratio in Pb+gas collisions, (c) model simulation as described in the text, (d) schematic illustration of the model. The data in panel (b) come from [3]. Other panels come from [7].

As it can be seen in Fig. 2(c), the model gives a fair description of the main features of the peripheral Pb+Pb data, Fig. 2(a). It is to be noted that the simulation presented here assumes a mixed set of initial conditions with 50% of events generated with  $t_E = 0.5$  fm/c and the remaining 50% generated with  $t_E = 1$  fm/c. This means that the pion formation site is placed “behind” the nearest spectator system as discussed in [2]. The only region where a more significant disagreement between data and model can be seen is  $x_F \approx 0.2$ , low  $p_T$ . This has been identified as resulting from the *fragmentation (break-up) of the spectator system*, and yielding, in fact, independent information on the space-time evolution of this process [10].

#### 4. $\frac{\pi^+}{\pi^-}$ ratios over full phase space

Taking into account the success of our model in describing the basic features of the experimental data, we decide that it may be applied with

some confidence to investigate the electromagnetic distortion of  $\frac{\pi^+}{\pi^-}$  ratios in the full range of  $x_F$ . The results of our simulations<sup>2</sup> are shown in Fig. 3. As apparent in Fig. 3(a), the two-dimensional electromagnetic distortion seen in the data in Section 3 is, in fact, part of a larger structure, consisting of two symmetric “holes” in the  $\frac{\pi^+}{\pi^-}$  ratio, accompanied by a possible rise at higher absolute  $x_F$ . It is interesting to note that the region of sensitivity to the electromagnetic effect extends, at high  $x_F$ , to relatively large transverse momenta, up to and above  $p_T = 1$  GeV/ $c$ .




Fig. 3. (Color online)  $\frac{\pi^+}{\pi^-}$  ratio in peripheral Pb+Pb reactions (a) and (b), intermediate  $p$ +Pb collisions (c) and central  $p$ +Pb collisions (d). Note that the simulation of Pb+Pb reactions extends further in  $p_T$  relative to our earlier work [6].

Comparison with Fig. 3(b) demonstrates the sensitivity of the spectator-induced electromagnetic distortion to the pion emission time  $t_E$ . This shows the important ability of the electromagnetic effect to constrain possible scenarios and to provide new information on the space-time evolution of the process of pion production, see also [1, 6, 7, 10].

Finally, the electromagnetic effect in intermediate and central  $p$ +Pb reactions is shown in Figs. 3(c) and 3(d). With the trivial exception of the “one-hole” structure imposed by the presence of a single spectator nucleus,

<sup>2</sup> Note: for simplicity, this version of our simulation neglects the differences between initial  $\pi^+$  and  $\pi^-$  spectra, resulting from the excess of neutrons in the Pb nucleus (these differences were taken into account in the previous section). This implies that any deviation of the  $\frac{\pi^+}{\pi^-}$  ratio from unity will solely be due to electromagnetic effects.

the similarity to peripheral Pb+Pb collisions, Fig. 3(a), is evident. Little dependence on pure geometry (centrality) of the  $p$ +Pb collision is apparent in the figure. This gives a relatively well defined situation for using the electromagnetic interaction as source of information on pion production and on its interplay with nuclear fragmentation, also in  $p + A$  collisions.

## 5. Conclusions and outlook

The spectator-induced electromagnetic effect discussed in this paper influences a number of observables in various types of nuclear collisions ( $p + A$ , Pb+gas, Pb+Pb, *etc.*). The well-defined nature of the electromagnetic interaction makes it a convenient tool to provide independent information on the space-time evolution of the reaction. However, only a few phenomena have been addressed here. Other examples can be quoted. Among these, the electromagnetic effect induced on kaon spectra [7], on azimuthal anisotropies (directed flow) [11], its possible presence in “ultra-peripheral” Pb+Pb collisions [12], and many others are presently under active investigation.

This work was supported by the Polish National Science Centre (on the basis of decision No. DEC-2011/03/B/ST2/02634).

## REFERENCES

- [1] A. Rybicki, *PoS EPS-HEP*, 031 (2009).
- [2] A. Rybicki, Habilitation Thesis, Report No. 2040/PH, The H. Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences, Kraków 2010, <http://www.ifj.edu.pl/publ/reports/2010/>
- [3] I. Sputowska, A. Rybicki, to appear in *Eur. Phys. J. Web. of Conf.*
- [4] S. Afanasev *et al.*, [NA49 Collab.], *Nucl. Instrum. Methods* **A430**, 210 (1999).
- [5] I. Sputowska, M. Sc. Thesis, AGH Univ. of Science and Technology, Kraków 2010. See also: A. Łászló, Ph.D. Thesis, KFKI Research Institute for Particle and Nuclear Physics, Budapest 2007.
- [6] A. Rybicki, A. Szczurek, *Phys. Rev.* **C75**, 054903 (2007).
- [7] A. Rybicki, A. Szczurek, E. Kozik, *Acta Phys. Pol. B Proc. Suppl.* **5**, 369 (2012).
- [8] C. Alt *et al.*, [NA49 Collab.], *Eur. Phys. J.* **C45**, 343 (2006).
- [9] O. Chvala [NA49 Collab.], *Eur. Phys. J.* **C33**, S615 (2004).
- [10] A. Rybicki, *Acta Phys. Pol. B* **42**, 867 (2011).
- [11] A. Rybicki, A. Szczurek, [arXiv:1303.7354](https://arxiv.org/abs/1303.7354) [nucl-th].
- [12] M. Klusek-Gawenda, A. Szczurek, *Phys. Rev.* **C82**, 014904 (2010).