

INFRARED BEHAVIOUR OF PROPAGATORS AND RUNNING COUPLING IN THE CONFORMAL WINDOW OF QCD*

MARKUS HOPFER, REINHARD ALKOFER

Institut für Physik, Karl-Franzens Universität Graz
Universitätsplatz 5, 8010 Graz, Austria

CHRISTIAN S. FISCHER

Institut für Theoretische Physik, Justus-Liebig Universität Giessen
Heinrich-Buff-Ring 16, 35392 Giessen, Germany

(Received May 13, 2014)

Using the Dyson–Schwinger approach we investigate Landau gauge QCD with a relatively large number of chiral quark flavours. A self-consistent treatment on the propagator level enables us to study unquenching effects via the quark loop diagram in the gluon equation. Above the critical number of fermion flavours, the non-perturbative running coupling develops a plateau over a wide momentum range. Correspondingly, the propagators follow a power law behaviour in this momentum range indicating conformal behaviour. Our value $N_f^{\text{crit}} = 4.5$ is strongly sensitive to the details of the quark–gluon vertex calling for more detailed investigations in future studies.

DOI:10.5506/APhysPolBSupp.7.613

PACS numbers: 11.15.–q, 11.30.Rd, 12.38.Aw

1. Introduction

Walking technicolor models have been introduced to overcome the phenomenological difficulties faced by the early technicolor formulations [1], see [2] for a recent review. These models exhibit an approximate scale invariance over a wide energy range as well as a proximity to an infrared fixed point, where the gauge coupling is slowly running, or walking. Asymptotically free gauge theories can be utilized to mimic these properties, where it is expected that QCD with a large number of chiral fermion flavours is a viable candidate.

* Presented at “Excited QCD 2014”, Bjelašnica Mountain, Sarajevo, Bosnia and Herzegovina, February 2–8, 2014.

By linking the Green functions of a quantum field theory, the Dyson–Schwinger framework offers an appropriate non-perturbative tool to explore a given theory over all energies ranging from the deep infrared to the perturbative regime. Since DSEs constitute an infinite set of coupled integral equations carefully chosen truncations have to be applied in order to treat the equations numerically. Therefore, a comparison with other non-perturbative methods is inevitable at some point in order to fine-tune the truncation and to minimize errors induced by it. Once an appropriate truncation scheme is established, the Dyson–Schwinger framework is a reliable and robust tool to explore the theory.

2. The system of coupled Dyson–Schwinger equations

The central object in the following investigation is the quark propagator DSE depicted in Fig. 1. Intimately connected is the dressed gluon propagator indicated by the wiggly line. By increasing the fermion flavours,



Fig. 1. The DSE for the quark propagator. All internal propagators are dressed. Coloured blobs denote full vertices.

back-coupling effects of quark degrees of freedom on the Yang–Mills sector become important and simple model descriptions of the gluon propagator without detailed knowledge of its flavour dependence will prove to be insufficient. In particular, a naive extrapolation of QCD results to larger flavour numbers as done in Ref. [3] seems to be questionable. Thus, a self-consistent incorporation of the corresponding gluon DSE becomes mandatory. In the following, we outline the coupled system of DSEs and refer to Ref. [4] for details on the self-consistent treatment.

2.1. The coupled system

The renormalized DSE for the quark propagator is given by¹

$$S^{-1}(p) = Z_2 S_0^{-1}(p) + g^2 Z_{1F} C_F \int \frac{d^4 q}{(2\pi)^4} \gamma^\mu S(q) \Gamma^\nu(q, p; k) D^{\mu\nu}(k). \quad (1)$$

Here, Z_2 and Z_{1F} are the renormalization constants for the quark wave function and the quark–gluon vertex, respectively. The colour trace yields a factor of $C_F = (N_c^2 - 1)/(2N_c)$ and the gluon momentum is defined via

¹ We follow the conventions and notation of Ref. [5].

$k_\mu = p_\mu - q_\mu$. The full quark propagator is given by $S^{-1}(p) = -i\not{p}A(p^2, \mu^2) + B(p^2, \mu^2)$, where the dressing functions A and B implicitly depend on the renormalization scale μ . The quark mass function is defined via $M(p^2) = B(p^2, \mu^2)/A(p^2, \mu^2)$ and is a renormalization scale independent quantity.

The gluon propagator $D^{\mu\nu}(p)$ is included self-consistently by solving the corresponding DSEs for the Yang–Mills system², where we employ the truncation scheme proposed in Refs. [5, 7]. Unquenching effects enter the gluon DSE via the quark-loop diagram as depicted in Fig. 2. One obtains a closed




Fig. 2. The truncated DSE for the gluon propagator.

system if the quark–gluon vertex and the three-gluon vertex is specified. As shown later in Sec. 3, the phase transition is quite insensitive to details of the three-gluon vertex and the main impact seems to come from different tensor structures immanent in the quark–gluon vertex. Although this object was at the focus of recent investigations [8], it is up to now still too ambitious to include it in a full self-consistent way due to its complicated multi-tensor structure. In order to proceed, we defer this desirable but also highly demanding task to future work and model the quark–gluon vertex according to Ref. [5]. The formal structure of the gluon DSE is given by

$$D_{\mu\nu}^{-1}(p) = Z_3 D_{0,\mu\nu}^{-1}(p) + \Pi_{\mu\nu}^{\text{YM}}(p) + \Pi_{\mu\nu}^{\text{quark}}(p),$$

where the gluon self-energy contribution stemming from the quark-loop reads

$$\Pi_{\mu\nu}^{\text{quark}}(p) = -g^2 \frac{N_f}{2} Z_{1F} \int \frac{d^4 q}{(2\pi)^4} \text{tr}_D [\gamma_\mu S(q) \Gamma_\nu(q, k; p) S(k)]. \quad (2)$$

We note that, in general, a truncated DSE system is plagued by spurious divergencies appearing in the kernels of the loop integrals. Based on a UV analysis, a safe way to remove these unwanted contributions is to modify the integral kernels by constructing appropriate compensation terms, *cf.* Refs. [5, 7]. For moderate flavour numbers the quark loop diagram is IR sub-leading. Hence, a direct modification of the corresponding integral kernels is possible. However, as soon as the system approaches N_f^{crit} , the quark loop becomes IR enhanced and shows similar IR scaling as the ghost loop. Hence, subtracting quadratic divergencies directly from the quark loop fails

² The numerical implementation is detailed in Refs. [4–6].

if one wants to probe the chiral phase transition. In Ref. [4] we give several complementary methods which are able to eliminate these artificial contributions in a safe way and which are also used throughout.

3. Results

We present results obtained from a self-consistent treatment of the DSE system on the propagator level. For the three-point functions models are employed, where we emphasize the important role of the quark–gluon vertex tensor structure. As shown in Fig. 3 (a), above $N_f^{\text{crit}} \approx 4.5$ dynamical mass is no longer generated and the systems enter a chirally symmetric phase. Increasing the effective quark–gluon interaction strength using models which include *only* the tree-level vertex structure γ^μ has virtually no impact on the location of the phase transition as shown in Fig. 3 (b). We furthermore note that different models for the gauge-boson vertex [9] tend to influence




Fig. 3. Upper panel: The infrared quark mass function $M(p^2)$ and the chiral condensate $\langle\bar{\psi}\psi\rangle$ for different flavours N_f . Lines are drawn to guide the eye. Right: Different quark–gluon vertex models which contain *only* the tree-level structure do not influence the phase transition. However, *additional* tensor structure increases N_f^{crit} . Lower panel: The gauge-boson vertex has minor impact on the location of the transition and even tends to decrease N_f^{crit} . Right: Results obtained with different bare quark masses. As expected, the phase transition gets washed out.

N_f^{crit} only slightly as detailed in Fig. 3 (c). On the other hand, including *additional* tensor structure in the quark–gluon vertex increases N_f^{crit} considerably. Thus, a detailed knowledge of the quark–gluon vertex is crucial in order to give reliable predictions for N_f^{crit} .

The phase transition manifests itself also in a drastic change of the propagators. In Fig. 4 we present results for the non-perturbative running coupling $\alpha(p^2) = \alpha(\mu^2)Z(p^2)G^2(p^2)$, the ghost dressing function $G(p^2)$, the gluon propagator $Z(p^2)/p^2$ and the inverse vector self-energy $A^{-1}(p^2)$ for different flavour values $N_f \in \{0, 4, 5\}$. By increasing N_f the coupling is lowered, where at $N_f \lesssim N_f^{\text{crit}}$ this lowering is significant and finally at N_f^{crit} a sudden drop occurs and a plateau is formed which develops over a wide momentum range. If N_f is further increased, the plateau is successively lowered. Thus, within the chirally symmetric phase, a scaling relation between the Yang–Mills propagators is established, *i.e.* these objects develop a power law behaviour in this momentum region as can be seen from Fig. 4 (b) and Fig. 4 (c). As shown in Fig. 4 (d), the quark wave-function renormalization is constant in this region as expected from an IR analysis.




Fig. 4. Results for the running coupling $\alpha(p^2)$, the ghost dressing function $G(p^2, \mu^2)$, the gluon propagator $Z(p^2, \mu^2)/p^2$ and the quark wave-function renormalization $A^{-1}(p^2, \mu^2)$. In the calculations we use a perturbative renormalization scale of $\mu^2 = 5 \times 10^4 \text{ GeV}^2$.

4. Conclusions

We presented results from an exploratory study of large N_f QCD using the Dyson–Schwinger framework in Landau gauge. A self-consistent treatment of the corresponding DSEs on the propagator level reveals a transition to a chirally symmetric phase for $N_f^{\text{crit}} \approx 4.5$. The non-perturbative running coupling develops a plateau in this regime, where, correspondingly, the propagators follow a power law indicating conformal behaviour. The critical fermion flavour number is sensitive to details of the quark–gluon vertex model, whereas the gauge-boson vertex seems to play a minor role. This emphasizes the need for a more complete calculation using a full quark–gluon vertex in upcoming studies.

We thank Markus Huber, Axel Maas, Valentin Mader, Francesco Sannino and Milan Vujanovic for valuable discussions. M.H. acknowledges support from the Doktoratskolleg “Hadrons in Vacuum, Nuclei and Stars” of the Austrian Science Fund, FWF DK W1203-N16.

REFERENCES

- [1] S. Weinberg, *Phys. Rev.* **D19**, 1277 (1979); L. Susskind, *Phys. Rev.* **D20**, 2619 (1979); S. Weinberg, *Phys. Rev.* **D13**, 974 (1976); S. Dimopoulos, L. Susskind, *Nucl. Phys.* **B155**, 237 (1979); E. Eichten, K.D. Lane, *Phys. Lett.* **B90**, 125 (1980); E. Farhi, L. Susskind, *Phys. Rep.* **74**, 277 (1981).
- [2] F. Sannino, *Acta Phys. Pol. B* **40**, 3533 (2009) [arXiv:0911.0931 [hep-ph]].
- [3] A. Bashir, A. Raya, J. Rodriguez-Quintero, *Phys. Rev.* **D88**, 054003 (2013) [arXiv:1302.5829 [hep-ph]].
- [4] M. Hopfer, C.S. Fischer, R. Alkofer, in preparation.
- [5] C.S. Fischer, R. Alkofer, *Phys. Rev.* **D67**, 094020 (2003) [arXiv:hep-ph/0301094]; C.S. Fischer, arXiv:hep-ph/0304233; C.S. Fischer, P. Watson, W. Cassing, *Phys. Rev.* **D72**, 094025 (2005) [arXiv:hep-ph/0509213].
- [6] M. Hopfer, R. Alkofer, G. Haase, *Comput. Phys. Commun.* **184**, 1183 (2013) [arXiv:1206.1779 [hep-ph]].
- [7] C.S. Fischer, R. Alkofer, H. Reinhardt, *Phys. Rev.* **D65**, 094008 (2002) [arXiv:hep-ph/0202195]; .
- [8] R. Williams, arXiv:1404.2545 [hep-ph]; M. Hopfer, A. Windisch, R. Alkofer, *PoS CONFINEMENT X*, 073 (2013) [arXiv:1301.3672 [hep-ph]]; R. Alkofer, C.S. Fischer, F.J. Llanes-Estrada, K. Schwenzer, *Ann. Phys.* **324**, 106 (2009) [arXiv:0804.3042 [hep-ph]].
- [9] M.Q. Huber, L. von Smekal, *J. High Energy Phys.* **04**, 149 (2013) [arXiv:1211.6092 [hep-th]]; *PoS CONFINEMENT X*, 062 (2013) [arXiv:1301.3080 [hep-th]].