

THE q -STATISTICS AND QCD THERMODYNAMICS AT LHC*

TRAMBAK BHATTACHARYYA, ARVIND KHUNTIA, PRAGATI SAHOO
PRAKHAR GARG, POOJA PAREEK, RAGHUNATH SAHOO[†]

Discipline of Physics, School of Basic Sciences, Indian Institute of Technology
Indore, Khandwa Road, Simrol, M.P.- 453552, India

JEAN CLEYMANS

UCT-CERN Research Centre and Department of Physics
University of Cape Town
Rondebosch 7701, South Africa

(Received May 4, 2016)

We perform a Taylor series expansion of Tsallis distribution by assuming the Tsallis parameter q close to 1. The q value shows the deviation of a system from a thermalised Boltzmann distribution. By taking up to first order in $(q - 1)$, we derive an analytical result for Tsallis distribution including radial flow. Further, in the present work, we also study the speed of sound (c_s) as a function of temperature using the non-extensive Tsallis statistics for different q values and for different mass cut-offs.

DOI:10.5506/APhysPolBSupp.9.177

1. Introduction

Tsallis distribution has been successful in explaining the transverse momentum spectra observed at the Relativistic Heavy Ion Collider (RHIC) and at the Large Hadron Collider (LHC) with a non-extensive parameter q . Here, q shows the deviation of a system from a thermalised Boltzmann system. We incorporate the radial flow in the Tsallis statistics by using Taylor series expansion to get an analytical expression. It is believed that c_s^2 in the hadronic medium decreases to zero in the first order phase transition from hadronic phase to QGP phase. Therefore, the study of c_s^2 is important to locate the QCD phase boundary.

* Presented by R. Sahoo at WPCF 2015: XI Workshop on Particle Correlations and Femtoscopy, Warszawa, Poland, November 3–7, 2015.

[†] Corresponding author: Raghunath.Sahoo@cern.ch

2. Taylor expansion of Tsallis distribution in $(q - 1)$

Tsallis non-extensive distribution function is given by

$$f = \left[1 + (q - 1) \frac{E - \mu}{T} \right]^{-\frac{1}{q-1}}. \quad (1)$$

One obtains the relevant thermodynamical quantities by using f^q [1]. Assuming the non-extensive parameter q close to 1, Tsallis distribution function, f^q , can be expanded in Taylor series in $(q - 1)$, which is given by

$$\begin{aligned} & \left[1 + (q - 1) \frac{E - \mu}{T} \right]^{-\frac{q}{q-1}} \\ & \simeq e^{-\frac{E-\mu}{T}} \left\{ 1 + (q - 1) \frac{1}{2} \frac{E - \mu}{T} \left(-2 + \frac{E - \mu}{T} \right) + \frac{(q - 1)^2}{2!} \right. \\ & \times \frac{1}{12} \left[\frac{E - \mu}{T} \right]^2 \left[24 - 20 \frac{E - \mu}{T} + 3 \left(\frac{E - \mu}{T} \right)^2 \right] + \mathcal{O} \{ (q - 1)^3 \} + \left. \right\} \quad (2) \end{aligned}$$

This result is used to calculate the momentum spectra [1]. In Fig. 1 (left), we show fits to the normalized differential π^- yields in (0–5)% Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the Tsallis (solid line) and Boltzmann distributions (dashed line). Also, fits with the Tsallis distribution keeping terms to first order (dash-dotted line) and second order in $(q - 1)$ (dotted line) are shown in Fig. 1 (left). In Fig. 1 (right), the fits are with the Tsallis distribution including flow keeping terms to first order in $(q - 1)$ (dashed line). The flow velocity is fixed at $v = 0.609$, with $T = 146$ MeV, $q = 1.030$




Fig. 1. Fits to the normalized differential π^- yields in (0–5)% Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV [1] with the Tsallis distribution without (left) and with (right) radial flow (dashed line).

and the radius of the volume is $R = 29.8$ fm [1]. The lower parts of the figures show the deviation from data points. For truncation of the series expansion up to the first order with pure Tsallis distribution, one needs two conditions to be satisfied [1], *i.e.* $|1 - q|\frac{E}{T} < 1$ and $q|1 - q|\left(\frac{E}{T}\right)^2 < 2$.

3. Momentum spectra including radial flow in $(q - 1)$

In order to incorporate the effect of radial flow on the momentum spectra and to see how it could improve to explain the momentum spectra obtained in Pb+Pb collisions, we have included a constant flow velocity, v , and the invariant yield is given by

$$\begin{aligned} \frac{1}{p_T} \frac{dN}{dp_T dy} = \frac{gV}{(2\pi)^2} & \left\{ 2T[rI_0(s)K_1(r) - sI_1(s)K_0(r)] \right. \\ & - (q-1)Tr^2I_0(s)[K_0(r) + K_2(r)] + 4(q-1)TrsI_1(s)K_1(r) \\ & - (q-1)Ts^2K_0(r)[I_0(s) + I_2(s)] + \frac{(q-1)}{4}Tr^3I_0(s)[K_3(r) + 3K_1(r)] \\ & - \frac{3(q-1)}{2}Tr^2s[K_2(r) + K_0(r)]I_1(s) + \frac{3(q-1)}{2}Ts^2r[I_0(s) \\ & \left. + I_2(s)]K_1(r) - \frac{(q-1)}{4}Ts^3[I_3(s) + 3I_1(s)]K_0(r) \right\}. \end{aligned} \quad (3)$$

$r \equiv \frac{\gamma m_T}{T}$, $s \equiv \frac{\gamma v p_T}{T}$, $I_n(s)$ and $K_n(r)$ are the modified Bessel functions of the first and second kind.

Figure 1 (left) shows that our model explains experimental data very well in the intermediate p_T region (0.5–2.5 GeV/c) but shows slight deviation in low and high p_T regions.

4. Speed of sound in a physical hadron resonance gas

For a physical hadron resonance gas, we use Tsallis form of Fermi–Dirac (FD) and Bose–Einstein (BE) statistics according to particle species [2], which is given by

$$f_T(E) \equiv \frac{1}{\exp_q \left(\frac{E - \mu}{T} \right) \pm 1}. \quad (4)$$

The \pm sign in the denominator stands for FD and BE respectively. In the limit $q \rightarrow 1$, \exp_q reduces to the standard exponential. The square of speed of sound is the change in pressure with change in energy density, which is given by

$$c_s^2(T) = \left(\frac{\partial P}{\partial \epsilon} \right)_V = \frac{s(T)}{C_V(T)}, \quad (5)$$

where, s is the entropy and C_V is the specific heat at constant volume. c_s^2 is plotted as a function of temperature in Fig. 2 (left) and Fig. 2 (right), taking different mass cut-offs for $q = 1.05$ and $q = 1.15$.




Fig. 2. The c_s^2 as a function of temperature for $q = 1.05$ (left) [2] and $q = 1.15$ (right) [2].

5. Summary

Taylor series expansion of Tsallis distribution in $(q - 1)$ is used to introduce the radial flow in an analytic way in addition to the study of the degree of deviation from equilibrium Boltzmann distribution. The p_T spectra for 0–5% central Pb+Pb collisions at 2.76 TeV is well-described with inclusion of radial flow up to first order in $(q - 1)$. The speed of sound for physical resonance gas has been calculated for different non-extensive parameters q . Further, taking higher q values, it seems to increase the speed of sound near the limiting temperature (T_H) as compared to the extensive Boltzmann statistics. For higher q values, the critical behaviour of speed of sound occurs earlier as compared to extensive statistics, which indicates an earlier phase transition in non-extensive statistics. This seems to be a function of degree of deviation from equilibrium statistics.

REFERENCES

- [1] T. Bhattacharyya *et al.*, *Eur. Phys. J. A* **52**, 30 (2016) and references therein.
- [2] A. Khuntia *et al.*, [arXiv:1602.01645 \[hep-ph\]](#) and references therein.