

WAVES ALONG FRACTAL COASTLINES: FROM FRACTAL ARITHMETIC TO WAVE EQUATIONS

MAREK CZACHOR

Department of Theoretical Physics and Quantum Information
Gdańsk University of Technology
G. Narutowicza 11/12, 80-233 Gdańsk, Poland

(Received December 12, 2018; accepted March 13, 2019)

Beginning with addition and multiplication intrinsic to a Koch-type curve, we formulate and solve wave equation describing wave propagation along a fractal coastline. As opposed to examples known from the literature, we do not replace the fractal by the continuum in which it is embedded. This seems to be the first example of a truly intrinsic description of wave propagation along a fractal curve. The theory is relativistically covariant under an appropriately defined Lorentz group.

DOI:10.5506/APhysPolB.50.813

1. Non-Newtonian calculus

Consider two sets \mathbb{X} and \mathbb{Y} whose cardinality is continuum, and a function $A : \mathbb{X} \rightarrow \mathbb{Y}$. There exist bijections $f_{\mathbb{X}}, f_{\mathbb{Y}}, g_{\mathbb{X}}, g_{\mathbb{Y}}$, such that the diagram

$$\begin{array}{ccccc}
 \mathbb{R} & \xrightarrow{\tilde{B}} & \mathbb{R} \\
 g_{\mathbb{X}} \uparrow & & \uparrow g_{\mathbb{Y}} \\
 \mathbb{X} & \xrightarrow{A} & \mathbb{Y} \\
 f_{\mathbb{X}} \downarrow & & \downarrow f_{\mathbb{Y}} \\
 \mathbb{R} & \xrightarrow{\tilde{A}} & \mathbb{R}
 \end{array}$$

is commutative. The functions \tilde{A} and \tilde{B} are defined by the diagram. It is natural to think of \mathbb{X} and \mathbb{Y} in terms of one-dimensional manifolds whose global charts are defined by the bijections.

In differential topology and geometry, a derivative of $A : \mathbb{X} \rightarrow \mathbb{Y}$ would be a function $A' : \mathbb{X} \rightarrow \mathbb{Y}$ defined by $\tilde{A}'(r) = d\tilde{A}(r)/dr$. Of course, since

$$\tilde{A} = f_{\mathbb{Y}} \circ g_{\mathbb{Y}}^{-1} \circ \tilde{B} \circ g_{\mathbb{X}} \circ f_{\mathbb{X}}^{-1} = \varphi_{\mathbb{Y}}^{-1} \circ \tilde{B} \circ \varphi_{\mathbb{X}},$$

a derivative of A can be equivalently defined in terms of \tilde{B} , provided $\varphi_{\mathbb{X}}$ and $\varphi_{\mathbb{Y}}^{-1}$ are at least C^1 . A transition between the two forms is determined by the chain rule for derivatives.

In the arithmetic approach to differentiation [1–9], one starts from a different perspective. In the first step, one employs the bijections to turn \mathbb{X} and \mathbb{Y} into fields isomorphic to \mathbb{R} . Explicitly, one defines the arithmetic operations in \mathbb{X} (addition, subtraction, multiplication, division) by

$$x \oplus_{\mathbb{X}} y = f_{\mathbb{X}}^{-1}(f_{\mathbb{X}}(x) + f_{\mathbb{X}}(y)), \quad (1)$$

$$x \ominus_{\mathbb{X}} y = f_{\mathbb{X}}^{-1}(f_{\mathbb{X}}(x) - f_{\mathbb{X}}(y)), \quad (2)$$

$$x \odot_{\mathbb{X}} y = f_{\mathbb{X}}^{-1}(f_{\mathbb{X}}(x)f_{\mathbb{X}}(y)), \quad (3)$$

$$x \oslash_{\mathbb{X}} y = f_{\mathbb{X}}^{-1}(f_{\mathbb{X}}(x)/f_{\mathbb{X}}(y)), \quad (4)$$

and analogously in \mathbb{Y} . This type of arithmetic is a special case of a general non-Diophantine arithmetic discussed by Burgin [10–13]. The case of a linear f was extensively studied in [14–16] with emphasis on distinguishing between numbers, treated abstractly, and their representations and values.

The topologies of \mathbb{X} and \mathbb{Y} are induced by the bijections from the topology of \mathbb{R} . Let the limit $x \rightarrow x_0 \in \mathbb{X}$ be defined by the formula

$$\lim_{x \rightarrow x_0} A(x) = f_{\mathbb{Y}}^{-1} \left(\lim_{r \rightarrow f_{\mathbb{X}}(x_0)} \tilde{A}(r) \right). \quad (5)$$

The derivative of A can be expressed in terms of limits in three equivalent ways

$$\begin{aligned} \frac{DA(x)}{Dx} &= \lim_{h \rightarrow 0} \left(A(x \oplus_{\mathbb{X}} f_{\mathbb{X}}^{-1}(h)) \ominus_{\mathbb{Y}} A(x) \right) \oslash_{\mathbb{Y}} f_{\mathbb{Y}}^{-1}(h) \\ &= \lim_{h \rightarrow 0} \left(A(x \oplus_{\mathbb{X}} h_{\mathbb{X}}) \ominus_{\mathbb{Y}} A(x) \right) \oslash_{\mathbb{Y}} h_{\mathbb{Y}} \\ &= \lim_{h \rightarrow 0_{\mathbb{X}}} \left(A(x \oplus_{\mathbb{X}} h) \ominus_{\mathbb{Y}} A(x) \right) \oslash_{\mathbb{Y}} f(h), \end{aligned} \quad (6)$$

where $f = f_{\mathbb{Y}}^{-1} \circ f_{\mathbb{X}}$. Here, $0_{\mathbb{X}}$ is the neutral element of addition in \mathbb{X} . This type of derivative was investigated in a systematic way for the first time in [1], for the case where \mathbb{X} and \mathbb{Y} were subsets of \mathbb{R} , while $f_{\mathbb{Y}}$ and $f_{\mathbb{X}}$ were continuous in the metric topology of \mathbb{R} . The derivative was rediscovered by myself in a fractal context [4]. The main difference between the formalism from [1] and my approach is that now the derivative is applicable to all sets whose cardinality is continuum, such as Sierpiński-type fractals [7], which obviously do not have to be subsets of \mathbb{R} , and in typical examples, the bijections are discontinuous in metric topologies of \mathbb{X} and \mathbb{Y} . This counterintuitive possibility opened by the non-Newtonian calculus is especially useful in fractal

applications. Just to give one example, a construction of Fourier transforms on arbitrary Cantor sets is in the non-Newtonian framework basically trivial [6], simultaneously circumventing various impossibility theorems known from the more traditional approach [17, 18]. The arithmetic perspective is simultaneously applicable to all the other aspects of mathematical modeling, including algebraic or probabilistic methods. The freedom of choice of arithmetic plays a role of a universal symmetry of any mathematical model.

An equivalent and very convenient form of the derivative is

$$\frac{DA(x)}{Dx} = f_{\mathbb{Y}}^{-1} \left(\frac{d\tilde{A}(f_{\mathbb{X}}(x))}{df_{\mathbb{X}}(x)} \right). \quad (7)$$

The derivative is Newtonian if \mathbb{X} and \mathbb{Y} are subsets of \mathbb{R} , and $f_{\mathbb{X}}(x) = x$, $f_{\mathbb{Y}}(y) = y$ are the identity maps. If the bijections are less trivial, one speaks of non-Newtonian derivatives.

Of particular interest is the non-Newtonian version of the chain rule. Consider the diagram

$$\begin{array}{ccccc} \mathbb{X} & \xrightarrow{A} & \mathbb{Y} & \xrightarrow{B} & \mathbb{Z} \\ f_{\mathbb{X}} \downarrow & & f_{\mathbb{Y}} \downarrow & & f_{\mathbb{Z}} \downarrow \\ \mathbb{R} & \xrightarrow{\tilde{A}} & \mathbb{R} & \xrightarrow{\tilde{B}} & \mathbb{R} \end{array},$$

then

$$\frac{D(B \circ A)(x)}{Dx} = f_{\mathbb{Z}}^{-1} \left[f_{\mathbb{Z}} \left(\frac{DB(A(x))}{DA(x)} \right) f_{\mathbb{Y}} \left(\frac{DA(x)}{Dx} \right) \right]. \quad (8)$$

For a composition of three functions,

$$\mathbb{W} \xrightarrow{A} \mathbb{X} \xrightarrow{B} \mathbb{Y} \xrightarrow{C} \mathbb{Z}, \quad (9)$$

one finds

$$\begin{aligned} & \frac{DC \circ B \circ A(x)}{Dx} = \\ & f_{\mathbb{Z}}^{-1} \left[f_{\mathbb{Z}} \left(\frac{DC[B(A(x))]}{DB(A(x))} \right) f_{\mathbb{Y}} \left(\frac{DB(A(x))}{DA(x)} \right) f_{\mathbb{X}} \left(\frac{DA(x)}{Dx} \right) \right]. \end{aligned} \quad (10)$$

The latter case is important since it allows us to better understand the structure of the non-Newtonian derivative. Indeed, let the three functions be the ones occurring in the definition of $\mathbb{X} \xrightarrow{A} \mathbb{Y}$, *i.e.*

$$\mathbb{X} \xrightarrow{f_{\mathbb{X}}} \mathbb{R} \xrightarrow{\tilde{A}} \mathbb{R} \xrightarrow{f_{\mathbb{Y}}^{-1}} \mathbb{Y}. \quad (11)$$

Now, directly from the definition, one checks that

$$\frac{Df_{\mathbb{X}}(x)}{Dx} = 1, \quad \frac{Df_{\mathbb{Y}}(x)}{Dx} = 1, \quad \frac{Df_{\mathbb{X}}^{-1}(x)}{Dx} = 1_{\mathbb{X}}, \quad \frac{Df_{\mathbb{Y}}^{-1}(x)}{Dx} = 1_{\mathbb{Y}}. \quad (12)$$

The chain rule implies

$$\begin{aligned} \frac{DA(x)}{Dx} &= \frac{D\left(f_{\mathbb{Y}}^{-1} \circ \tilde{A} \circ f_{\mathbb{X}}\right)(x)}{Dx} \\ &= f_{\mathbb{Y}}^{-1} \left[f_{\mathbb{Y}} \left(\frac{Df_{\mathbb{Y}}^{-1} \left[\tilde{A}(f_{\mathbb{X}}(x)) \right]}{D\tilde{A}(f_{\mathbb{X}}(x))} \right) f_{\mathbb{R}} \left(\frac{D\tilde{A}(f_{\mathbb{X}}(x))}{Df_{\mathbb{X}}(x)} \right) f_{\mathbb{R}} \left(\frac{Df_{\mathbb{X}}(x)}{Dx} \right) \right]. \end{aligned}$$

The arithmetic in \mathbb{R} is Diophantine, $f_{\mathbb{R}}(x) = x$, and thus

$$\frac{D\tilde{A}(f_{\mathbb{X}}(x))}{Df_{\mathbb{X}}(x)} = \frac{d\tilde{A}(f_{\mathbb{X}}(x))}{df_{\mathbb{X}}(x)}$$

is Newtonian. Derivatives (12) imply

$$\frac{DA(x)}{Dx} = f_{\mathbb{Y}}^{-1} \left(\frac{d\tilde{A}(f_{\mathbb{X}}(x))}{df_{\mathbb{X}}(x)} \right),$$

and we reconstruct our definition of the derivative. One concludes that the bijections behave as identity maps with respect to non-Newtonian derivatives they define, no matter how weird the bijections themselves actually are.

The integral is defined in a way guaranteeing the fundamental laws of calculus, relating derivatives and integrals

$$\int_Y^X A(x) Dx = f_{\mathbb{Y}}^{-1} \left(\int_{f_{\mathbb{X}}(Y)}^{f_{\mathbb{X}}(X)} \tilde{A}(x) dx \right), \quad (13)$$

where $\int \tilde{A}(x) dx$ is the usual (say, Lebesgue) integral of a function $\tilde{A} : \mathbb{R} \rightarrow \mathbb{R}$. One proves that

$$\frac{D}{DX} \int_Y^X A(x) Dx = A(X), \quad (14)$$

$$\int_Y^X \frac{DA(x)}{Dx} Dx = A(X) \ominus_{\mathbb{Y}} A(Y). \quad (15)$$

Let us now see how it works in the simple but instructive case of $f(x) = x^3$. The manifold in question is $\mathbb{X} = \mathbb{R}$. Let the two (global) charts be given by $f(x) = x^3$ and $g(x) = x$. Their composition $g \circ f^{-1}$ is not a diffeomorphism if the differentiation is understood in the Newtonian way. Apparently, $f(x) = x^3$ does not define a differentiable structure on \mathbb{R} . In the standard Newtonian formalism, the only structure we have at our disposal is C^0 .

The arithmetic approach begins with arithmetic operations intrinsic to \mathbb{X}

$$x \oplus y = f^{-1}(f(x) + f(y)) = \sqrt[3]{x^3 + y^3}, \quad (16)$$

$$x \ominus y = f^{-1}(f(x) - f(y)) = \sqrt[3]{x^3 - y^3}, \quad (17)$$

$$x \odot y = f^{-1}(f(x)f(y)) = \sqrt[3]{x^3 y^3} = xy, \quad (18)$$

$$x \oslash y = f^{-1}(f(x)/f(y)) = \sqrt[3]{x^3/y^3} = x/y. \quad (19)$$

Let us stress again that f is, by construction, a field isomorphism of $(\mathbb{R}, +, \cdot)$ and $(\mathbb{R}, \oplus, \odot)$. Therefore, \oplus and \odot are commutative and associative, and \odot is distributive with respect to \oplus . The neutral elements of \oplus and \odot , $0'$ and $1'$, are the standard ones: $0' = f^{-1}(0) = \sqrt[3]{0} = 0$, $1' = f^{-1}(1) = \sqrt[3]{1} = 1$. Although multiplication is unchanged, the link between addition and multiplication is a subtle one, as can be seen in the following example:

$$\begin{aligned} x \oplus \dots \oplus x &= \sqrt[3]{x^3 + \dots + x^3} \quad (n \text{ times}) \\ &= \sqrt[3]{nx} = f^{-1}(n)x. \end{aligned} \quad (20)$$

The inverse bijection $f^{-1}(x) = \sqrt[3]{x}$ is continuous but not Newtonian differentiable at $x = 0$, hence the loss of the Newtonian diffeomorphism property. Still, the derivative of a function $A : \mathbb{X} \rightarrow \mathbb{X}$,

$$\frac{DA(x)}{Dx} = \lim_{h \rightarrow 0} \left(A(x \oplus h) \ominus A(x) \right) \oslash h, \quad (21)$$

is well defined. The non-Newtonian D/Dx satisfies all the basic rules of differentiation, of course with respect to the new arithmetic:

(a) The Leibniz rule.

$$\begin{aligned} \frac{DA(x)B(x)}{Dx} &= \lim_{h \rightarrow 0} \left(A(x \oplus h)B(x \oplus h) \ominus A(x)B(x) \right) \oslash h \\ &= \sqrt[3]{\left(\frac{DA(x)}{Dx}B(x) \right)^3 + \left(A(x)\frac{DB(x)}{Dx} \right)^3} \\ &= \frac{DA(x)}{Dx}B(x) \oplus A(x)\frac{DB(x)}{Dx}. \end{aligned} \quad (22)$$

(b) Linearity.

$$\begin{aligned}
 \frac{DA(x) \oplus B(x)}{Dx} &= \lim_{h \rightarrow 0} \left(A(x \oplus h) \oplus B(x \oplus h) \ominus (A(x) \oplus B(x)) \right) \oslash h \\
 &= \sqrt[3]{\left(\frac{DA(x)}{Dx} \right)^3 + \left(\frac{DB(x)}{Dx} \right)^3} \\
 &= \frac{DA(x)}{Dx} \oplus \frac{DB(x)}{Dx}.
 \end{aligned} \tag{23}$$

(c) The chain rule. Denoting

$$H = B(x \oplus h) \ominus B(x), \tag{24}$$

we obtain

$$\begin{aligned}
 \frac{DA(B(x))}{Dx} &= \lim_{h \rightarrow 0} \sqrt[3]{\frac{\left(A(B(x) \oplus H) \right)^3 - \left(A(B(x)) \right)^3}{h^3}} \\
 &= \lim_{H \rightarrow 0} \frac{A(B(x) \oplus H) \ominus A(B(x))}{H} \lim_{h \rightarrow 0} \frac{B(x \oplus h) \ominus B(x)}{h} \\
 &= \frac{DA(B(x))}{DB(x)} \frac{DB(x)}{Dx}.
 \end{aligned} \tag{25}$$

The non-Newtonian derivate has interesting implications for differential equations. For example, the unique solution of

$$\frac{DA(x)}{Dx} = A(x), \quad A(0) = 1 \tag{26}$$

is

$$A(x) = e^{x^3/3} = f^{-1} \left(e^{f(x)} \right), \tag{27}$$

as one can verify directly from definition (21). The exponent satisfies the usual law

$$A(x_1 \oplus x_2) = e^{(x_1^3 + x_2^3)/3} = A(x_1) \odot A(x_2). \tag{28}$$

One can similarly verify that

$$\text{Sin } x = \sqrt[3]{\sin(x^3)}, \tag{29}$$

$$\text{Cos } x = \sqrt[3]{\cos(x^3)} \tag{30}$$

satisfy

$$\frac{D\text{Sin } x}{Dx} = \text{Cos } x, \quad (31)$$

$$\frac{D\text{Cos } x}{Dx} = \ominus\text{Sin } x = -\text{Sin } x, \quad (32)$$

where $\ominus x = 0 \ominus x = \sqrt[3]{-x^3} = -x$, and

$$\text{Sin}^2 x \oplus \text{Cos}^2 x = \sqrt[3]{\sin^2(x^3) + \cos^2(x^3)} = 1. \quad (33)$$

$\text{Sin } x$ and $\text{Cos } x$ are essentially the chirp signals known from signal analysis (Fig. 1 and Fig. 2).




Fig. 1. The circle $x \mapsto (\text{Cos } x, \text{Sin } x)$, $0 \leq x < (2\pi)^{1/3}$, with trigonometric functions given by (29)–(30).

It is instructive to compare (31) with the Newtonian derivative

$$\frac{d\text{Sin } x}{dx} = \frac{x^2 \cos(x^3)}{\sin^{\frac{2}{3}}(x^3)}, \quad (34)$$

defined with respect to the ‘standard’ arithmetic (Fig. 2).




Fig. 2. The non-Newtonian derivative $\text{DSin } x/Dx = \text{Cos } x$ (full, Eq. (31)), as compared to the standard Newtonian $\text{dSin } x/dx$ (dashed, Eq. (34)). The singular behavior of the dashed curve follows from Newtonian non-differentiability of $f^{-1}(x) = \sqrt[3]{x}$ at $x = 0$. In contrast, the non-Newtonian derivative is non-singular since f and f^{-1} get differentiated in a non-Newtonian way, yielding trivial derivatives.

Even more intriguing examples occur if one considers derivatives of functions $A : \mathbb{X} \rightarrow \mathbb{Y}$ where the domain and the image of A involve different arithmetics. Let $\mathbb{X} = \mathbb{R}_+$, $\mathbb{Y} = \mathbb{R}$, $f_{\mathbb{X}}(x) = \ln x$, $f_{\mathbb{Y}}(x) = x^3$. The arithmetic operations in \mathbb{X} read explicitly

$$x_1 \oplus_{\mathbb{X}} x_2 = f_{\mathbb{X}}^{-1}(f_{\mathbb{X}}(x_1) + f_{\mathbb{X}}(x_2)) = e^{\ln x_1 + \ln x_2} = x_1 x_2, \quad (35)$$

$$x_1 \ominus_{\mathbb{X}} x_2 = f_{\mathbb{X}}^{-1}(f_{\mathbb{X}}(x_1) - f_{\mathbb{X}}(x_2)) = e^{\ln x_1 - \ln x_2} = x_1/x_2, \quad (36)$$

$$x_1 \odot_{\mathbb{X}} x_2 = f_{\mathbb{X}}^{-1}(f_{\mathbb{X}}(x_1)f_{\mathbb{X}}(x_2)) = e^{\ln x_1 \ln x_2} = x_1^{\ln x_2} = x_2^{\ln x_1}, \quad (37)$$

$$x_1 \oslash_{\mathbb{X}} x_2 = f_{\mathbb{X}}^{-1}(f_{\mathbb{X}}(x_1)/f_{\mathbb{X}}(x_2)) = e^{\ln x_1 / \ln x_2} = x_1^{1/\ln x_2}. \quad (38)$$

Neutral elements in \mathbb{X} are given by

$$1_{\mathbb{X}} = f_{\mathbb{X}}^{-1}(1) = e^1 = e, \quad (39)$$

$$0_{\mathbb{X}} = f_{\mathbb{X}}^{-1}(0) = e^0 = 1. \quad (40)$$

A negative of $x \in \mathbb{X}$ is defined as

$$\ominus_{\mathbb{X}} x = 0_{\mathbb{X}} \ominus_{\mathbb{X}} x = f_{\mathbb{X}}^{-1}(-f_{\mathbb{X}}(x)) = e^{-\ln x} = 1/x \in \mathbb{R}_+. \quad (41)$$

As we can see, numbers negative with respect to the arithmetic from \mathbb{X} are positive if treated in the usual sense. The unique solution $A : \mathbb{X} \rightarrow \mathbb{Y}$ of

$$\frac{\text{DA}(x)}{\text{D}x} = A(x), \quad A(0_{\mathbb{X}}) = 1_{\mathbb{Y}} \quad (42)$$

turns out to be

$$A(x) = f_{\mathbb{Y}}^{-1} \left(e^{f_{\mathbb{X}}(x)} \right) = \sqrt[3]{e^{\ln x}} = \sqrt[3]{x}. \quad (43)$$

Indeed, first of all,

$$A(0_{\mathbb{X}}) = \sqrt[3]{1} = 1 = 1_{\mathbb{Y}}. \quad (44)$$

Recalling that multiplication in \mathbb{Y} is unchanged, we check directly from definition (cf. [7]):

$$\begin{aligned} \frac{DA(x)}{Dx} &= \lim_{h \rightarrow 0} \left(A(x \oplus_{\mathbb{X}} f_{\mathbb{X}}^{-1}(h)) \ominus_{\mathbb{Y}} A(x) \right) \oslash_{\mathbb{Y}} f_{\mathbb{Y}}^{-1}(h) \\ &= \lim_{h \rightarrow 0} \left(\sqrt[3]{x \oplus_{\mathbb{X}} e^h} \ominus_{\mathbb{Y}} \sqrt[3]{x} \right) / \sqrt[3]{h} \\ &= \lim_{h \rightarrow 0} \sqrt[3]{x \frac{e^h - 1}{h}} = \sqrt[3]{x} = A(x). \end{aligned} \quad (45)$$

The exponent satisfies

$$\begin{aligned} A(x_1 \oplus_{\mathbb{X}} x_2) &= A(x_1 x_2) = \sqrt[3]{x_1 x_2} = \sqrt[3]{x_1} \sqrt[3]{x_2} = A(x_1) A(x_2) \\ &= A(x_1) \odot_{\mathbb{Y}} A(x_2), \end{aligned} \quad (46)$$

as expected. The results are counterintuitive but consistent. The bijection $f_{\mathbb{X}}(x) = \ln x$ is a simplest example of an information channel associated with human or animal nervous system (the Weber–Fechner law; this is why decibels correspond to a logarithmic scale [19]).

As final two examples consider first $f_{\mathbb{X}}(x) = x$, $f_{\mathbb{Y}}(x) = \ln x$. The non-Newtonian derivative reads explicitly

$$\begin{aligned} \frac{DA(x)}{Dx} &= \lim_{h \rightarrow 0} \left(A(x+h) \ominus_{\mathbb{Y}} A(x) \right) \oslash_{\mathbb{Y}} h_{\mathbb{Y}} \\ &= \lim_{h \rightarrow 0} e^{(\ln A(x+h) - \ln A(x))/h} = e^{A'(x)/A(x)}. \end{aligned} \quad (47)$$

Here, $A'(x) = dA(x)/dx$ is the Newtonian derivative. Let us now solve

$$\frac{DA(x)}{Dx} = A(x), \quad A(0) = 1_{\mathbb{Y}} = f_{\mathbb{Y}}^{-1}(1) = e, \quad (48)$$

an equation equivalent to

$$e^{A'(x)/A(x)} = A(x). \quad (49)$$

By the general formula, we know that this must be the non-Newtonian exponent

$$A(x) = f_{\mathbb{Y}}^{-1} \left(e^{f_{\mathbb{X}}(x)} \right) = e^{e^x}. \quad (50)$$

Secondly, let $f_{\mathbb{X}}(x) = \ln x = f_{\mathbb{Y}}(x)$. Then

$$\frac{DA(x)}{Dx} = e^{xA'(x)/A(x)}. \quad (51)$$

Here, values of non-Newtonian and Newtonian exponents coincide,

$$A(x) = f_{\mathbb{Y}}^{-1} \left(e^{f_{\mathbb{X}}(x)} \right) = e^{e^{\ln x}} = e^x, \quad (52)$$

but their domains are different. Both types of differentiation have been extensively studied in the literature, with numerous applications [20–25]. The variety of applications, from signal processing to economics, is not that surprising if one realizes that $\ln x$ represents a neuronal information channel [19]. The two non-Newtonian derivatives represent here a perception of change, and not the change itself.

Armed with these intuitions, we are ready to apply the formalism to waves on Koch-type fractals.

2. Koch curve supported on unit interval

For convenience, we represent \mathbb{R}^2 by \mathbb{C} . Let us begin with the Koch curve $K_{[0,1]} \subset \mathbb{C}$, beginning at 0 and ending at 1 (Fig. 3).

A point $z \in K_{[0,1]}$ can be parametrized by a real number in quaternary representation

$$y = (0.q_1 \dots q_j \dots)_4 \in [0, 1], \quad (53)$$

where $q_k = 0, 1, 2, 3$. The parametrization is defined by a bijection $g : [0, 1] \rightarrow K_{[0,1]}$, $z = g(y)$, constructed as follows. Consider $a = e^{i\alpha}$, $0 \leq \alpha \leq \pi/2$, $L = 1/(2 + 2 \cos \alpha)$, and

$$\hat{0}(z) = Lz, \quad (54)$$

$$\hat{1}(z) = L(1 + az), \quad (55)$$

$$\hat{2}(z) = L(1 + a + \bar{a}z), \quad (56)$$

$$\hat{3}(z) = L(1 + 2 \cos \alpha + z). \quad (57)$$

An n -digit point $z \in K_{[0,1]}$ corresponding to $y = (0.q_1 \dots q_n)_4$, $q_n \neq 0$, is given by

$$\hat{q}_1 \circ \dots \circ \hat{q}_n(0) = g(y) \quad (58)$$




Fig. 3. Koch curves and their generator (the upper inset) parametrized by α and corresponding to (54)–(58). From highest to lowest: $\alpha = \pi/2.5$, $\alpha = \pi/3$, $\alpha = \pi/4$, $\alpha = \pi/6$.

(value at 0 of the composition of maps). If $y_n = (0.q_1 \dots q_n)_4$ is a Cauchy sequence convergent to $y = \lim_{n \rightarrow \infty} y_n$, then $g(y) = \lim_{n \rightarrow \infty} g(y_n)$. Curves from Fig. 3 are the images $g([0, 1])$ for various α . g is one-one, so it defines the inverse bijection $g^{-1} = f : K_{[0,1]} \rightarrow [0, 1]$.

In order to have a better feel of our bijection, let us have a look at the relation between the standard $\pi/3$ Koch curve and its quaternary parametrization, as illustrated in Fig. 4. Decreasing the initiator [26, 27] of the Koch curve three times is equivalent to dividing each vertex number by four (*i.e.* shifting left the decimal separator by one position). The bijection is, therefore, equivalent to parametrization of the Koch curve by its Hausdorff integral, in exact analogy to the construction of Epstein and Śniatycki [26, 27]. The authors of [26, 27] begin with the integral and obtain derivatives by means of the fundamental theorem of calculus. The arithmetic approach begins with the derivative, and then the integral is defined through the fundamental theorem of non-Newtonian calculus.

Another formalism that has to be mentioned in this context begins with the notion of a mass function [28], a concept in some respects similar to the Hausdorff measure, but easier to compute. Interestingly, it can be shown that the resulting ‘ F^α -calculus’ implicitly involves a bijection (see Sec. 6 in [28]), playing exactly the same role as the bijections occurring in a general non-Newtonian calculus, namely conjugating non-Newtonian derivatives and integrals with their Newtonian counterparts. The bijection is then, more or less explicitly, applied to Langevin, Newton and Schrödinger equations on fractal curves and space-times [29–31]. However, the formalism heavily




Fig. 4. Link between a vertex position in the $\alpha = \pi/3$ Koch curve and its numbering by $y = (0.q_1 \dots q_j \dots)_4 \in [0, 1]$ in quaternary representation. Rescaling the unit segment three times, we obtain a smaller copy of the Koch curve. The corresponding vertices of the two curves are numbered by identical digits, with digital separators shifted by one position. The number y can be thus regarded as a Hausdorff measure of the part of the Koch curve extending between the origin and the vertex, if we normalize the measure to 1 on the segment $[0, 1]$. The rule applies to all the Koch curves generated by (54)–(57).

depends on structure and parametrization of the fractal sets, and reduces to the ordinary Newtonian one on non-fractal domains. This should be contrasted with the general non-Newtonian approach, working whenever the sets in question have cardinality of continuum. Moreover, the arithmetic perspective, advocated in the present work, is not restricted to calculus only, but includes algebraic or probabilistic aspects as well. This is why the non-Newtonian wave equation will be shown to be Lorentz covariant, even if fractal space-time is modeled by a Cartesian product of different fractals.

Let us finally note here that the more traditional approaches to fractal analysis [32, 33] have not managed to formulate any calculus on fractals of a Koch-curve type.

For $\alpha = \pi/3$, we obtain the standard curve, generated by equilateral triangles. Similarity dimension of a curve generated by (54)–(58) is given by (Fig. 5)

$$D = \frac{\log 4}{\log(2 + 2 \cos \alpha)}. \quad (59)$$

There are many ways of extending the Koch curve from $K_{[0,1]}$ to $K_{\mathbb{R}}$. For example, let $K_{[k,k+1]}$ be the curve $K_{[0,1]}$ shifted according to $z \mapsto z+k$, $k \in \mathbb{Z}$. Then $K_{\mathbb{R}} = \cup_{k \in \mathbb{Z}} K_{[k,k+1]}$ is a periodic Koch curve, with the bijection $f : K_{\mathbb{R}} \rightarrow \mathbb{R}$ constructed from appropriately shifted maps g defined above. Non-periodic but self-similar extensions can be obtained by shifts and rescalings. From our point of view, the only condition we impose on f is the continuity of $g = f^{-1}$ at 0, *i.e.* $\lim_{y \rightarrow 0^-} g(y) = \lim_{y \rightarrow 0^+} g(y) = g(0)$. We take $g(0) = 0$.




Fig. 5. Similarity dimension D and the length L of the generator from Fig. 1 as functions of α . The horizontal lines show the values for the standard $\pi/3$ Koch curve.

Combining the generalized Koch curves, we can construct a curve which is in a one–one relation with \mathbb{R} , with explicitly given bijection f , and whose fractal dimensions vary from segment to segment in a prescribed way. This type of generalization may be useful for applications involving realistic coastlines, whose fractal dimensions coincide with the data described by the Richardson law [34]. In what follows, we will concentrate on the simple case $\alpha = \pi/3$, $L = 1/3$, of the standard Koch curve.

3. Wave equation on Koch curves

First of all, let us assume we discuss a real-valued field, whose evolution on the Koch curve $\mathbb{X} = K_{\mathbb{R}}$ is described with respect to a ‘normal’ non-fractal time t . The field is thus represented by $\mathbb{R} \times \mathbb{X} \mapsto \Phi_t(x) \in \mathbb{R}$, with $x \in \mathbb{X}$. Since $\mathbb{Y} = \mathbb{R}$, we take $f_{\mathbb{Y}} = \text{id}_{\mathbb{R}}$. (Although $f_{\mathbb{Y}}(y) = y^3$ or any other

bijection would do as well, leading to a different behavior of the wave.) The wave equation is

$$\frac{1}{c^2} \frac{d^2}{dt^2} \Phi_t(x) - \frac{D^2}{Dx^2} \Phi_t(x) = 0, \quad (60)$$

where

$$\frac{d}{dt} \Phi_t(x) = \frac{\lim_{h \rightarrow 0} (\Phi_{t+h}(x) - \Phi_t(x))}{h}, \quad (61)$$

$$\frac{D}{Dx} \Phi_t(x) = \frac{\lim_{h \rightarrow 0} (\Phi_t(x \oplus_{\mathbb{X}} f_{\mathbb{X}}^{-1}(h)) - \Phi_t(x))}{h}. \quad (62)$$

We search solutions in the form (here $y = ct$)

$$\Phi_t(x) = A(x, y) + B(x, y), \quad (63)$$

where

$$\left(\frac{d}{dy} - \frac{D}{Dx} \right) A(x, y) = \left(\frac{d}{dy} + \frac{D}{Dx} \right) B(x, y) \equiv 0, \quad (64)$$

suggesting simply

$$A(x, y) = a(f_{\mathbb{X}}(x) + y), \quad (65)$$

$$B(x, y) = b(f_{\mathbb{X}}(x) - y) \quad (66)$$

for some twice differentiable $a, b : \mathbb{R} \rightarrow \mathbb{R}$.

Indeed, from definitions

$$\begin{aligned} \frac{D}{Dx} A(x, y) &= \lim_{h \rightarrow 0} \frac{A(x \oplus_{\mathbb{X}} f_{\mathbb{X}}^{-1}(h), y) - A(x, y)}{h} \\ &= \lim_{h \rightarrow 0} \frac{a(f_{\mathbb{X}}(x) + h + y) - a(f_{\mathbb{X}}(x) + y)}{h} \\ &\equiv \frac{d}{dy} a(f_{\mathbb{X}}(x) + y) = \frac{d}{dy} A(x, y). \end{aligned} \quad (67)$$

One similarly verifies that d/dy and D/Dx commute, and

$$\frac{D}{Dx} B(x, y) \equiv -\frac{d}{dy} B(x, y). \quad (68)$$

Figure 6 shows the dynamics of $\Phi_t(x)$ with $a = 0$. The energy of the wave is given by

$$E = \frac{1}{2} \int_{f_{\mathbb{X}}^{-1}(-\infty)}^{f_{\mathbb{X}}^{-1}(\infty)} \left(\frac{1}{c^2} \left| \frac{d\Phi_t(x)}{dt} \right|^2 + \left| \frac{D\Phi_t(x)}{Dx} \right|^2 \right) Dx, \tag{69}$$

where the integral is defined by (13).




Fig. 6. ‘Aurora borealis wave’: Six snapshots of $\Phi_t(x)$ propagating to the right along the Koch curve. The upper plot shows the corresponding function b occurring in (66).

Let us explicitly check the time independence of E for the particular case of $\Phi_t(x) = a(f_{\mathbb{X}}(x) + ct)$. Let $a'(x) = da(x)/dx$ be the Newtonian derivative. Then,

$$\begin{aligned} E &= \int_{-\infty}^{\infty} |a'(f_{\mathbb{X}} \circ f_{\mathbb{X}}^{-1}(x) + ct)|^2 dx \\ &= \int_{-\infty}^{\infty} |a'(x)|^2 dx \end{aligned} \quad (70)$$

is independent of time, as it should be.

4. Lorentz covariance

In our model, space-time consists of points $(x^0, x^1) = (ct, x) \in \mathbb{R} \times \mathbb{X}$, with $(x^0, f_{\mathbb{X}}(x^1)) \in \mathbb{R}^2$. A Lorentz transformation $x' = \mathcal{L}(x)$, $\mathcal{L} : \mathbb{R} \times \mathbb{X} \rightarrow \mathbb{R} \times \mathbb{X}$, is defined by

$$\begin{pmatrix} x'^0 \\ x'^1 \end{pmatrix} = \begin{pmatrix} L^0_0 x^0 + L^0_1 f_{\mathbb{X}}(x^1) \\ f_{\mathbb{X}}^{-1}(L^1_0 x^0 + L^1_1 f_{\mathbb{X}}(x^1)) \end{pmatrix}, \quad (71)$$

or, equivalently, by

$$\begin{pmatrix} x'^0 \\ f_{\mathbb{X}}(x'^1) \end{pmatrix} = \begin{pmatrix} L^0_0 & L^0_1 \\ L^1_0 & L^1_1 \end{pmatrix} \begin{pmatrix} x^0 \\ f_{\mathbb{X}}(x^1) \end{pmatrix}, \quad (72)$$

where $L \in \text{SO}(1, 1)$. Equation (71) implements a nonlinear action of the group $\text{SO}(1, 1)$, and reduces to the usual representation if $\mathbb{X} = \mathbb{R}$ and $f_{\mathbb{X}}(x^1) = x^1$. Transformations (71) form a group.

In order to prove Lorentz invariance of the wave equation, let us first note that its solution

$$\begin{aligned} \Phi_t(x) &= a(f_{\mathbb{X}}(x^1) + x^0) + b(f_{\mathbb{X}}(x^1) - x^0) \\ &= \phi(x^0, f_{\mathbb{X}}(x^1)) \end{aligned} \quad (73)$$

defines a function ϕ , satisfying (due to triviality of $f_{\mathbb{Y}}$)

$$\frac{D\Phi_t(x)}{Dx} = \frac{\partial \phi(x^0, f_{\mathbb{X}}(x^1))}{\partial f_{\mathbb{X}}(x^1)}, \quad (74)$$

$$\frac{1}{c} \frac{d\Phi_t(x)}{dt} = \frac{\partial \phi(x^0, f_{\mathbb{X}}(x^1))}{\partial x^0}. \quad (75)$$

Accordingly, the wave equation takes the standard form of

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial f_{\mathbb{X}}(x^1)^2} \right) \phi(x^0, f_{\mathbb{X}}(x^1)) = 0. \quad (76)$$

It is invariant under (72) if ϕ transforms by

$$\phi'(x'^0, f_{\mathbb{X}}(x'^1)) = \phi(x^0, f_{\mathbb{X}}(x^1)), \quad (77)$$

which is equivalent to the scalar-field transformation $\Phi'_{t'}(x') = \Phi_t(x)$.

Replacing $\mathbb{R} \times \mathbb{X}$ by a more general case $\mathbb{X}_0 \times \mathbb{X}_1$, $f_{\mathbb{X}_j} : \mathbb{X}_j \rightarrow \mathbb{R}$ (cf. [30]), one arrives at a Lorentz invariant wave equation (with both space-time derivatives appropriately defined), and Lorentz transformations

$$\begin{pmatrix} x'^0 \\ x'^1 \end{pmatrix} = \begin{pmatrix} f_{\mathbb{X}_0}^{-1} \left(L^0_0 f_{\mathbb{X}_0}(x^0) + L^0_1 f_{\mathbb{X}_1}(x^1) \right) \\ f_{\mathbb{X}_1}^{-1} \left(L^1_0 f_{\mathbb{X}_0}(x^0) + L^1_1 f_{\mathbb{X}_1}(x^1) \right) \end{pmatrix}. \quad (78)$$

A generalization to space-times constructed by Cartesian products of arbitrary numbers of fractals is now obvious.

5. Conclusions

To conclude, we have obtained a wave that propagates along a Koch-type curve. The wave possesses finite conserved energy and satisfies the usual wave equation, formulated with respect to appropriately defined derivatives. The derivatives are not the ones we know from the standard mathematical literature of the subject, but are very natural and easy to work with. The solution we have found is the general one, a fact following from the standard form (76) of the wave equation. The velocity of the wave is intriguing. On the one hand, it is described by the parameter c in the wave equation. On the other hand, however, the length of any piece of a fractal coast is infinite and yet the wave moves from point to point in a finite time, and with speed that *looks* finite and natural. This is possible since the fractal sum $z = x \oplus_{\mathbb{X}} y$ of two points in a Koch curve is uniquely defined in spite of the apparently ‘infinite’ distances between x , y , z and the origin 0. Another interesting aspect of the resulting motion is the lack of difficulties with combining non-fractal time with fractal space. Lorentz transformations in the corresponding space-time have been constructed, and Lorentz invariance of the wave equation has been proved. Fractal arithmetic automatically tames the infinities inherent in the length of the curve. It would not be very surprising if our fractal calculus found applications also in other branches of physics, where finite physical results are buried in apparently infinite theoretical predictions.

REFERENCES

- [1] M. Grossman, R. Katz, *Non-Newtonian Calculus*, Lee Press, Pigeon Cove, 1972.
- [2] M. Grossman, *The First Nonlinear System of Differential and Integral Calculus*, Mathco, Rockport (1979).
- [3] M. Grossman, *Bigeometric Calculus: A System with Scale-Free Derivative*, Archimedes Foundation, Rockport, 1983.
- [4] M. Czachor, *Quantum Stud.: Math. Found.* **3**, 123 (2016).
- [5] D. Aerts, M. Czachor, M. Kuna, *Chaos, Solitons Fract.* **83**, 201 (2016).
- [6] D. Aerts, M. Czachor, M. Kuna, *Chaos, Solitons Fract.* **91**, 461 (2016).
- [7] D. Aerts, M. Czachor, M. Kuna, *Rep. Math. Phys.* **81**, 357 (2018).
- [8] M. Czachor, *Int. J. Theor. Phys.* **56**, 1364 (2017).
- [9] Z. Domański, M. Błaszak, [arXiv:1706.00980 \[math-ph\]](#).
- [10] M. Burgin, Non-classical Models of Natural Numbers, *Russ. Math. Surveys* **32**, 209-210 (1977) (in Russian).
- [11] M. Burgin, *Non-Diophantine Arithmetics, or is it Possible that $2 + 2$ is not Equal to 4 ?*, Ukrainian Academy of Information Sciences, Kiev 1997 (in Russian).
- [12] M. Burgin, [arXiv:1010.3287 \[math.GM\]](#).
- [13] M. Burgin, G. Meissner, *Appl. Math.* **8**, 133 (2017).
- [14] P. Benioff, *Int. J. Theor. Phys.* **50**, 1887 (2011).
- [15] P. Benioff, *Quantum Stud.: Math. Found.* **2**, 289 (2015).
- [16] P. Benioff, *Quantum Inf. Proc.* **15**, 1081 (2016).
- [17] P.E.T. Jorgensen, S. Pedersen, *J. Anal. Math.* **75**, 185 (1998).
- [18] P.E.T. Jorgensen, *Analysis and Probability: Wavelets, Signals, Fractals*, Springer, New York 2006.
- [19] M. Czachor, *Information Processing and Fechner's Problem as a Choice of Arithmetic*, in: M. Burgin, W. Hofkirchner (Eds.), *Information Studies and the Quest for Interdisciplinarity: Unity Through Diversity*, World Scientific, Singapore 2017, pp. 363–372.
- [20] D. Filip, C. Piatecki, *Math. Aeterna* **4**, 101 (2014).
- [21] D. Aniszewska, *Nonlinear Dyn.* **50**, 265 (2007).
- [22] A.E. Bashirov, E. Mısırlı, A. Özyapıcı, *J. Math. Anal. Appl.* **337**, 36 (2008).
- [23] L. Florack, H. van Assen, *J. Math. Imaging Vis.* **42**, 64 (2012).
- [24] A. Ozyapıcı, B. Bilgehan, *Numer. Algorithms* **71**, 475 (2016).
- [25] N. Yalcina, E. Celikb, A. Gokdogana, *Optik* **127**, 9984 (2016).
- [26] M. Epstein, J. Śniatycki, *Physica D* **220**, 54 (2006).
- [27] M. Epstein, J. Śniatycki, *Chaos, Solitons Fract.* **38**, 334 (2008).
- [28] A. Parvate, S. Satin, A.D. Gangal, *Fractals* **19**, 15 (2011).

- [29] S. Satin, A.D. Gangal, *Fractals* **24**, 1650028 (2016).
- [30] A.K. Golmankhaneh, A.K. Golmankhaneh, D. Baleanu, *Int. J. Theor. Phys.* **54**, 1275 (2015).
- [31] A.K. Golmankhaneh, *Turk. J. Phys.* **41**, 418 (2017).
- [32] J. Kigami, *Analysis on Fractals*, Cambridge Tracts in Mathematics, vol. 143, Cambridge University Press, Cambridge 2001.
- [33] R.S. Strichartz, *Differential Equations on Fractals*, Princeton University Press, Princeton 2006.
- [34] B. Mandelbrot, *Science* **155**, 636 (1967).