

HYDRODYNAMICS FORMALISM WITH SPIN DYNAMICS*

RAJEEV SINGH

Institute of Nuclear Physics Polish Academy of Sciences, 31-342 Kraków, Poland

(Received November 25, 2020)

We review the key steps of the relativistic fluid dynamics formalism with spin degrees of freedom initiated recently. We obtain equations of motion of the expansion of the system from the underlying definitions of quantum kinetic theory for the equilibrium phase-space distribution functions. We investigate the dynamics of spin polarization of the system in the Bjorken hydrodynamical background.

DOI:10.5506/APhysPolBSupp.14.461

1. Introduction

Spin polarization experimental measurements of Λ hyperons recently taken by the STAR Collaboration [1–4] have created a huge interest in the spin polarization studies and in studies correlating between the vorticity and particle spin polarization in relativistic heavy-ion collisions [5–41]; for reviews, see [42–46]. Thermal-based models [47–50], which precisely explain the global polarization of particles, are able to explain differential results correctly [4]. These models assume the condition that particle spin polarization emitted at freeze-out hypersurface is defined by the thermodynamical quantity which is named as thermal vorticity [6, 51], not considering the fact that it may independently evolve during the expansion of the fluid. In this article, we follow scheme proposed in Refs. [52–57] and analyze such a possibility of spin polarization evolution using relativistic hydrodynamics framework with spin.

2. Distribution functions in equilibrium

If we know phase-space distribution function for the system's equilibrium state, then it is possible to derive relativistic hydrodynamics from the underlying kinetic theory definitions [58]. Following ideas developed by Becattini *et al.* [6], we take into consideration the following distribution functions for

* Presented at the on-line meeting *Criticality in QCD and the Hadron Resonance Gas*, Wrocław, Poland, July 29–31, 2020.

the relativistic systems of spin 1/2 massive particles (and antiparticles) in the local equilibrium state:

$$f_{rs}^+(x, p) = \bar{u}_r(p)X^+u_s(p), \quad f_{rs}^-(x, p) = -\bar{v}_s(p)X^-v_r(p), \quad (1)$$

where x and p is the space-time position coordinate and the four-vector momentum, respectively, with $u_r(p)$ and $v_r(p)$ being the Dirac bispinors ($r, s = 1, 2$). Dirac bispinors follow the normalization conditions as $\bar{u}_r(p)u_s(p) = \delta_{rs}$ and $\bar{v}_r(p)v_s(p) = -\delta_{rs}$, here δ_{rs} is the Dirac delta function and X^\pm have the following form in terms of relativistic Boltzmann distributions:

$$X^\pm = \exp \left[\pm \xi(x) - \beta_\mu(x)p^\mu \pm \frac{1}{2}\omega_{\mu\nu}(x)\Sigma^{\mu\nu} \right],$$

where $\beta^\mu \equiv U^\mu/T$ and $\xi \equiv \mu/T$, here T , μ and U^μ is the temperature, baryon chemical potential and four-vector velocity, respectively, and $\omega_{\mu\nu}$ is the second rank asymmetric tensor known as a spin polarization tensor with $\Sigma^{\mu\nu} \equiv \frac{i}{4}[\gamma^\mu, \gamma^\nu]$ being the spin operator.

With the help of the expressions from Ref. [59] and Eqs. (1), we can obtain the Wigner functions in equilibrium as follows:

$$\begin{aligned} \mathcal{W}_{\text{eq}}^\pm(x, k) &= \frac{e^{\pm\xi}}{4m} \int dP e^{-\beta \cdot p} \delta^{(4)}(k \mp p) \\ &\times \left[2m(m \pm \not{p}) \cosh(\zeta) \pm \frac{\sinh(\zeta)}{2\zeta} \omega_{\mu\nu} (\not{p} \pm m) \Sigma^{\mu\nu} (\not{p} \pm m) \right], \quad (2) \end{aligned}$$

where k being the off mass-shell particle four-momentum, $dP = d^3p/((2\pi)^3 E_p)$ is the invariant measure with $E_p = \sqrt{m^2 + \mathbf{p}^2}$ denoting the on-shell energy of the particle, and $\zeta = \frac{1}{2\sqrt{2}}\sqrt{\omega_{\mu\nu}\omega^{\mu\nu}}$ in terms of spin polarization tensor.

The Wigner function can also be expanded using the Clifford algebra expansion (2)

$$\begin{aligned} \mathcal{W}_{\text{eq}}^\pm(x, k) &= \frac{1}{4} \left[\mathcal{F}_{\text{eq}}^\pm(x, k) + i\gamma_5 \mathcal{P}_{\text{eq}}^\pm(x, k) + \gamma^\mu \mathcal{V}_{\text{eq},\mu}^\pm(x, k) \right. \\ &\quad \left. + \gamma_5 \gamma^\mu \mathcal{A}_{\text{eq},\mu}^\pm(x, k) + \Sigma^{\mu\nu} \mathcal{S}_{\text{eq},\mu\nu}^\pm(x, k) \right], \end{aligned}$$

where $\mathcal{X} \in \{\mathcal{F}, \mathcal{P}, \mathcal{V}_\mu, \mathcal{A}_\mu, \mathcal{S}_{\nu\mu}\}$ are the coefficient functions of the Wigner function, which can be obtained from the trace of $\mathcal{W}_{\text{eq}}^\pm(x, k)$ multiplying first by: $\{\mathbf{1}, -i\gamma_5, \gamma_\mu, \gamma_\mu\gamma_5, 2\Sigma_{\mu\nu}\}$.

3. Kinetic and hydrodynamical equations

The kinetic equation to be followed by Wigner function is

$$(\gamma_\mu K^\mu - m) \mathcal{W}(x, k) = C[\mathcal{W}(x, k)], \quad (3)$$

with $K^\mu = k^\mu + \frac{i\hbar}{2} \partial^\mu$. For a global equilibrium state, the Wigner function follows exactly Eq. (3) with the collision term $C[\mathcal{W}(x, k)] = 0$. The widely used method of treating Eq. (3) is the semi-classical expansion method of the coefficient functions of the Wigner function

$$\mathcal{X} = \mathcal{X}^{(0)} + \hbar \mathcal{X}^{(1)} + \hbar^2 \mathcal{X}^{(2)} + \dots$$

Up to the first order (*i.e.* next-to-leading order) in \hbar , the treatment of Eq. (3) gives the following kinetic equations to be followed by the two independent coefficients which are: \mathcal{F}_{eq} and $\mathcal{A}_{\text{eq}}^\nu$,

$$k^\mu \partial_\mu \mathcal{F}_{\text{eq}}(x, k) = 0, \quad k^\mu \partial_\mu \mathcal{A}_{\text{eq}}^\nu(x, k) = 0, \quad k_\nu \mathcal{A}_{\text{eq}}^\nu(x, k) = 0. \quad (4)$$

In the case of the global equilibrium, Eqs. (4) are satisfied exactly which in turn yields the conditions that β_μ is a Killing vector, whereas ξ and $\omega_{\mu\nu}$ are constants, but $\omega_{\mu\nu}$ does not necessarily be equal to thermal vorticity $\varpi_{\mu\nu} = -\frac{1}{2}(\partial_\mu \beta_\nu - \partial_\nu \beta_\mu) = \text{const}$. However, in the case of a local equilibrium, Eqs. (4) are not exactly followed. Here, we follow [60], and by permitting β , ξ and ω dependence on x , we need to have only certain moments in the momentum space of the kinetic equations (4) which are satisfied, which lead to conservation laws for charge, energy-linear momentum and spin [56]

$$\partial_\mu N^\mu = 0, \quad (5)$$

$$\partial_\mu T_{\text{GLW}}^{\mu\nu} = 0, \quad (6)$$

$$\partial_\lambda S_{\text{GLW}}^{\lambda, \alpha\beta} = 0. \quad (7)$$

Here, the baryon current, the energy momentum and the spin tensors are based on the forms by the de Groot–van Leeuwen–van Weert (GLW) [59]

$$N^\alpha = nU^\alpha, \quad (8)$$

$$T_{\text{GLW}}^{\alpha\beta} = (\varepsilon + P)U^\alpha U^\beta - P g^{\alpha\beta}, \quad (9)$$

$$S_{\text{GLW}}^{\alpha, \beta\gamma} = \cosh(\xi) \left[n_{(0)} U^\alpha \omega^{\beta\gamma} + \mathcal{A}_{(0)} U^\alpha U^\delta U^{[\beta} \omega^{\gamma]}_\delta \right. \quad (10)$$

$$\left. + \mathcal{B}_{(0)} \left(U^{[\beta} \Delta^{\alpha\delta} \omega^{\gamma]}_\delta + U^\alpha \Delta^{\delta[\beta} \omega^{\gamma]}_\delta + U^\delta \Delta^{\alpha[\beta} \omega^{\gamma]}_\delta \right) \right], \quad (11)$$

where $\Delta^{\alpha\beta} = g^{\alpha\beta} - U^\alpha U^\beta$ is the spatial projection operator which is orthogonal to the hydrodynamic flow 4-vector U .

For the case of polarization tensor in the leading order, the baryon number density, the energy density, and the pressure are expressed, respectively, as

$$n = \sinh(\xi) n_{(0)}(T), \quad (12)$$

$$\varepsilon = \cosh(\xi) \varepsilon_{(0)}(T), \quad (13)$$

$$P = \cosh(\xi) P_{(0)}(T), \quad (14)$$

where for spin-less and neutral massive Boltzmann particles, thermodynamical properties are defined by [61]

$$n_{(0)}(T) = \frac{2 T^3}{\pi^2} \hat{m}^2 K_2(\hat{m}), \quad (15)$$

$$\varepsilon_{(0)}(T) = \frac{2 T^4}{\pi^2} \hat{m}^2 [3K_2(\hat{m}) + \hat{m}K_1(\hat{m})], \quad (16)$$

$$P_{(0)}(T) = T n_{(0)}(T). \quad (17)$$

Here, $K_1(\hat{m})$ and $K_2(\hat{m})$ are modified Bessel functions of 1st and 2nd kind respectively. The

$$\mathcal{B}_{(0)} = -\frac{2}{\hat{m}^2} s_{(0)}(T), \quad \mathcal{A}_{(0)} = -3\mathcal{B}_{(0)} + 2n_{(0)}(T) \quad (18)$$

with entropy density $s_{(0)} = (\varepsilon_{(0)} + P_{(0)})/T$ and $\hat{m} = m/T$.

4. Bjorken expansion set-up

Since the spin polarization tensor $\omega_{\mu\nu}$ is a 2nd rank asymmetric tensor, as in analogy to the Faraday electromagnetic field strength tensor, it can be written into electric-like (κ) and magnetic-like (ω) components

$$\omega_{\mu\nu} = \kappa_\mu U_\nu - \kappa_\nu U_\mu + \epsilon_{\mu\nu\alpha\beta} U^\alpha \omega^\beta, \quad (19)$$

where κ and ω are 4-vectors, orthogonal to fluid flow vector U_μ . For longitudinal boost-invariant and transversely homogeneous systems [62, 63], one can write the following basis vectors:

$$\begin{aligned} U^\alpha &= \frac{1}{\tau} (t, 0, 0, z) = (\cosh(\eta), 0, 0, \sinh(\eta)), \\ X^\alpha &= (0, 1, 0, 0), \\ Y^\alpha &= (0, 0, 1, 0), \\ Z^\alpha &= \frac{1}{\tau} (z, 0, 0, t) = (\sinh(\eta), 0, 0, \cosh(\eta)), \end{aligned} \quad (20)$$

where longitudinal proper time is defined as $\tau = \sqrt{t^2 - z^2}$ and the space-time rapidity is defined as $\eta = \frac{1}{2} \ln((t+z)/(t-z))$. The normalization conditions satisfied by the basis vectors (20) are

$$\begin{aligned} U \cdot U &= 1, \\ X \cdot X &= Y \cdot Y = Z \cdot Z = -1, \\ X \cdot U &= Y \cdot U = Z \cdot U = 0, \\ X \cdot Y &= Y \cdot Z = Z \cdot X = 0. \end{aligned} \quad (21)$$

Using the fact that κ and ω are orthogonal to U_μ and Eqs. (21), κ^μ and ω^μ can be written as

$$\begin{aligned} \kappa^\alpha &= C_{\kappa X}(\tau)X^\alpha + C_{\kappa Y}(\tau)Y^\alpha + C_{\kappa Z}(\tau)Z^\alpha, \\ \omega^\alpha &= C_{\omega X}(\tau)X^\alpha + C_{\omega Y}(\tau)Y^\alpha + C_{\omega Z}(\tau)Z^\alpha, \end{aligned} \tag{22}$$

where one can notice that the scalar functions depend only on proper time (τ).

Putting Eqs. (22) into Eq. (7) and then using the projection method, we project the resulting tensor on different combination of basis vectors $U_\alpha X_\beta$, $U_\alpha Y_\beta$, $U_\alpha Z_\beta$, $Y_\alpha Z_\beta$, $X_\alpha Z_\beta$ and $X_\alpha Y_\beta$, and get the six equations of motions as

$$\text{diag}(\mathcal{L}, \mathcal{L}, \mathcal{L}, \mathcal{P}, \mathcal{P}, \mathcal{P}) \dot{\mathbf{C}} = \text{diag}(\mathcal{Q}_1, \mathcal{Q}_1, \mathcal{Q}_2, \mathcal{R}_1, \mathcal{R}_1, \mathcal{R}_2) \mathbf{C}, \tag{23}$$

where $\mathbf{C} = (C_{\kappa X}, C_{\kappa Y}, C_{\kappa Z}, C_{\omega X}, C_{\omega Y}, C_{\omega Z})$, $(\dot{\dots}) \equiv U \cdot \partial = \partial_\tau$ and

$$\begin{aligned} \mathcal{L}(\tau) &= \mathcal{A}_1 - \frac{1}{2}\mathcal{A}_2 - \mathcal{A}_3, \\ \mathcal{P}(\tau) &= \mathcal{A}_1, \\ \mathcal{Q}_1(\tau) &= - \left[\dot{\mathcal{L}} + \frac{1}{\tau} \left(\mathcal{L} + \frac{1}{2}\mathcal{A}_3 \right) \right], \\ \mathcal{Q}_2(\tau) &= - \left(\dot{\mathcal{L}} + \frac{\mathcal{L}}{\tau} \right), \\ \mathcal{R}_1(\tau) &= - \left[\dot{\mathcal{P}} + \frac{1}{\tau} \left(\mathcal{P} - \frac{1}{2}\mathcal{A}_3 \right) \right], \\ \mathcal{R}_2(\tau) &= - \left(\dot{\mathcal{P}} + \frac{\mathcal{P}}{\tau} \right), \end{aligned}$$

with

$$\begin{aligned} \mathcal{A}_1 &= \cosh(\xi) (n_{(0)} - \mathcal{B}_{(0)}), \\ \mathcal{A}_2 &= \cosh(\xi) (\mathcal{A}_{(0)} - 3\mathcal{B}_{(0)}), \\ \mathcal{A}_3 &= \cosh(\xi) \mathcal{B}_{(0)}. \end{aligned}$$

Equation (23) implies that the C functions evolve independently of each other for the case of Bjorken flow, and $C_{\kappa X}$ and $C_{\kappa Y}$ (similarly $C_{\omega X}$ and $C_{\omega Y}$) follow the same form of evolution equations due to the rotational invariance.

The charge current conservation (5) for the Bjorken-type flow is expressed as

$$\frac{dn}{d\tau} + \frac{n}{\tau} = 0, \tag{24}$$

whereas the energy and linear momentum conservation law (6) (after projecting on U) yields

$$\frac{d\varepsilon}{d\tau} + \frac{(\varepsilon + P)}{\tau} = 0. \tag{25}$$

5. Particle spin polarization at freeze-out

To calculate the mean spin polarization per particle, the following formula is used [56]:

$$\langle \pi_\mu \rangle = E_p \frac{d\Pi_\mu(p)}{d^3p} \Big/ E_p \frac{d\mathcal{N}(p)}{d^3p}, \tag{26}$$

with $E_p \frac{d\Pi_\mu(p)}{d^3p}$ being the total value of the Pauli–Lubański vector (after integrating over the freeze-out hypersurface, $\Delta\Sigma_\lambda$),

$$E_p \frac{d\Pi_\mu(p)}{d^3p} = -\frac{\cosh(\xi)}{(2\pi)^3 m} \int \Delta\Sigma_\lambda p^\lambda e^{-\beta \cdot p} \tilde{\omega}_{\mu\beta} p^\beta,$$

and

$$E_p \frac{d\mathcal{N}(p)}{d^3p} = \frac{4 \cosh(\xi)}{(2\pi)^3} \int \Delta\Sigma_\lambda p^\lambda e^{-\beta \cdot p}$$

is the total momentum density of both particles and antiparticles with four-momentum given as $p^\lambda = (m_T \cosh y_p, p_x, p_y, m_T \sinh y_p)$.

After performing the canonical boost [64] of (26), we obtain the polarization vector $\langle \pi_\mu^* \rangle$ in the local rest frame of the particle as

$$\langle \pi_\mu^* \rangle = -\frac{1}{8m} \begin{bmatrix} 0 \\ \left(\frac{p_x \sinh y_p}{b}\right) a_i + \left(\frac{\chi p_x \cosh y_p}{b}\right) a_j + 2C_{\kappa Z} p_y - \chi C_{\omega X} m_T \\ \left(\frac{p_y \sinh y_p}{b}\right) a_i + \left(\frac{\chi p_y \cosh y_p}{b}\right) a_j - 2C_{\kappa Z} p_x - \chi C_{\omega Y} m_T \\ - \left(\frac{m \cosh y_p + m_T}{b}\right) a_i - \left(\frac{\chi m \sinh y_p}{b}\right) a_j \end{bmatrix}, \tag{27}$$

with $a_i = \chi (C_{\kappa X} p_y - C_{\kappa Y} p_x) + 2C_{\omega Z} m_T$, $a_j = C_{\omega X} p_x + C_{\omega Y} p_y$, $b = m_T \cosh y_p + m$, and $\chi = (K_0(\hat{m}_T) + K_2(\hat{m}_T)) / K_1(\hat{m}_T)$ and $\hat{m}_T = m_T / T$.

6. Results

Here, we show the solutions of differential equations (23), (24), and (25). System is initialized at the initial proper time $\tau_0 = 1$ fm with initial temperature and the initial baryon chemical potential as $T_0 = T(\tau_0) = 150$ MeV and $\mu_0 = \mu(\tau_0) = 800$ MeV, respectively. Here, the system is assumed to be formed with Λ particles having mass $m = 1116$ MeV. In Fig. 1, the proper-time dependence of temperature and baryon chemical potential is depicted, where the temperature decreases with the proper time, whereas the ratio of baryon chemical potential and temperature increases with the proper time. From Fig. 2, the proper-time dependence of the C functions can be known which describe the system's spin polarization evolution.




Fig. 1. (Colour on-line) Dependence of the temperature re-scaled by its initial value (solid black line) and the ratio of baryon chemical potential over temperature re-scaled by the initial ratio (dotted blue line) on the proper time.




Fig. 2. (Colour on-line) Dependence of scalar functions $C_{\kappa X}$ (solid black line), $C_{\kappa Z}$ (dash-dotted blue line), $C_{\omega X}$ (dotted red line) and $C_{\omega Z}$ (dashed green line) on the proper time.

Using the information of the thermodynamic parameters and C coefficients evolution, we can calculate the different components of the mean polarization vector in the rest frame of the particles $\langle \pi_\mu^* \rangle$ at freeze-out, see Fig. 3. We note that $\langle \pi_y^* \rangle$ is negative reflecting the system's initial spin polarization. Due to the Bjorken symmetry which we have assumed in our calculations in this article, the longitudinal component ($\langle \pi_z^* \rangle$) of the mean polarization vector is vanishing which is not in agreement with the quadrupole structure of the longitudinal component of the spin polarization seen in the experiment. However, we note here that $\langle \pi_x^* \rangle$ shows quadrupole structure. We nevertheless see and note that the Bjorken set-up is very simple to address the measurements done by the experiment.




Fig. 3. Different components of the mean polarization of Λ particles in the rest frame of the particle obtained with the initial values $\mu_0 = 800$ MeV, $T_0 = 155$ MeV, $C_{\kappa,0} = (0, 0, 0)$ and $C_{\omega,0} = (0, 0.1, 0)$ for $y_p = 0$.

7. Summary

We briefly presented the key ingredients of relativistic perfect-fluid hydrodynamics with spin framework initiated recently. From the definitions of kinetic theory for the equilibrium phase-space distribution functions in the local equilibrium, we obtained the equations of motions for the expansion of the system. For the case of the Bjorken-type flow, we investigated the system's spin polarization dynamics, which in turn shows that the scalar functions describing the dynamics of the spin polarization evolve independently of each other. These results are used to obtain the particle spin polarization at the freeze-out hypersurface. We, however, note that within the current, a simple set-up of Bjorken symmetry experimental measurements cannot be addressed properly.

R.S. would like to thank Wojciech Florkowski, Radosław Ryblewski and Avdhesh Kumar for clarifying discussions. This research is supported in part by the National Science Centre, Poland (NCN) grants No. 2016/23/B/ST2/00717 and No. 2018/30/E/ST2/00432.

REFERENCES

- [1] STAR Collaboration (L. Adamczyk *et al.*), *Nature* **548**, 62 (2017).
- [2] STAR Collaboration (J. Adam *et al.*), *Phys. Rev. C* **98**, 014910 (2018).
- [3] STAR Collaboration (T. Niida), *Nucl. Phys. A* **982**, 511 (2019).
- [4] STAR Collaboration (J. Adam *et al.*), *Phys. Rev. Lett.* **123**, 132301 (2019).
- [5] F. Becattini, L. Tinti, *Ann. Phys.* **325**, 1566 (2010).
- [6] F. Becattini *et al.*, *Ann. Phys.* **338**, 32 (2013).
- [7] D. Montenegro *et al.*, *Phys. Rev. D* **96**, 056012 (2017); *Addendum ibid.* **96**, 079901 (2017).
- [8] D. Montenegro, L. Tinti, G. Torrieri, *Phys. Rev. D* **96**, 076016 (2017).
- [9] F. Becattini, W. Florkowski, E. Speranza, *Phys. Lett. B* **789**, 419 (2019).
- [10] B. Boldizsár, M.I. Nagy, M. Csanád, *Universe* **5**, 101 (2019).
- [11] S.Y.F. Liu, Y. Sun, C.M. Ko, *Phys. Rev. Lett.* **125**, 062301 (2020).
- [12] W. Florkowski *et al.*, *Phys. Rev. C* **100**, 054907 (2019).
- [13] H.Z. Wu *et al.*, *Phys. Rev. Res.* **1**, 033058 (2019).
- [14] F. Becattini, G. Cao, E. Speranza, *Eur. Phys. J. C* **79**, 741 (2019).
- [15] J.j. Zhang *et al.*, *Phys. Rev. C* **100**, 064904 (2019).
- [16] K. Fukushima, S. Pu, [arXiv:2001.00359](https://arxiv.org/abs/2001.00359) [hep-ph].
- [17] W. Florkowski, A. Kumar, R. Ryblewski, *Acta Phys. Pol. B* **51**, 945 (2020).
- [18] S. Li, H.U. Yee, *Phys. Rev. D* **100**, 056022 (2019).
- [19] K. Hattori, Y. Hidaka, D.L. Yang, *Phys. Rev. D* **100**, 096011 (2019).
- [20] N. Weickgenannt *et al.*, *Phys. Rev. D* **100**, 056018 (2019).
- [21] N. Weickgenannt *et al.*, [arXiv:2005.01506](https://arxiv.org/abs/2005.01506) [hep-ph].
- [22] S. Shi, C. Gale, S. Jeon, [arXiv:2008.08618](https://arxiv.org/abs/2008.08618) [nucl-th].
- [23] K. Hattori *et al.*, *Phys. Lett. B* **795**, 100 (2019).
- [24] V.E. Ambrus, *J. High Energy Phys.* **2008**, 016 (2020).
- [25] X.L. Sheng, L. Oliva, Q. Wang, *Phys. Rev. D* **101**, 096005 (2020).
- [26] Y.B. Ivanov, V.D. Toneev, A.A. Soldatov, *Phys. Atom. Nucl.* **83**, 179 (2020).
- [27] Y. Xie, D. Wang, L.P. Csernai, *Eur. Phys. J. C* **80**, 39 (2020).
- [28] G.Y. Prokhorov, O.V. Teryaev, V.I. Zakharov, *Phys. Rev. D* **99**, 071901 (2019).
- [29] G.Y. Prokhorov, O.V. Teryaev, V.I. Zakharov, *J. High Energy Phys.* **1902**, 146 (2019).
- [30] G.Y. Prokhorov, O.V. Teryaev, V.I. Zakharov, *J. High Energy Phys.* **2020**, 137 (2020).
- [31] D.L. Yang, *Phys. Rev. D* **98**, 076019 (2018).
- [32] Y. Liu, X. Huang, *Nucl. Sci. Tech.* **31**, 56 (2020).
- [33] S. Tabatabaee, N. Sadooghi, *Phys. Rev. D* **101**, 076022 (2020).

- [34] S. Bhadury *et al.*, [arXiv:2002.03937](#) [hep-ph].
- [35] Y. Liu, K. Mameda, X. Huang, *Chin. Phys. C* **44**, 094101 (2020).
- [36] D. Yang, K. Hattori, Y. Hidaka, *J. High Energy Phys.* **2020**, 070 (2020).
- [37] X. Deng *et al.*, *Phys. Rev. C* **101**, 064908 (2020).
- [38] ExHIC-P Collaboration (H. Taya *et al.*), *Phys. Rev. C* **102**, 021901 (2020).
- [39] J.H. Gao *et al.*, [arXiv:2009.04803](#) [nucl-th].
- [40] S. Bhadury *et al.*, [arXiv:2008.10976](#) [nucl-th].
- [41] D. Montenegro, G. Torrieri, *Phys. Rev. D* **102**, 036007 (2020).
- [42] F. Becattini, M.A. Lisa, [arXiv:2003.03640](#) [nucl-ex].
- [43] L. Tinti, W. Florkowski, [arXiv:2007.04029](#) [nucl-th].
- [44] E. Speranza, N. Weickgenannt, [arXiv:2007.00138](#) [nucl-th].
- [45] J.H. Gao *et al.*, *Nucl. Sci. Tech.* **31**, 90 (2020).
- [46] F. Becattini, [arXiv:2004.04050](#) [hep-th].
- [47] F. Becattini *et al.*, *Phys. Rev. C* **95**, 054902 (2017).
- [48] I. Karpenko, F. Becattini, *Eur. Phys. J. C* **77**, 213 (2017).
- [49] H. Li *et al.*, *Phys. Rev. C* **96**, 054908 (2017).
- [50] Y. Xie, D. Wang, L.P. Csernai, *Phys. Rev. C* **95**, 031901 (2017).
- [51] F. Becattini, F. Piccinini, J. Rizzo, *Phys. Rev. C* **77**, 024906 (2008).
- [52] W. Florkowski *et al.*, *Phys. Rev. C* **97**, 041901 (2018).
- [53] W. Florkowski *et al.*, *Phys. Rev. D* **97**, 116017 (2018).
- [54] W. Florkowski, E. Speranza, F. Becattini, *Acta Phys. Pol. B* **49**, 1409 (2018).
- [55] W. Florkowski, R. Ryblewski, A. Kumar, *Prog. Part. Nucl. Phys.* **108**, 103709 (2019).
- [56] W. Florkowski, A. Kumar, R. Ryblewski, *Phys. Rev. C* **98**, 044906 (2018).
- [57] W. Florkowski *et al.*, *Phys. Rev. C* **99**, 044910 (2019).
- [58] W. Florkowski, M.P. Heller, M. Spalinski, *Rep. Prog. Phys.* **81**, 046001 (2018).
- [59] S.R. De Groot, W.A. Van Leeuwen, C.G. Van Weert, «Relativistic Kinetic Theory, Principles, Applications», *North-Holland*, Amsterdam 1980.
- [60] G. Denicol *et al.*, *Phys. Rev. D* **85**, 114047 (2012).
- [61] W. Florkowski, «Phenomenology of Ultra-Relativistic Heavy-Ion Collisions», *Singapore: World Scientific*, 2010.
- [62] J.D. Bjorken, *Phys. Rev. D* **27**, 140 (1983).
- [63] L. Rezzolla, O. Zanotti, «Relativistic Hydrodynamics», *Oxford University Press*, 2013.
- [64] E. Leader, «Spin in Particle Physics», *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. Vol. 15*, 2011 pp. 1–500.